

Measuring Mental Health Exasperation in the Era of Lockdowns

2:30PM EST | November 18th, 2021 | 2021 SyS Symposium

Moderator: Hussain Yusuf, M.D., M.P.H. – CDC/DDPHSS/CSELS/DHIS/PEB

Presenters:

Isabel Chung, MPH – Chicago Department of Public Health

Mwedusasa Mtenga, MPH – Florida Department of Health

Katie McDaniel, MPH – Florida Department of Health

Alan Mai, MPH – Florida Department of Health

Vasiliki Georgoulas-Sherry, PhD – Washington State Department of Health

Mary Megliola Franzen, MPH – Washington State Department of Health





Isabel Chung, MPH
Chicago Department of Public
Health






Trends in Emergency Department Utilization for Mental Health, Substance Use and Violent Victimization in Chicago, IL Before and at the Beginning of the COVID-19 Pandemic in 2020

Isabel Chung, MPH; Jocelyn Wilder, MPH; Wilnise Jasmin, MD, MBA, MPH



Emergency Department Visits in Chicago, 2019 - 2020

- The total number of Emergency Department (ED) visits in Chicago decreased by 21.5% from 2019 to 2020.
- Two pivotal dates in 2020:
 1. 3/21/20: the Illinois Stay-at-Home order and start of first COVID-19 surge.
 2. 10/5/20: start of the second and largest COVID-19 surge





Emergency Department Utilization for Behavioral Health, Chicago, IL, 2019 - 2020

Analytic Goal:

To examine how ED utilization was affected for those with acute behavioral health needs during the pandemic.

Methods:


- Selected 7 syndromic indicators from the National Syndromic Surveillance Program (NSSP) covering 3 behavioral health domains:
 - I. Mental Health
 - A. CDC Suicide Attempt, v1
 - B. SDC Disaster-related Mental Health v1
 - II. Violent Victimization
 - A. CDC Suspected Child Abuse & Neglect, v1
 - B. CDC Assault-related Firearm Injury, v1
 - C. CDC Intimate Partner Violence, v2
 - III. Substance Use
 - A. CDC Alcohol v1
 - B. CDC All Drugs v2

Methods, (continued):

- Indicators were examined both separately and collectively with a Behavioral Health composite score (total number of ED visits summed across all 7 NSSP indicators)
- Three distinct time periods:
 - A. Time 1 (T1): Before the IL stay-at-home order (reference period)
 - B. Time 2 (T2): After the IL stay-at-home order and up until the second surge
 - C. Time 3 (T3): During the second surge
- Metric Definitions:
 - A. Utilization rates: Number of ED visits for an indicator per 100,000
 - B. Percent change: Magnitude of change for each time period when compared to the same time period in 2019 expressed as a percent
 - C. Fold change: Magnitude of change between two time periods in 2020 expressed as a ratio



Emergency Department Utilization for Behavioral Health, Chicago, IL, 2020






Emergency Department Utilization for Behavioral Health, Chicago, IL, 2019 - 2020

Behavioral Health composite score:

- Total number of ED visits summed across all 7 Behavioral Health syndromes
- ED utilization for behavioral health increased by 19.3% in 2020 but decreased by 1.3% for all other types of ED visits.


Weekly ED Utilization Rates for Behavioral Health
Chicago, IL, 2019-2020






Emergency Department Utilization for Behavioral Health, Chicago, IL, 2019 - 2020

Weekly Rates for Behavioral Health ED Utilization, Chicago, IL, 2019-2020



Weekly Percent Change for Behavioral Health ED Utilization, Chicago, IL, 2019-2020



Emergency Department Utilization for Behavioral Health, Chicago, IL, 2019 – 2020

Behavioral Health Domain	Behavioral Health Syndrome	Total Number Visits in 2020 (N)	Annual mean weekly rate, 2020 (N per 100,000 visits)	Annual Percent Change in Mean Weekly Rates: 2019 to 2020	Average percent change for T1	Average percent change for T2	Average percent change for T3	Fold Change: T1 to T2	Fold Change: T1 to T3
OVERALL	Behavioral Health composite score of 7 Syndromes	77,042	7,670	+19.3%	14.2%	25.9%	+7.9%	1.8x (↑)	1.8x (↓)
Mental Health	CDC Suicide Attempt, v1	2,682	267	+21.7%	+4.4%	+31.3%	+17.0%	7.1x (↑)	3.9x (↑)
	SDC Disaster-related Mental Health v1	25,786	2,575	+14.7%	+2.2%	+20.1%	+13.9%	9.2x (↑)	6.4x (↑)
Violent Victimization	CDC Suspected Child Abuse & Neglect, v1	946	93	-0.7%	-10.0%	+10.2%	-14.8%	2.1x (↑)	1.5x (↓)
	CDC Assault-related Firearm Injury, v1	1,227	127	106.6%	+13.2%	+166.6%	+24.1%	12.7x (↑)	1.8x (↑)
	CDC Intimate Partner Violence, v2	313	31	14.0%	-2.1%	+29.1%	-6.9%	15.0x (↑)	3.3x (↓)
Substance Use	CDC Alcohol v1	28,208	2,799	6.7%	+6.5%	+10.7%	-2.6%	1.6x (↑)	1.4x (↓)
	CDC All Drugs v2	17,870	1,777	53.0%	+66.5%	+63.3%	+20.2%	0.95 (↓)	3.3x (↓)

(↑): Increase in fold change across two time periods;

(↓): Decrease in fold change across two time periods



Emergency Department Utilization for Behavioral Health, Chicago, IL, 2019 - 2020

Conclusions:

1. When compared to the first 3 months of 2020 (T1), ED utilization for behavioral health in Chicago:
 - Increased 1.8- fold during the 7 months following the IL stay-at-home order (T2) but then
 - Decreased by the same amount in last 3 months of 2020 (T3) during the second surge of COVID-19
2. Out of the three behavioral health domains, ED utilization rates for violent victimization increased the most in T2 followed by mental health.
3. In T3, ED utilization rates for violent victimization decreased the most followed by a slight decrease in mental health rates.
4. Large changes in substance use rates were not seen throughout 2020 although these ED visits accounted for the greatest proportion (60%) of all behavioral health visits.



Emergency Department Utilization for Behavioral Health, Chicago, IL, 2019 - 2020

Discussion:

1. Acknowledgment of data limitations:
 1. Data is cross-sectional. Therefore, causation cannot be inferred.
 2. ED utilization as a population indicator may not capture all contributing factors leading up to each visit.
 3. Chief complaint data used for NSSP-ESSENCE (Electronic Surveillance System for the Early Notification of Community-Based Epidemics) queries may impact sensitivity/specificity of indicators.
2. Multiple events related to the pandemic including the stay-at-home order, disruptions in healthcare and social service delivery systems, or major socioeconomic events (e.g. loss of jobs or loved ones) may have increased experiences of traumatic stress among many Chicagoans.
3. As seen in other large urban settings nationwide, events rooted in social injustice, such as the deaths of Breonna Taylor and George Floyd, and mounting political tensions around the 2020 elections may have added to increased social unrest and traumatic stress.
4. Acute changes in behavioral health outcomes may be manifestations of increased levels of traumatic stress.
5. Further analyses to stratify rates by demographics and geographics can be used to identify subpopulations who may be disproportionately experiencing worsening behavioral health. This information can also be used to bolster support systems cross-cutting multiple sectors (e.g. economic, behavioral health and social services) in ways that will promote coping and resilience.



Emergency Department Utilization for Behavioral Health, Chicago, IL, 2019 - 2020

Acknowledgements:

- Matthew Richards, LCSW, MDiv, Deputy Commissioner of Behavioral Health, Chicago Department of Public Health
- Nik Prachand, MPH, Director of Epidemiology, Chicago Department of Public Health
- Kathryn Jackson, MS, Biostatistician, Northwestern University
- Blair Turner, MPH, Epidemiologist II, Chicago Department of Public Health
- Livia Verklan McInnes, GIS Analyst, Chicago Department of Public Health

Data Source:

CDC National Syndromic Surveillance Program

Contact Information:

Isabel Chung, MPH, Epidemiologist III,
Chicago Department of Public Health
Isabel.chung@cityofchicago.org



Mwedusasa Mtenga, MPH
Florida Department of Health





Katie McDaniel, MPH & Alan Mai
Florida Department of Health



Department of Health

Increase in Emergency Department Visits due to Suspected Suicide Attempt Among Persons Ages 12-17 Years, Florida, January 2019–May 2021

**2021 Syndromic Surveillance Symposium
November 18, 2021**

Katie McDaniel, MPH

Senior Environmental Epidemiologist

Alan Mai, MPH

Senior Injury Epidemiologist



Background

- From 2010 to 2019, the suicide rate among Florida youth 12-17 years of age increased from 2.6 to 5.1 per 100,000.
- The number of Florida youth who think about or attempt suicide is even higher than the number of deaths by suicide.
 - ❖ According to the 2019 Florida Youth Risk Behavior Survey:
 - 7.9% of high school students reported a suicide attempt.
 - 11.8% made a plan on how to attempt suicide.
 - 15.6% seriously considered attempting suicide.
 - 33.7% experienced feelings of sadness or hopelessness almost every day.



Background, continued

- During 2020, the proportion of emergency department (ED) visits for suspected suicide attempts among youth 12-17 years of age spiked nationally and persisted into 2021.
- Reinforced the need for increased attention to the factors that exacerbate the risk for suicidal behavior among youth.
- To assess whether trends observed nationally occurred among Florida youth, Florida's syndromic surveillance data were analyzed.



Methods

- Syndromic surveillance data from Florida's Electronic Surveillance System for the Early Notification of Community-based Epidemics (ESSENCE-FL) were used.
- Examined patterns in ED visits for suspected suicide attempts among youth 12-17 years of age.
- Began with 5-year trend data.
- More focused analyses were then conducted on January 1, 2019 to May 13, 2021 time period.
 - ❖ Allowed for comparisons to national trends.




Methods, continued

- ESSENCE-FL captures data from approximately 99% of Florida's ED facilities.
- Suspected suicide attempts were identified using chief complaint terms and administrative discharge diagnosis codes.
 - ❖ Developed by the Centers for Disease Control and Prevention (CDC).



5-Year Trends

Monthly emergency department (ED) visits for suspected suicide attempts among Florida youth ages 12-17 years, January 2016-May 2021



**Count and percentage change of ED visits for suspected suicide attempts
among youth ages 12-17 years, Florida, January 2019-May 2021^a**

Group	Count			Percentage Change from 2019	
	2019	2020	2021	2020 ^b	2021 ^c
Total	3,043	4,339	2,905	42.59	122.43
Girls	2,298	3,435	2,330	49.48	134.88
Boys	745	902	575	21.07	83.12
White^d	2,044	2,988	1,987	46.18	133.22
Black^d	543	791	571	45.67	127.49
Other Race^d	594	792	541	22.81	70.94
Hispanic^e	456	560	347	33.33	114.68



^a Data are provisional and maybe subject to change.

^b Percentage change compared to January-December 2019.


^c Percentage change compared to January-May 2019.

^d Includes Hispanic and non-Hispanic.

^e Includes all races.




Results by Race/Ethnicity



Note: Includes Hispanic and non-Hispanic.




Results by Race/Ethnicity, continued



Note: Includes Hispanic and non-Hispanic.




Results by Race/Ethnicity, continued



Note: Includes Hispanic and non-Hispanic.



Results by Race/Ethnicity, continued



Note: Includes all races.



Limitations

- Data are provisional and may be subject to change.
- Not all facilities report to ESSENCE-FL.
- Results may be impacted by coding and reporting practices from each facility.
- Data may not represent true prevalence of suspected suicide attempts.



Limitations, continued

- Not able to determine single causal links between the many factors influencing youth suicide.
 - ❖ Adverse childhood experiences
 - ❖ Economic support
 - ❖ Access and delivery of evidence-based care
 - ❖ Coping skills and social connectedness



Discussion

- Findings presented align with national trends.
- Suicidal behaviors among youth are of particular concern.
 - ❖ Suicide attempts are a leading risk factor for death by suicide.
- Public health surveillance data are important to identify emergent factors.
 - ❖ Syndromic data allow for near real-time monitoring of trends.



Discussion, continued

- These data can inform suicide prevention initiatives.
 - ❖ Enable programs to quickly enhance approaches and implement evidence-based strategies to meet the needs of youth during times of emergency or disruption.
- Data to action
 - ❖ Build targeted initiatives for those most affected.
 - ❖ Partnerships and resources:
 - County Health Departments
 - Department of Children and Families
 - Department of Education
 - Community Partners



Contact Information

Katie McDaniel, MPH

Senior Environmental Epidemiologist

Katherine.McDaniel@flhealth.gov

Alan Mai, MPH

Senior Injury Epidemiologist

Alan.Mai@flhealth.gov





Vasiliki Georgoulas-Sherry, PhD &
Mary Megliola Franzen, MPH
Washington State Department of Health





WASHINGTON STATE DEPARTMENT OF HEALTH'S SYNDROMIC
SURVEILLANCE IN GUIDING THE BEHAVIORAL HEALTH
RESPONSE DURING THE PANDEMIC

Vasiliki Georgoulas-Sherry, PhD & Mary Franzen, MPH
Washington State Department of Health
Division of Emergency Preparedness, Resilience, and Response



Background

- The effect of the novel coronavirus (SARS-CoV-2) pandemic has produced significant health concerns, negatively impacting individuals in a multitude of ways, creating new hardships and barriers for individuals. Even if individuals are not physically affected by COVID-19, the implications for behavioral health remain.
- As the ongoing and constantly changing nature of the pandemic continues, the unique characteristics of this pandemic trend toward anxiety and loneliness as significant behavioral health outcomes.
- This subsequently has led to behavioral health problems being considered as critical and prevalent public health concerns, especially as the pandemic continues to considerably impact the community. The pandemic has presented uncommon behavioral health impacts like no other, and this requires different mitigation, planning, and response.

Methods

- Since April 2020, the Washington State Department of Health (WA DOH) has utilized the Rapid Health Information Network (RHINO) to conduct syndromic surveillance on five behavioral health syndrome queries that can be detected in emergency department settings:
 - psychological distress,
 - suicidal ideation,
 - suspected suicide attempts,
 - alcohol-related conditions, and
 - suspected all-drug overdoses
- through deidentified patient records.

Data Source & Definitions

The Department of Health collects syndromic surveillance data in near real-time from hospitals and clinics across Washington. The data are always subject to updates. Key data elements reported include patient demographic information, chief complaint, and coded diagnoses. This [data collection system](#) is the only source of emergency department (ED) data for Washington.

Visits of interest per 10,000 ED visits can help provide insights into:

1. behavioral health impacts since the implementation of the “Stay Home, Stay Healthy” order from March 23, 2020 (CDC Week 13),
2. seasonal shifts year-over-year,
3. new visit trends due to COVID-19 symptoms and diagnosis,
4. perceptions of disease transmission and risk, as well as,
5. the relative frequency of these indicators for 2019, 2020, and 2021.

Limitations:

Because the volume of visits across care settings varied widely during 2020 and to date in 2021, rates presented in this report may not reflect the true magnitude and direction of trends for behavioral health conditions and should be interpreted cautiously.

Furthermore, caution should be taken when evaluating data as fewer overall ED visits beginning in March 2020 (CDC Week 10-14) can skew results of data.

Furthermore, ED visits count for suicidal ideation, suspected suicide attempt, psychological distress, and suspected overdoses might show an increase in awareness in mental health experiences, thus taking a larger share of the total ED visits.

Suicidal ideation: Created by partners in the CDC's National Center for Injury Prevention to support states and jurisdictions to query visits related to suicidal ideation, or thoughts or plans of engaging in suicide-related behavior. Full details are available at <https://knowledgerepository.syndromicsurveillance.org/CDC-suicide-attempt-v1-syndromedefinitioncommittee>.

Suspected suicide attempt: Created by partners in the CDC's National Center for Injury Prevention to support states and jurisdictions to query visits related to a suicide attempt, or self-directed and potentially injurious behavior with any intent to die as a result of the behavior. Full details are available at <https://knowledgerepository.syndromicsurveillance.org/CDC-suicide-ideation-v1-syndromedefinitioncommittee>.


Psychological distress: This is a Syndrome Definition Committee community-developed syndrome definition for mental health conditions likely to increase in emergency department frequency during and after natural or human-caused disaster events. Full details are available at <https://knowledgerepository.syndromicsurveillance.org/disaster-related-mental-health-v1-syndromedefinitioncommittee>.

Suspected all drugs overdoses: This definition specifies overdoses for any drug, including heroin, opioid, and stimulants. It is indexed in the Electronic Surveillance System for the Early Notification of Community-Based Epidemics (ESSENCE) platform as CDC All Drug v1. Full details available at <https://knowledgerepository.syndromicsurveillance.org/cdc-all-drug-v1>.


Results

- Syndromic data evaluating psychological distress, suicidal ideation, suspected suicide attempts, alcohol-related conditions, and suspected all-drug overdoses from 2019 to 2021 has shown that visit volumes appear to be trending back to reflect levels more typical of care-seeking behavior before COVID-19, and therefore, rates may be converging with previous years' rates.
- While rate (calculated as per 10,000 ED visits) in all five behavioral health indicators peaked in the beginning months of 2021 and were typically higher than the rates in 2019 or 2020, the Summer months show decreases in these rates and remain lower than rates in 2019 or 2020.


Results: Psychological Distress




Results: Suicidal Ideation




Results: Suspected Suicide Attempts



Results: Alcohol-Related Conditions



Results: Suspected All-Drug Overdoses



Implications

- The utilization of syndromic surveillance for behavioral health can show with near real-time data the potential impacts of the COVID-19 pandemic on mental health and substance use issues which can:
 - support population-level strategic planning and intervention,
 - mitigate the behavioral health impacts of the COVID-19 pandemic,
 - impact forecasts used by health partners and state agencies for strategic planning and intervention, and
 - help public health partners such as behavioral health providers and response teams respond quickly to potential behavioral health outcomes of the COVID-19 pandemic by creating:
 - provider alerts,
 - social media messaging,
 - behavioral health reference guides, and
 - other health communications (youth-, older adults-, regional-, and statewide-focused reports on behavioral health).



To request this document in another format, call 1-800-525-0127. Deaf or hard of hearing customers, please call 711 (Washington Relay) or email civil.rights@doh.wa.gov.

National Syndromic Surveillance Program

Community of Practice

Resources that Advance the Science and
Practice of Syndromic Surveillance



2021

Syndromic Surveillance Symposium

Week of November 15, 2021



KAHUIINA
CONSULTING