

Chun-Zhen Hua ORCID iD: 0000-0001-9311-6229

Epidemiological features and viral shedding in children with SARS-CoV-2 infection

Chun-Zhen Hua^{1*}, Zi-Ping Miao^{2*}, Ji-Shan Zheng³, Qian Huang⁴, Qing-Feng Sun⁵, Hong-Ping Lu⁶, Fei-Fei Su⁷, Wei-Hong Wang⁸, Lie-Ping Huang⁹, Zhi-Wei Xu¹⁰, Le-Dan Ji¹¹, Hong-Ping Zhang¹², Xiao-Wei Yang¹³, Ming-Hui Li¹⁴, Yue-Yan Mao¹⁵, Man-Zhen Ying¹⁶, Sheng Ye¹⁷, Qiang Shu^{18#}, En-Fu Chen^{2#}, Jian-Feng Liang¹⁹, Wei Wang²⁰, Zhi-Min Chen²¹, Wei Li²², Jun-Fen Fu^{23#}

*Chun-Zhen Hua and Zi-Ping Miao contributed equally to this study.

Affiliations:

¹Department of Infectious Diseases, Children's Hospital, Zhejiang University School of Medicine, National Clinical Research Center for Child Health, Hangzhou 310003, Zhejiang, P.R. China; (huachunzhen@zju.edu.cn)

²Department of Infectious Diseases, Zhejiang Provincial Center for Disease Control and Prevention, Hangzhou 310003, Zhejiang, P.R. China; (zpmiao@cdc.zj.cn)

³Department of Infectious Diseases, Ningbo Women and Children's Hospital, Ningbo 315012, Zhejiang, P.R. China; (zjs222@163.com)

⁴Department of Infectious Diseases, Hangzhou Xixi Hospital, Hangzhou 310023, Zhejiang, P.R. China; (13575746522@163.com)

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1002/jmv.26180.

This article is protected by copyright. All rights reserved.

⁵Department of Infectious Diseases, Ruian People's Hospital, Wenzhou 325200, Zhejiang, P.R. China; (sunxue0806@126.com)

⁶Department of Pediatrics, Taizhou Hospital, Taizhou 317000, Zhejiang, P.R. China; (luhp@enzemed.com)

⁷Department of Infectious Diseases, Wenzhou Central Hospital, Wenzhou 325000, Zhejiang, P.R. China; (feifeisuzs@163.com)

⁸Department of Infectious Diseases, Huzhou Central Hospital, Huzhou 313000, Zhejiang, P.R. China; (hzwwh0606@163.com)

⁹Department of Pediatrics, Zhoushan Women and Children's Hospital, Zhoushan 316000, Zhejiang, P.R. China; (huanglp816@126.com)

¹⁰Department of Infectious Diseases, The Second Affiliated Hospital & Yuying Children's Hospital of Wenzhou Medical University, Wenzhou 325027, Zhejiang, P.R. China; (xuzw1971@sina.com)

¹¹Department of Pediatrics, Yueqing People's Hospital, Wenzhou 325600, Zhejiang, P.R. China; (jiledan@163.com)

¹²Department of Pediatrics, Wenzhou People's Hospital, Wenzhou 325000, Zhejiang, P.R. China; (zjzhp@126.com)

¹³Department of Pediatrics, The First People's Hospital of Wenling, Taizhou 317500, Zhejiang,

P.R. China;(yxw7007@126.com)

¹⁴Department of Infectious Diseases, Shaoxing People's Hospital, Shaoxing 312000, Zhejiang, P.R.

China;(lmhxx@163.com)

¹⁵Department of Pediatrics, The First People's Hospital of Yuhang, Hangzhou 311100, Zhejiang,

P.R. China;(hzmaoyy@163.com)

¹⁶Department of Infectious Diseases, Dongyang people's hospital, Jinhua 321100, Zhejiang, P.R.

China; (dm.0579@163.com)

¹⁷Department of Critical Care Medicine, The Children's Hospital, Zhejiang University School of

Medicine; National Clinical Research Center for Child Health, Hangzhou 310003, P.R

China.(E-mail: yeshengchina@zju.edu.cn)

¹⁸ Data information center, The Children's Hospital, Zhejiang University School of Medicine;

National Clinical Research Center for Child Health, Hangzhou 310003, P.R China.(E-mail:

shuqiang@zju.edu.cn)

¹⁹Statistics Unit, The Children's Hospital, Zhejiang University School of Medicine; National

Clinical Research Center for Child Health, Hangzhou 310003, P.R China. (E-mail:

liangjianfeng@zju.edu.cn)

²⁰Department of medical administration, The Children's Hospital, Zhejiang University School of Medicine; National Clinical Research Center for Child Health, Hangzhou 310003, P.R China.
(E-mail: wangw_hz930@zju.edu.cn)

²¹Department of respiratory, The Children's Hospital, Zhejiang University School of Medicine; National Clinical Research Center for Child Health, Hangzhou 310003, P.R China. (E-mail: chenzhimin6@163.com)

²² Department of laboratory center, The Children's Hospital, Zhejiang University School of Medicine; National Clinical Research Center for Child Health, Hangzhou 310003, P.R China. (E-mail: yuki3456@zju.edu.cn)

²³Data information center, The Children's Hospital, Zhejiang University School of Medicine; National Clinical Research Center for Child Health, Hangzhou 310003, P.R China. (E-mail: fjf68@zju.edu.cn)

Running Title: Viral shedding in children with SARS-CoV-2 infection

Ethics approval and consent to participate: The study was approved by both the institutional ethics board of Zhejiang Provincial CDC (T-043-R) and ethics commission of all designated hospitals for recruiting COVID-19 patients. Oral consents were obtained from the guardians of children.

Conflicts of Interest: The authors have no conflicts of interest to declare.

This article is protected by copyright. All rights reserved.

Address for correspondence:

Qiang Shu, Data Information Center, Children's Hospital, Zhejiang University School of Medicine; National Clinical research Center for Child Health, Hangzhou 310003, P.R China.

Phone: 0086-0571-86670006; Fax: 0086-571-87033296. E-mail: shuqiang@zju.edu.cn

En-Fu Chen, Department of Infectious Disease, Zhejiang Provincial Center for Disease Control and Prevention, Hangzhou 310003 Zhejiang, P.R. China. Phone: 0086-571-87115130. Email: enfchen@cdc.zj.cn

Jun-Fen Fu, Data information center, Children's Hospital, Zhejiang University School of Medicine; National Clinical research Center for Child Health, Hangzhou 310003, P.R China.

Phone: 0086-0571-88873013; Fax: 0086-571-87033296. E-mail: fjf68@zju.edu.cn

Evidence before this study

Corona virus disease 2019 (COVID-19) is a highly contagious disease and has caused pandemics all over the world. Children had milder clinical symptoms, better clinical outcome, and more common fecal viral excretion. However, their susceptibility for SARS-CoV-2 infection is a matter of debate, and if antiviral therapy affects viral shedding outcomes is unknown.

Added value of this study

We investigated all contacts in the families with SARS-CoV-2 infected members, and analyzed the epidemical aspects in children in Zhejiang Province, China. We found that incidence of infection in children contacts was 13.2% (43/325, 95%CI 9.5-16.9), which was significant lower than that in

adult contacts (21.2%, 108/ 510, 95%CI 17.6-24.7) in the same families. The mean incubation period for children was 9.1 days. Fecal SARS-CoV-2 RNA detection was positive in 91.4% (32/35) cases and some children had viral excretion time exceeded 70 days. Viral clearance time was not different among the groups treated with different antiviral regimens. No subsequent infection was observed in family contacts of fecal-viral-excretion children.

Implications of all the available evidence

Children have lower susceptibility for SARS-CoV-2 infection, longer incubation and fecal viral excretion time. Positive results of fecal SARS-CoV-2 RNA detection are not used as indication for hospitalization or quarantine. No effect on fecal viral clearance was observed when treated with antiviral regimens in the study.

Abstract

Background A pandemic of SARS-CoV-2 infection broke out all over the world, however, epidemiological data and viral shedding in pediatric patients are limited.

Methods We conducted a retrospective, multi-center study, and followed up with all children from the families with SARS-CoV-2 infected members in Zhejiang Province, China. All infections were confirmed by testing the SARS-CoV-2 RNA with RT-PCR method, and epidemiological data between children and adults in the same families was compared. Effect of antiviral therapy was evaluated observationally and fecal viral excretion times among groups with different antiviral regimens were compared with Kaplan Meier plot.

Results By February 29, 2020, 1298 cases from 883 families were confirmed with SARS-CoV-2 infection and 314 of which were families with children. Incidence of infection in child close contacts was significantly lower than that in adult contacts (13.2% vs 21.2%). The mean age of 43 pediatric cases was 8.2 years and mean incubation period was 9.1 days. Forty (93.0%) were family clustering. Thirty-three children had COVID-19 (20 pneumonia) with mild symptoms and 10 were asymptomatic. Fecal SARS-CoV-2 RNA detection was positive in 91.4% (32/35) cases and some children had viral excretion time over 70 days. Viral clearance time was not different among the groups treated with different antiviral regimens. No subsequent infection was observed in family contacts of fecal-viral-excreting children.

Conclusion Children have lower susceptibility of SARS-CoV-2 infection, longer incubation and fecal viral excretion time. Positive results of fecal SARS-CoV-2 RNA detection were not used as indication for hospitalization or quarantine.

Funding Zhejiang University Special Scientific Research fund; Ruian Science and Technology Bureau.

Keywords: SARS-CoV-2; COVID-19; Epidemiological characteristics; Children; Viral excretion; Antiviral therapy; Viral clearance

Text

Recently, a pandemic of corona virus disease 2019 (COVID-19) broke out all over the world. By June 10, 2020, more than 7 million cases infected with SARS-CoV-2 were reported worldwide ¹.

Reports on COVID-19 exploded and most of them were focused on adults²⁻⁶. Research involving pediatric cases with SARS-CoV-2 infection was rare^{5,7}. A recent review of 72,314 cases by the Chinese Center for Disease Control and Prevention (CDC) showed that less than 2% of the patients were younger than 19 years⁸. Dong et al analyzed the epidemiology of COVID-19 among children in China, reported that up to February 8, 2020, the total confirmed cases (≤ 16 years) was 574 national wide⁹. The low incidence of pediatric COVID-19 has perplexed clinicians, epidemiologists and scientist¹⁰. Most of the researchers believe that children are susceptible for SARS-CoV-2 infection^{7,9,10}, the absence of pediatric patients with COVID-19 should be attributed to low exposure to COVID-19 patients⁴. Xu et al⁷ screened 745 children and 3,174 adults by nasopharyngeal swab real-time PCR with reverse transcription (RT-PCR) for SARS-CoV-2 infection, found that confirmed cases were much fewer in children after exposure to COVID-19 patients than that in adults. Thus, if children have lower susceptibility for SARS-CoV-2 infection becomes a matter of debate. During the progress of COVID-19 outbreak, Zhejiang was one of the hard-hit provinces⁹. The number of confirmed cases rose rapidly along with the return of merchants and college students who lived in Wuhan before the Spring Festival, which made it possible to perform a family clustering study on the susceptibility to SARS-CoV-2 infection in children. During the preparation of our manuscript, a report on clinical and epidemiological features of 36 children (≤ 16 years) with COVID-19 in Zhejiang, China, the same region as our study, was published on the journal of Lancet Infectious Disease¹¹. The data showed that children with COVID-19 had mild or asymptomatic disease accompanied by pneumonia in about half the cases. Certainly, there was some overlap of patients' population in both studies. Our study had

enrolled all pediatric cases in Zhejiang Province, which were about 30% more cases (≤ 14 years) than the previous study¹¹. In addition to the similar clinical characteristic as described by Qiu et al., we have added analysis on the susceptibility of SARS-CoV-2 infection in children contacts and adult contacts in the same families, investigation on viral shedding and evaluation the effectiveness on viral clearance of different antiviral treatments by long time of follow-up. Our study addressed some of the most essential and un-answered questions that were not reported by Qiu et al.¹¹. Thus, the objective of the current study was to describe the epidemiological and viral clearance aspects in all children from the families with SARS-CoV-2 infected members in Zhejiang province.

Methods

Study design and participants

A retrospective, multi-center study including all pediatric cases (≤ 14 years) with SARS-CoV-2 infection, accompanied by follow up, was designed in Zhejiang province, China. There was no selection of any sort on cases. Demographic information and epidemiological data of all pediatric cases were exacted from the electronic master database (updated daily) established by the Zhejiang Provincial CDC. All cases were confirmed based on positive results of a real-time reverse transcription PCR (RT-PCR) assay of SARS-CoV-2 RNA from respiratory specimens¹² and all cases with confirmed infection should be hospitalized at local designated hospitals according to the national policy. The epidemiological data was obtained from CDC records by Miao ZP (who was both health authorities from CDC and researchers in the study). Clinical data was obtained from the patients' medical records by the attending (who were researchers in the study too). Clinical

outcomes were followed up (till now) after the children were discharged (outpatients follow up by attending and telephone follow up by Hua CZ). Children's epidemiological information was confirmed again with children's guardians by direct telephone communication during the follow-up process. The incubation period was from the initial exposure to the illness onset day. Initial exposure was defined as the day when children was exposed to the confirmed patients for those who occasionally visited; or previous 3 days when the first patient had illness onset in the family for those who lived together¹³. All data were cross-checked by two researchers (Hua CZ and Miao ZP).

This study was approved by both the institutional ethics board of Zhejiang Provincial CDC (T-043-R) and ethics commission of all designated hospitals for recruiting COVID-19 patients. Individual privacy was protected during the study. Oral consents were obtained from the guardians of the children.

Statistical analysis

SPSS software 20.0 (IMM, US) was used in the study. Age and time variables were described as mean (SD) if they were normally distributed and compared with *t* test, or expressed as median (IQR) if they were not normally distributed and days in hospital among groups with different antiviral therapeutic regimens were compared with the Kruskal-Wallis test. Categorical variables were described as number (%) and the prevalence of SARS-CoV-2 infection was compared by χ^2 test between children group and adult group. Fisher's exact test was used in comparing the positive rates of RT-PCR results in feces among groups with different antiviral regimes. A two-sided α of

less than 0.05 was considered statistically significant. Epidemic trend analysis was conducted with R software version 3.5.3 (R Foundation for Statistical Computing)

Role of the funding source

The funder of the study had no role in study design, data collection, data analysis, data interpretation, or writing of the report. The corresponding authors had full access to all the data in the study and had final responsibility for the decision to submit for publication.

Results

1. Epidemiological features of SARS-CoV-2 infection in children By February 29, 2020, 1298 cases infected with SARS-CoV-2 were confirmed in Zhejiang province, China. The patients aged from 3m20d to 96 years old, 43 of which (3.3%, 43/1298) were children (≤ 14 years). The epidemic trend of SARS-CoV-2 infection in children and adults in Zhejiang Province was shown in Fig 1.

All these 1298 cases were from 883 families excluding the prisoners, elderly people in gerocomium and students on campus, 714 families (80.9%) of which were successfully followed up and 314 families had 417 children living together. Two of which were neonates born by pregnant women with COVID-19. Specimens evaluated from umbilical cord blood, neonatal blood, stool, and nasopharynx specimen, all were negative for SARS-CoV-2 RNA by RT-PCR.

The physical conditions of the children contacts and adult contacts from the 314 families with SARS-CoV-2 infected members in Zhejiang Province are shown in Table 1. All SARS-CoV-2 infection cases in the present study were confirmed by real-time reverse transcription PCR assay

This article is protected by copyright. All rights reserved.

with respiratory specimen. Incidence of SARS-CoV-2 infection in children contacts (13.2%, 43/325, 95%CI 9.5-16.9) was lower than that in those adults contacts (21.2%, 108/510, 95%CI 17.6-24.7), and the difference was statistically significant ($\chi^2=8.46$, $P=0.004$).

The 43 cases who were not more than 14 years were analyzed, and 26 (60.5%, 26/43) were male.

The children's ages were from 3m20d to 14 y with a mean of 8.16 years (SD: 4.07). Forty children (93.0%) belong to family clustering cases and had long-term exposure (lived together with the family members with COVID-19 or had lived in epidemic areas, ie, Wuhan, or other cities in Hubei province). Three (7.0%, 3/43) children had short-term exposure to COVID-19 patients directly (occasional visits or in public transportations). The incubation periods were ranging from 4 days to 21 days, with a mean of 9.1 (SD: 3.7) days. Three patients (9.4%, 3/32) had incubation stages for more than 14 days. Twenty children (46.5%, 20/43) were imported cases: 13 children lived in Hubei province (12 in Wuhan city); 6 children were on vacation for 5-10 days in Hubei province (3 in Wuhan); and 1 child lived in Henan province and came to Hangzhou for medical advice after his father, who is a doctor and treated suspected COVID-19 patients from Wuhan in his private clinic in countryside, thought they might have been infected by SARS-CoV-2. Feces were tested for SARS-CoV-2 RNA in 35 children by RT-PCR more than one time, and positive results were found in 32 (91.4%, 32/35) children.

2. Clinical characteristics of pediatric COVID-19 Signs and symptoms in 43 children with SARS-CoV-2 infection are shown in Table 2. Two patients had vomiting, diarrhea, abdominal pain after receiving oral lopinavir/ritonavir and arbidol, and atomization inhalation with interferon- α 2b

in combination, and their gastrointestinal symptoms disappeared after lopinavir/ritonavir being removed or replaced with darunavir/cobicistat tablets. Liver function abnormality occurred in 3 patients after they received oral lopinavir/ritonavir and arbidol, and atomization inhalation with interferon- α 2b (lopinavir/ritonavir was replaced with darunavir/cobicistatin one patient because of gastrointestinal symptoms). Thirty-six (83.7%, 36/43) children received chest computed tomography (CT) examination 1-8 times when they were in hospital or during the outpatient follow-up. Twenty children had imaging evidence for pneumonia, and 9 had patchy shadow mainly in the peripheral lung fields. Three (15.0%, 3/20) children were asymptomatic and did not have imaging finding during the first week in hospital, but had patchy shadow or unilateral ground-glass opacity by chest CT when rechecked before being discharged (3 patients) or after being discharged (1 patient). Unilateral pneumonia were found by CT in 3 children whose chest X-rays were normal. The history of illness and the time when imaging change occurred in 20 patients with SARS-CoV-2 pneumonia are shown in Table 3.

Treatment and outcome

All children were treated in isolation, and stayed in designated hospitals for recruiting COVID-19 patients. Only seven (16.3%, 7/43) of the children were hospitalized in the infectious disease ward in the children's hospital or in pediatric ward in general hospitals. Thirty-eight children (88.4%, 38/43) received antiviral treatment, including 21 with monotherapy, 17 with two or more than two antiviral drugs (Table 4), The duration of antiviral treatment ranged from 5 to 31 days with a mean as 14.9 (SD: 7.4) days. Five patients were not prescribed with any antiviral drugs. No patient was

administered with corticosteroid or intravenous immunoglobulin. Oral Chinese medicine was used in 20 cases for 2-35 days. By March 6, 2020, all of the 43 children were discharged with favorable outcome. Their hospital stay ranged from 3 to 32 days with a mean of 20.2 (SD: 7.9) days. Criteria for discharge were based on viral clearance in respiratory samples from upper respiratory tract, improvement of clinical symptoms and chest radiographic evidence. Eighteen children (51.4%, 18/35) had positive results of SARS-CoV-2 RT-PCR in feces when they were discharged. Alanine aminotransferase increased from normal to 106 U/L in one child after he was discharged and had received more than 30 days of oral Chinese medicine (He had had liver function abnormality after receiving antiviral treatment in hospital and had recovered when being discharged). All children were kept in quarantine for another 2 weeks, which included arrangement at resorts by the government for 35 and home quarantine for the other eight patients.

By the end of April 20, 41 were followed up by telephone for at least twice, and one patient still had positive fecal RT-PCR results. On day 7 and day 14 after being discharged, the results of fecal RT-PCR switched to negative in 17.6% (3/17) and 33.3% (6/18) of the children, respectively. Positive fecal viral excretions were confirmed by RT-PCR for more than 70 days in one child since illness onset (As of April 20, 2020, follow up are going on). Fig2 shows the clearance curves of SARS-cov-2 in feces in cases treated with different antiviral regimens. None of their family contacts developed new infection during the quarantine periods. Imaging evidence for pneumonia was found in 1 child (he was asymptomatic at that time) when he was re-checked by CT 7 days after being discharged.

Discussion

Currently, a few studies on pediatric SARS-CoV-2 infection were reported accompanied by the global pandemic of COVID-19 in the world, though the numbers of cases included in these studies were usually small because of the low incidence in children. Xu et al ⁷ studied 745 children and 3,174 adults who had either close contact with diagnosed patients or had family clusters and found that positive rate in adults was 2.7-fold higher than that in children. Similarly, in the present study, we investigated all of the contacts in the same families and found that the incidence of SARS-CoV-2 infection in children contacts was 13.2%, which was much lower than that in adults contacts and was in accordance with the finding by Xu et al ⁷ and Lu et al ¹⁴, and indicated that children are less susceptible to SARS-CoV-2 infection. SARS-CoV-2 uses the cell receptor Angiotensin Converting Enzyme 2 (ACE2) and Transmembrane Serine Protease 2 (TMPRSS2) for infecting cells. Children were found to have significantly lower expression of SARS-CoV-2 receptors-ACE2 and TMPRSS2 in the upper and lower airways than that of adults ¹⁵, and the reduced airway tissue expression of ACE2 and TMPRSS2 may be the reason why children had lower risk of infection ^{16, 17}. With the implementation of policy to restrict the movement of people, the epidemic trend of COVID-19 in children was controlled much better than that in adults, and no new cases were confirmed since February 10, 2020, which may be due to the low susceptibility, as well as the reduced exposure. Most of the children were at school ages and were in family clusters. The epidemic peak in children was two weeks later than that in adults, which might be associated with the longer incubation stage ^{7, 18}. The median incubation period in adults usually was 3.0-6.4 days ^{2-6, 19}. Conversely, it was much longer in children ⁷. The median incubation period was 9.1

days with a range as 4~21 days in the present study, which could partly explain why the number of pediatric cases was peaked two weeks after that of adult cases. A long incubation period in children indicated that quarantine period for children who had contacted confirmed patients should be longer.

Similar with the findings in previous studies^{5,7,20}, symptoms in most of the pediatric patients with COVID-19 were mild, or even none. Most of patients did not seek medical care, and they were diagnosed because of their exposure history. Fever and cough were the most common symptoms, which were milder or even transient. Headaches, fatigue and chest tightness, which were common symptoms in adults, were rare in children in our study. All of the patients recovered soon. Leukopenia and lymphocytopenia were not common and most of the children had normal C-reactive protein, which indicated the inflammatory response was mild.

Although children with SARS-CoV-2 infection had mild symptoms or even were asymptomatic, chest CT was still widely used. Most of the children received chest CT more than one time.

Accordingly, more pneumonia, including “asymptomatic pneumonia”, was confirmed based on abnormalities in chest CT imaging. Image manifestation was found upon admission in most adult patients, and the typical findings of chest CT images were bilateral pneumonia, multiple lobular ground-glass opacity and subsegmental areas of consolidation^{2-6,17,21}. Twenty percent of the children with pneumonia in our study were confirmed at the second weeks in hospital, or even after being discharged. The most common findings were unilateral pneumonia, isolated ground-glass opacity. Transient fever or respiratory symptoms were not present in half of the

patients when pneumonic images were found by chest CT or X-ray, indicating that change in pulmonary image might be later than the appearance of clinical symptom. The possible progress of SARS-CoV-2, an emerging virus, in children was not clear, which might lead to the over examination by chest CT. Chest CT helps to find more cases with COVID-19 from children with mild symptoms or even without any symptoms²¹. Even so, when we look back on these children with SARS-CoV-2 infection, we question the necessity of chest CT in most cases with mild symptoms or even without any symptoms.

As previously mentioned, COVID-19 is an emerging disease and little was known about it in children when it broke out. Thus, an effective treatment has not been established. Symptoms in adults were severer, and antiviral drugs, such as oral lopinavir/ritonavir, oral arbidol and atomization inhalation with interferon, were widely used in adults in Zhejiang province.

Accordingly, pediatric cases, including asymptomatic children, most of which were hospitalized in infectious disease ward in general hospitals, were given antiviral drugs. It was difficult to analyze the effectiveness of antiviral therapy in children by the duration it needed to improve clinical symptoms, because their symptoms were mild and transient. As an alternative, we evaluated the outcome of treatment by analyzing the persistent respiratory and fecal viral shedding, and found that there was no difference among the groups received no antiviral therapy, or received one, two, three or more than three antiviral drugs. Furthermore, gastrointestinal reactions and liver function abnormality occurred in some children after receiving ≥ 3 antiviral drugs. Given that no antiviral treatment for coronavirus infection has been proven to be effective, we advise that, antiviral treatment might not be necessary for COVID-19 children without severe symptoms²². The best

way to treat pediatric mild SARS-CoV-2 infection might be doing nothing except close monitoring and isolation.

In this study, 18 children were discharged with positive results of fecal SARS-CoV-2 RNA detection. All of them were quarantined for another 2-4 weeks at resorts or at home. No new patient was found among their family members with close contact. According to previous studies, no evidence of fecal-oral transmission has been found for respiratory virus, the human coronavirus SARS or MERS. Similarly, there is no conclusive evidence that SARS-CoV-2 can cause illness by ingesting contaminated food or water to date. Therefore, we believe that a negative result of SARS-CoV-2 RNA detection in feces is not necessary for patients, whose clinical signs and symptoms had disappeared, to be discharged or released from isolation²³. The level of expression of the viral receptor (ACE2) and TMPRSS2, especially in the nasal tissue, may be critical for the ability of the virus to transmit and replicate. The reason for prolonged viral shedding in feces in children is unknown yet. As the receptor of SARS-CoV-2, ACE2, was abundantly expressed in gastric, duodenal and rectal epithelia in COVID-19 patients²⁴, which may lead to virus internalization and accumulation in these organs. It might explain the prolonged fecal virus shedding but needs to be further investigated.

In the present study, there were two neonates delivered by mothers with COVID-19, fortunately, neither clinical nor laboratory evidence for SARS-CoV-2 infection was confirmed in these babies. Till now, no case with vertical transmission was identified among pregnant women infected with SARS-CoV or MERS-CoV²⁵. However, during the epidemic of COVID-19 in China, a neonate,

delivered by a SARS-CoV-2 infected pregnant woman in Wuhan, was confirmed with COVID-19 at the age of 30-hour-old. The baby had shortness of breath together with abnormal chest imaging and liver function abnormalities²⁶. The possible route of SARS-CoV-2 transmission between the mother and neonate was not conclusive²⁵.

Our study has some limitations. First, the size of the cases was small, and all of the 43 children with SARS-CoV-2 infection were hospitalized in 15 local designated hospitals according to the principle of localization management, the program for pediatric cases was not run across hospitals. The diagnosis was confirmed with respiratory tract specimens and paired rectal swabs or feces specimens were not obtained for all children. The interval time of SARS-CoV-2 RNA detection was also variable. Therefore the durations of viral excretion through the gastrointestinal and respiratory tracts were not accurate. Second, not all children in the family with infected members were checked by real-time RT-PCR for SARS-CoV-2 RNA. Adolescent was not separated from the adult group because of the small sample size. Finally, the finding that antiviral therapy did not affect viral shedding outcomes was evaluated observationally, further study based on clinical trials of antiviral therapies is needed.

In conclusion, we found that susceptibility of SARS-CoV-2 infection in children was lower, and the incubation periods were longer than that in adults. The clinical symptoms in pediatric cases were mild. Chest CT or X-ray is helpful for diagnosing SARS-CoV-2 pneumonia; however, the necessity is questionable because most patients had mild symptoms and would be self-healing. Benefit of antiviral treatment on improving the clinical signs or shortening the duration of potential

viral excretion was not conclusive and further study is needed. At last, positive results of fecal SARS-CoV-2RNA detection should not be used as indication for hospitalization or quarantine.

Contributors

HCZ and MZP contributed to writing of the paper and took responsibility for the integrity of the data and the accuracy of the data analysis. ZJS, HQ, SQF, LHP, SFF, WWH, HLP, XZW, JLD, ZHP, YXW, LMH, MYY, YMZ and YS contributed to treatment of the patients, acquisition and interpretation of clinical data. LJF contributed to analysis of data. WW and CZM contributed to critical revision of the paper. SQ, CEF and FJF conceived the idea for and designed the study and had full access to all data in the study. All authors reviewed and approved the final version.

Declaration of interests

All authors declare no competing interests.

Data sharing

All authors declare that we this submission could be shared directly with the WHO.

The data that support the findings of this study are available from the corresponding author upon reasonable request. Participant data without names and identifiers will be made available after approval from the corresponding authors and Zhejiang Provincial CDC.

Acknowledgments

This work was supported by Zhejiang University Special Scientific Research fund for COVID-19 Prevention and Control, and funding from Ruian Science and Technology Bureau

This article is protected by copyright. All rights reserved.

(MS2020023). We acknowledge all health-care workers involved in the diagnosis and treatment of patients in Zhejiang Province. We thank Cai-Mei Fang, Medical Education Center, Health Commission of Zhejiang Province; Yi Jiang, Wenzhou People's Hospital; Qiao-Er Luo, Ningbo Women and Children's Hospital; Ming-Bin Xu, Changnan Chinese Medicine Hospital; Hui-Li Li, Changnan People's Hospital; Zhe Zhang, Ningbo Yinzhou People's Hospital; Shun-Zhi, Chen, Shaoxing People's Hospital; Yan-Hua Zheng, Zhoushan Women and Children's Hospital; Xiao-Qiang Zhang, The First People's Hospital of Yuhang; for their help in data collection. We thank Ying-Jie Lu, Harvard Medical School and Boston Children's Hospital, for his critical reading of the manuscript.

References

1. WHO. Coronavirus disease (COVID-19) Situation Report - 142. https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200610-covid-19-sitrep-142.pdf?sfvrsn=180898cd_2.
2. Chan JF, Yuan S, Kok KH, et al. A familial cluster of pneumonia associated with the 2019 novel coronavirus indicating person-to-person transmission: a study of a family cluster. *Lancet*. 2020; 395:514-23. doi: 10.1016/S0140-6736(20)30154-9.
3. Zhu N, Zhang D, Wang W, et al. Coronavirus Investigating, and Research Team. A novel coronavirus from patients with pneumonia in China, 2019. *N Engl J Med*. 2020; 382:727-33. doi: 10.1056/NEJMoa2001017.
4. Huang C, Wang Y, Li X, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet*. 2020; 395:497-506. doi: 10.1016/S0140-6736(20)30183-5.

5. Guan WJ, Ni ZY, Hu Y, et al. Clinical characteristics of 2019 novel coronavirus infection in China. *N Engl J Med*. 2020; 382:1708-20. doi: 10.1056/NEJMoa2002032.
6. Chen N, Zhou M, Dong X, et al. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet*. 2020; 395: 507-13. doi: 10.1016/S0140-6736(20)30211-7.
7. Xu Y, Li X, Zhu B, et al. Characteristics of pediatric SARS-CoV-2 infection and potential evidence for persistent fecal viral shedding. *Nat Med*. 2020;26:502-5. doi: 10.1038/s41591-020-0817-4.
8. Wu Z, McGoogan JM. Characteristics of and important lessons from the coronavirus disease 2019 (COVID-19) outbreak in China: summary of a report of 72 314 cases from the Chinese Center for Disease Control and Prevention. *JAMA*. 2020; 323: 1239-42. doi: 10.1001/jama.2020.2648.
9. Dong Y, Mo X, Hu Y, et al. Epidemiology of COVID-19 among children in China. *Pediatrics*. 2020;145(6):e20200702. doi: 10.1542/peds.2020-0702.
10. Kelvin AA, Halperin S. COVID-19 in children: the link in the transmission chain. *Lancet Infect Dis*. 2020;20(6):633-4. doi: 10.1016/S1473-3099(20)30236-X.
11. Qiu H, Wu J, Hong L, Luo Y, Song Q, Chen D. Clinical and epidemiological features of 36 children with coronavirus disease 2019 (COVID-19) in Zhejiang, China: an observational cohort study. *Lancet Infect Dis*. 2020; 20(6):689-96. doi: 10.1016/S1473-3099(20)30198-5.
12. WHO. Laboratory testing for 2019 novel coronavirus (2019-nCoV) in suspected human cases[EB/OL]. (2020-01-14). https://www.who.int/docs/default-source/coronaviruse/20200114-interim-laboratory-guidance-version.pdf?sfvrsn=6967c39b_4&download=true.
13. National Health Commission and National Administrative Office of Chinese Tradition Medicine. National Recommendations for Diagnosis and Treatment of pneumonia caused by 2019-nCoV (the 7th edition). <http://www.gov.cn/zhengce/zhengceku/2020-03/04/5486705/files/ae61004f930d47598711a0d4cbf874a9.pdf>.

14. Lu X, Zhang L, Du H, et al. SARS-CoV-2 Infection in Children. *N Engl J Med*. 2020;382:1663-5. doi: 10.1056/NEJMc2005073.
15. Sharif-Askari NS, Sharif-Askari FS, Alabed M, et al. Expression of SARS-CoV-2 receptor, ACE2, and TMPRSS2 in the lung airways is lower in children compared to adults and increases due to smoking and COPD. *Molecular Therapy: Methods & Clinical Development*, <https://doi.org/10.1016/j.omtm.2020.05.013>
16. Li W, Zhang B, Lu J, et al. The characteristics of household transmission of COVID-19. *Clin Infect Dis*. 2020 Apr 17;ciaa450. doi: 10.1093/cid/ciaa450. [Online ahead of print]
17. Zhang J, Litvinova M, Liang Y, et al. Changes in contact patterns shape the dynamics of the COVID-19 outbreak in China. *Science*. 2020 Apr 29;eabb8001. doi: 10.1126/science.abb8001. [Online ahead of print]
18. Cai J, Xu J, Lin D, et al. A Case Series of children with 2019 novel coronavirus infection: clinical and epidemiological features. *Clin Infect Dis*. 2020 Feb 28. doi: 10.1093/cid/ciaa198. [Epub ahead of print].
19. Backer JA, Klinkenberg D, Wallinga J. Incubation period of 2019 novel coronavirus (2019-nCoV) infections among travellers from Wuhan, China, 20-28 January 2020. *Euro Surveill*. 2020;25(5):2000062. doi: 10.2807/1560-7917.ES.2020.25.5.2000062.
20. Ai T, Yang Z, Hou H, et al. Correlation of Chest CT and RT-PCR Testing in Coronavirus Disease 2019 (COVID-19) in China: A Report of 1014 Cases. *Radiology*. 2020;Feb 26:200642. doi: 10.1148/radiol.2020200642. [Epub ahead of print].
21. Chua F, Armstrong-James D, Desai SR, et al. The role of CT in case ascertainment and management of COVID-19 pneumonia in the UK: insights from high-incidence regions. *Lancet Respir Med*. 2020;8:438-40. doi: 10.1016/S2213-2600(20)30132-6.

22.Chen ZM, Fu JF, Shu Q, et al. Diagnosis and treatment recommendations for pediatric respiratory infection caused by the 2019 novel coronavirus. *World J Pediatr.* 2020; Feb 5:1-7. doi: 10.1007/s12519-020-00345-5. [Epub ahead of print].

23.Steinbuch Y. Chinese baby tests positive for coronavirus 30 hours after birth. Available online: <https://nypost.com/2020/02/05/chinese-baby-tests-positive-for-coronavirus-30-hours-after-birth/> (accessed 9 February 2020).

24.Xiao F, Tang M, Zheng X, et al. Evidence for gastrointestinal infection of SARS - CoV - 2. *Gastroenterology.* 2020; 158(6):1831-3. doi:10.1053/j.gastro. 2020.02.055.

25.Wu Y, Guo C, Tang L, et al. Prolonged presence of SARS-CoV-2 viral RNA in faecal samples. *Lancet Gastroenterol Hepatol.* 2020;5:434-5. doi: 10.1016/S2468-1253(20)30083-2.

26.David A. Schwartz, Ashley L. Graham. Potential maternal and infant outcomes from coronavirus 2019-nCoV (SARS-CoV-2) infecting pregnant women: lessons from SARS, MERS, and other human coronavirus infections. *Viruses* 2020;12(2):194. doi: 10.3390/v12020194.

Legends

Fig 1 Epidemic trend of SARS-CoV-2 infection in children and adults in Zhejiang province




Fig 2 The clearance curves of SARS-cov-2 in feces in cases treated with different antiviral regiments




Table 1 The prevalence of infection in the children contacts and adult contacts from the 314

families with SARS-CoV-2 infected members in Zhejiang Province, China

Physical condition	Total	Children	Adults	Statistic values
	n	n (%)	n (%)	
SARS-CoV-2 infection	151	43 (10.3%)*	108 (17.1%)	$\chi^2=9.36, P=0.002\#$
COVID-19	133	33 (76.7%)	100 (92.6%)	$\chi^2=7.35, P=0.007^{**}$
Asymptomatic infection	18	10 (23.3%)	8 (7.4%)	
Negative results of RT-PCR	684	282 (67.6%)	402(63.6%)	$\chi^2=1.79, P=0.18\#$
Not tested	214	92 (22.1%)	122(19.3%)	$\chi^2=1.18, P=0.28\#$

Total (n)	1049	417	632
-----------	------	-----	-----

Notes: * Eight of which were brothers and sisters from 4 families. # Comparison on incidence between children contacts and adult contacts; ** Comparison on constituent ratio of COVID-19 between pediatric cases and adult cases with SARS-CoV-2 infection

Table 2 Signs, symptoms, imaging finding, leucocytes and lymphocytes counts in 43 pediatric patients with COVID-19 in Zhejiang Province

Signs and symptoms	Initial symptoms or signs on admission (n)	During the whole course (n)
Asymptomatic	10	10
Fever	17	21
Highest temperature (°C)		
≤37.4°C	14	10
37.5-38.9°C	14	17
≥39°C	3	4
Cough	5	12
Stuffy running nose	4	7
Fatigue	1	3

Diarrhea	1	3
Vomiting and abdominal pain	0	2
Liver function abnormality*	1	4
Shortness of breath, chest tightness, and headache	0	1
Imaging finding	16	20
Bilateral involvement	3	4
Unilateral involvement	13	16
Bilateral or unilateral ground-glass opacity	7	9
Leucocytes $<4 \times 10^9$ /L (n)	4	5
lymphocytes $<1.2 \times 10^9$ /L (n)	1	3

Note: *Alanine aminotransferase, ALT 54-112 U/L, Aspartate aminotransferase, AST 57-124 U/L

Table 3 Several time points in 20 patients with SARS-CoV-2 pneumonia

	Minimum	Maximum	Mean (SD)
Days from illness onset to visit	1	10	3.3 (2.0)
Days from be exposed to illness onset	4	21	9.2 (4.1)
Symptomatic duration (days)	1	10	4.6 (2.4)
Days from exposure to imaging finding	5	36	7 (6,10)
Days from illness onset or positive RT-PCR finding to imaging finding	1	30	3 (1,6)
Days in hospital	3	32	20.2 (7.9)
Days from the illness onset to results of RT-PCR in respiratory samples turned to negative	3	28	14.0 (6.6)

Note Four children had “asymptomatic” pneumonia, one of which was diagnosed after being discharged.

Table 4 Effectiveness on potential viral excretion time in children treated with different therapeutic regimens

	Without antiviral therapy (n=5)	Antiviral monotherapy* (n=21)	With two or more than two antiviral drugs# (n=17)	Statistic values
Days in hospital	18.0 (11.0)	19.9(6.2)	18.5(8.8)	$\chi^2=0.14$, $P=0.69$
Days of the course when antiviral treatment initiated	-	2.4 (1.9)	2.3 (1.3)	$t=0.12$, $P=0.90$
Days for antiviral treatment	-	13.2 (7.7)	15.9 (7.3)	$t=1.12$, $P=0.27$
Days when results of respiratory RT-PCR turned to negative	11.2 (5.2)	16.8(7.7)	12.7(6.6)	$\chi^2= 5.62$, $P=0.06$
Days when results of fecal RT-PCR turned to negative	38.0 (17.8)	30.2 (9.4)	28.8(14.7)	$\chi^2=1.37$, $P=0.51$
Numbers of the patients with positive RT-PCR results in feces when being discharged	2 / 3	9 / 18	6 / 12	$P=0.34$

Note: A, atomization inhalation with interferon- α 2b; B, oral lopinavir and ritonavir tablets; C, oral arbidol; D, oral or intravenous ribavirin. * 19 cases with A, one case with B and D respectively; #

Six cases with A and B; five cases with A and C; one cases with A and D. ** Two cases with A, B and C; One cases with A, C and D; One case with A, B and C as initial treatment, and switched to A, C, and oral darunavir/cobicistat in combination later because of serious vomiting, abdominal pain and diarrhea; One case with A, B, C and oral oseltamivir.