
Title: Clinical characteristics and factors associated with long-term viral excretion in patients with SARS-CoV-2 infection: a single center 28-day study

Authors: Ding Shi^{a,#}, Wenrui Wu^{a#}, Qing Wang^{a#}, Kaijin Xu^a, Jiaojiao Xie^a, Jingjing Wu^a, Longxian Lv^a, Jifang Sheng^a, Jing Guo^a, Kaicen Wang^a, Daiqiong Fang^b, Yating Li^a, & Lanjuan Li^{*,a}

Affiliations:

^a State Key Laboratory for Diagnosis and Treatment of Infectious Diseases, National Clinical Research Center for Infectious Diseases, Collaborative Innovation Center for Diagnosis and Treatment of Infectious Diseases, The First Affiliated Hospital, College of Medicine, Zhejiang University, Hangzhou, 310003, China

^b Department of Endocrinology and Metabolism, The First Affiliated Hospital, College of Medicine, Zhejiang University, Hangzhou, China

[#]Ding Shi, Wenrui Wu, Qing Wang contributed equally to this manuscript.

*Corresponding author: Lanjuan Li E-mail: ljli@zju.edu.cn

Tel.: 0571-87236458 Fax: 86-571-87081933

Summary: Male sex, immunoglobulin use, APACHE II score and lymphocyte count were independent factors associated with the duration of SARS-CoV-2 RNA shedding. The current antiviral protocol and low-to-moderate dosage of corticosteroid had little effect on the duration of viral excretion.

Corresponding author contact information:

Lanjuan Li, MD

Professor of Infectious Diseases,

State Key Laboratory for Diagnosis and Treatment of Infectious Diseases

First Affiliated Hospital Zhejiang University College of Medicine

79 Qingchun Road, Hangzhou 310003, P.R. China

Tel: 86-571-87709001 Fax: 86-571-87081933

Email: ljli@zju.edu.cn

Alternate Corresponding Author:

Ding Shi, MD/PHD

State Key Laboratory for Diagnosis and Treatment of Infectious Diseases

First Affiliated Hospital Zhejiang University College of Medicine

79 Qingchun Road, Hangzhou 310003, P.R. China

Email: shiding@zju.edu.cn

Clinical characteristics and factors associated with long-term viral excretion in patients with SARS-CoV-2 infection: a single center 28-day study

Abstract:

Background: Despite the ongoing spread of COVID-19, knowledge about factors affecting prolonged viral excretion is limited.

Methods: In this study, we retrospectively collected data from 99 hospitalized patients with COVID-19 between January 19 and February 17 in Zhejiang Province, China. We classified them into two groups based on whether the virus test results eventually became negative. Cox proportional hazards regression was used to evaluate factors associated with SARS-CoV-2 shedding.

Results: Among 99 patients, 61 patients had SARS-CoV-2 clearance (virus-negative group), but 38 patients had sustained positive results (virus-positive group). The median duration of SARS-CoV-2 excretion was 15 days (IQR 12-19) among the virus-negative patients. The shedding time was significantly increased if fecal SARS-CoV-2 RNA test results was positive. Male sex (HR, 0.58 [95% CI, 0.35-0.98]), immunoglobulin use (HR, 0.42 [95% CI, 0.24-0.76]), APACHE II score (HR, 0.89 [95% CI, 0.84-0.96]), and lymphocyte count (HR, 1.81 [95% CI, 1.05-3.1]) were independent factors associated with a prolonged duration of SARS-CoV-2 shedding. Antiviral therapy and corticosteroid treatment were not independent factors.

Conclusions: SARS-CoV-2 RNA clearance time was associated with sex, disease severity and lymphocyte function. The current antiviral protocol and low-to-moderate dosage of corticosteroid had little effect on the duration of viral excretion.

Keywords: SARS-CoV-2, viral excretion, disease severity, risk factors, lymphocyte function

Introduction:

Since December 2019, a pneumonia of unknown cause sparked an outbreak in Wuhan, China, which spread rapidly to other provinces of China and then to other countries. The causative pathogen was identified as a novel β -coronavirus, which was named the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2)[1], and the disease was named coronavirus disease 2019 (COVID-19). The basic clinical and epidemiologic characteristics of newly emerging COVID-19 had been reported in different cases[2, 3]. The SARS-CoV-2 infection causes severe respiratory symptoms, which indicates that the respiratory tract is the main source of human-to-human transmission. In addition, SARS-CoV-2 RNA has been detected in stool specimens as well as gastrointestinal tissue, which indicates that the fecal-oral route might be another nonnegligible transmission method[4]. So far, there is no vaccine or effective antiviral treatment for SARS-CoV-2.

Little information is available on the duration of SARS-CoV-2 excretion and the factors influencing viral shedding. It was reported that the RNA of Middle East respiratory syndrome coronavirus (MERS-CoV) could be detected in respiratory specimens for more than one month[5]. Severe acute respiratory syndrome coronavirus (SARS-CoV) RNA could be detected in sputum for 20 days and even more than 100 days in stool specimens from illness onset[6]. Prolonged viral shedding in the respiratory tract is associated with severe outcomes in patients with seasonal influenza and MERS[7, 8]. Features of viral shedding are useful for the evaluation of disease pathogenesis, disease course characteristics and treatment effects for SARS-CoV-2 infection.

Hence, we conducted a study on 99 hospitalized patients with confirmed SARS-CoV-2 infection identified between January 19 and February 17, 2020. We compared the clinical characteristics between patients with negative virus detection during hospitalization and those with persistent positive virus detection to evaluate the impact of different factors on the duration of SARS-CoV-2 shedding.

Patients and methods

We performed a single-center, retrospective cohort study from January 19 to February 17, 2020 of SARS-CoV-2 outbreak in Zhejiang Province at the First Affiliated Hospital of Zhejiang University. A total of 99 SARS-CoV-2 infected patients, including 15 patients transferred from other hospitals, were recruited in this study. All patients were observed for 28 days after admission unless they met the discharge criteria. COVID-19 was diagnosed on the basis of interim WHO guidance[9]. Only patients with laboratory confirmed infection, including one asymptomatic patient, were enrolled in this study. The asymptomatic patient was admitted to the hospital with his relatives and presented with fever and cough symptoms after admission for 2 days. The criteria for discharge were the absence of fever for at least 3 days, improvements in chest CT and clinical symptoms, and two consecutive negative results for SARS-CoV-2 RNA in respiratory samples obtained at least 1 day apart. Respiratory specimens (nasopharyngeal swabs, sputum, or endotracheal aspirates) and stool specimens were collected daily to determine the amount of SARS-CoV-2 RNA by using real-time reverse transcription polymerase chain reaction (RT-PCR), as previously described[3]. We defined COVID-

19 patients as virus-negative if the respiratory specimens were SARS-CoV-2 RNA negative across more than two intervals. Otherwise, we defined patients as virus-positive if their specimens were persistently SARS-CoV-2 RNA positive during hospitalization.

Data collection

The clinical data included demographic characteristics, medical comorbidities, laboratory findings, progression, and treatment of clinical illness. Documented medical comorbidities included hypertension, diabetes mellitus, cardiac diseases, chronic lung disease, pregnancy, and immunosuppressive disease. Complete blood counts, coagulation profiles, inflammatory indexes including C-reactive protein (CRP) and procalcitonin (PCT) as well as serum biochemical parameters including renal and liver function, creatine kinase, lactate dehydrogenase were tested for all patients at admission. This study was approved by the National Health Commission of China and Ethics Commission of the First Hospital of Zhejiang Province. Written informed consent was obtained from all participating patients.

Shedding duration

Shedding duration was defined as the time from illness onset to viral shedding cessation. Illness included but was not limited to the following symptoms: fever, sore throat, cough, runny nose, nausea and vomiting, diarrhea and fatigue. Illness onset was defined as the first appearance of the symptoms or the first RT-PCR positive result for the virus

for asymptomatic patients. Shedding cessation was defined as the occurrence of two consecutive RT-PCR negative results of respiratory specimens in a 24-hour interval.

Statistical Analyses

For most variables, descriptive statistics such as the median with interquartile range (IQR; for data with skewed distribution) and proportion (%) were calculated. The T test, Mann-Whitney U test, and Kruskal-Wallis test were used for continuous variables. The χ^2 test and Fisher's exact test were used for categorical variables. To identify risk factors associated with a prolonged duration of SARS-CoV-2 RNA shedding, we performed a time-dependent Cox proportional hazards model adjusted for baseline covariates. Outcomes were defined as time to SARS-CoV-2 RNA negativity, and we censored patients if they never cleared SARS-CoV-2 RNA during hospitalization. Kaplan-Meier curves were used to estimate the cumulative SARS-CoV-2 RNA negativity rate and the stratified log-rank statistic to compare the differences in viral clearance between the two groups. Statistical analyses were performed using SPSS software, version 20.0. For all analyses, probabilities were two-tailed, and a two-tailed *P* value of < 0.05 was considered significant.

Results

Patients characteristics

A total of 99 patients with SARS-CoV-2 infection were included in this study from January 19 to February 17, 2020. During hospitalization, 61 patients (61.6%) had

SARS-CoV-2 clearance (virus-negative group), but 38 patients (38.4%) still tested positive for SARS-CoV-2 RNA (virus-positive group). The demographic and clinical characteristics of COVID-19 patients are summarized in Table 1. The median age was 54 years (IQR, 39-46), and 22.2% of the patients were aged >65 years. The median age of patients in the virus-positive group was significantly greater than that of those in the virus-negative group (61.5 [50.3-72.3] vs 50 [38.5-59]; $P < 0.01$). The number of elderly patients (>65 years old) in the virus-positive group was significantly higher than that in the virus-negative group ($P < 0.01$). A total of 61.6% of patients were male, but no significant difference in sex was found between the two groups. Hypertension ($n=36$, 36.4%) and diabetes mellitus ($n=16$, 16.2%) were the most common comorbidities among the COVID-19 patients. Additionally, a few patients had chronic lung disease, cardiac disease, immunosuppressive disease and pregnancy, but there were no significant differences between the two groups (Table 1).

Laboratory Abnormalities

Among laboratory indicators tested at admission, inflammatory indexes such as CRP ($P=0.01$) and PCT ($P=0.01$) of the persistently virus positive patients were significantly higher than those of virus-negative patients. The oxygenation index ($P=0.03$), lymphocyte counts ($P=0.001$), albumin ($P=0.03$), CD4+ T cells ($P=0.01$), CD8+ T cells ($P=0.02$), and B cells ($P=0.01$) were significantly lower in virus-positive group than in the virus-negative group (Table 1). Other laboratory indicators, such as leukocyte count, hemoglobin, platelet count, coagulation profile, creatine, liver function, and

myocardial enzyme showed no difference between the two groups (Supplementary Table 1). From the results, persistent virus-positive patients showed lower immune cell counts and higher inflammation levels.

Prolonged duration of SARS-CoV-2 Shedding

Only 1 patient (1.6%) tested SARS-CoV-2 negative within 5 days, 9 patients (14.7%) tested negative within 10 days, and 49 (80.3%) tested negative within 20 days from illness onset. In addition, a small subset of 12 patients with SARS-CoV-2 had detectable levels of the virus up to 30 days from symptom onset. The median time of persistent viral shedding was 16 days (IQR 13-21). The virus-positive group showed a median viral shedding time of 19 days, which was significantly longer than that in the virus-negative group (median 15, IQR 12-19, $P=0.002$, Table 2). No apparent difference in time from illness onset to COVID-19 diagnosis was found between the two groups. SARS-CoV-2 RNA was positively detected in the stool specimens of 21 patients (21.2%), nine of whom were virus negative (14.8%) and twelve of whom were virus positive (31.6%, $P<0.01$). The results of fecal SARS-CoV-2 RNA testing turned negative in only 4 patients (19.0%) before February 17 and remained positive in 11 patients (18.0%) after testing for sputum viral RNA turned negative.

Treatment disparity between two groups

A majority of the patients (97.9%) received antiviral treatment (ARV), and 81 patients (81.8%) received two combined antiviral treatments with arbidol (200 mg 3 times daily)

and lopinavir and ritonavir (LPV/RTV, 400 mg twice daily and 100 mg twice daily, respectively), whereas 16 patients (16.2%) received single antiviral treatment with arbidol or LPV/RTV (Table 2). Darunavir (800 mg once daily) was prescribed to replace LPV/RTV if patients experienced significant drug side effects. The duration from illness onset to ARV start was 6 days (IQR 4-8.5). There was no significant difference in ARV start time between the virus-positive and virus-negative groups. A total of 77 patients (77.8%) received corticosteroid treatment with an initial dosage of 60 mg/d (IQR 40-80), and the time from illness onset to corticosteroid treatment was 8 days (IQR 6-10). There was no significant difference in the application, initial dosage or start time of corticosteroid treatment between the two groups (Table 2). Forty-nine patients (49.5%) were treated with antibiotics, and the median time from illness onset to antibiotic use was 9 days (IQR 6-13). There was no difference in antibiotic use proportion or antibiotic start time between the two groups ($P>0.05$). In addition, 43 patients (43.4%) received intravenous immunoglobulin treatment. The proportion of patients receiving immunoglobulin therapy in the virus-positive group was much higher than that in the virus-negative group (63.2% vs 31.1%, $P=0.002$).

Clinical outcome

The patients with persistent SARS-CoV-2 RNA positivity showed greater disease severity (Table 2). Among 99 COVID-19 patients, a total of 30 (30.3%) were transferred to the intensive care unit (ICU), and the patients in the virus-positive group had a higher rate of ICU admission than the virus-negative group (52.6% vs. 16.4%,

$P < 0.01$). The median length of ICU stay for the 30 patients was 7.5 days (IQR 4-11), and in the virus-positive group, the length of ICU stay was 8.5 days (IQR 6.3-11), which was longer than that in the virus-negative group (4 days, IQR 3-5.8, $P < 0.01$). The Acute Physiology and Chronic Health Evaluation II (APACHE II) score was used to evaluate the severity of hospitalized patients as well as to predict mortality[10]. The APACHE II score was 9.5 (IQR 5-15) in virus-positive group, which was much higher than that of virus-negative patients, with a score of 5 (IQR 2-8). Mechanical ventilation was performed in a minority of patients (12.1%). Compared with the virus-negative group, the virus-positive group had a higher rate of mechanical ventilation (26.3% vs. 3.3%, $P < 0.01$). Moreover, all patients receiving extracorporeal membrane oxygenation (ECMO) treatment were from the virus-positive group (15.8% vs. 0%, $P < 0.01$), indicating more severe lung injury and oxygenation.

Risk factors for prolonged viral shedding

We conducted a multivariable time-dependent Cox proportional hazards model to identify risk factors associated with the duration of SARS-CoV-2 RNA shedding (Table 3). Male sex (HR, 0.58 [95% CI, 0.35-0.98]), immunoglobulin use (HR, 0.42 [95% CI, 0.24-0.76]), APACHE II score (HR, 0.89 [95% CI, 0.84-0.96]), and lymphocyte count (HR, 1.81 [95% CI, 1.05-3.1]) were independent factors associated with the duration of SARS-CoV-2 RNA shedding, whereas current smoking, hypertension, diabetes, days from illness onset to ARV therapy, ARV combination therapy, corticosteroid treatment, and days from illness onset to corticosteroid treatment and antibiotic treatment were not

independent factors (Table 3). Interestingly, the use of immunoglobulin may prolong the duration of viral shedding. We found that the disease tended to be more severe in patients prescribed immunoglobulins, and these patients were older with higher APACHE II scores and worse immune function than those who did not use immunoglobulins (Supplementary Table 2).

Using Kaplan-Meier survival analysis, we found SARS-CoV-2 RNA clearance was associated with disease severity, which was significantly delayed in patients who were transferred to ICU and whose APACHE II scores >10 compared with that of patients in the general ward and whose APACHE II scores ≤ 10 (log-rank test, $P<0.001$; Figure 1A, 1B). Compared with patients with a high lymphocyte count ($> 0.5 \times 10^9/L$), patients with low immunity (lymphocyte count $\leq 0.5 \times 10^9/L$) showed a significant delay in virus clearance (log-rank test, $P=0.002$; Figure 1C). It took longer for fecal viral RNA-positive patients to clear SARS-CoV-2 RNA than fecal viral RNA-negative patients (log-rank test, $P=0.003$; Figure 1D).

Discussion

Despite the ongoing spread of COVID-19, knowledge of the factors affecting prolonged viral shedding is still limited. Thus, we conducted a retrospective, single-center study of 99 hospitalized patients with confirmed SARS-CoV-2 infection identified between January 19 and February 17, 2020 in Zhejiang Province, China. We identified that male sex, immunoglobulin use, APACHE II score, and lymphopenia were independent risk factors associated with the duration of SARS-CoV-2 RNA shedding, whereas ARV

combination therapy and corticosteroid treatment were not independent factors. In addition, patients with fecal viral RNA tested positive needed more time to clear SARS-CoV-2.

Consistent with previous findings, median age of hospitalized patients with SARS-CoV-2 infection was 54 years, and 22.2% patients were aged > 65 years [11]. Patients with persistent SARS-CoV-2 RNA positivity were older. Although age >65 years was considered as a risk factor associated with disease severity [2]. However, our study did not find age to be a risk factor for viral clearance. The key difference in age between the two groups may be related to weak immune response as well as coexisting illnesses. A majority of COVID-19 patients had hypertension and diabetes, and a small proportion had chronic lung disease, cardiac disease, immunosuppression or pregnancy. COVID-19 patients with underlying comorbidities, including hypertension, diabetes, cardiovascular disease, were more likely to be transferred to ICU [12]. Underlying diseases has been shown to be related to prolonged viral shedding in SARS patients[6]. However, we found no significant differences between virus-positive patients and virus-negative patients with regard to comorbidity. This result suggests that underlying diseases such as hypertension and diabetes may be related to the patient's disease prognosis but have little to do with the duration of viral shedding.

Male sex was found to be an independent risk factor for prolonged SARS-CoV-2 viral shedding. In SARS-CoV infection, males experience higher mortality than females[13].

Imbalanced levels of angiotensin-converting enzyme 2 (ACE2) between males and females may play an important role in the sex-based differences in the response to the disease[14]. ACE2 has been identified as the host receptor of SARS-CoV, which regulates both cross-species and human-to-human transmissions in SARS-CoV infection [15]. SARS-CoV-2 also use the ACE2 receptor to facilitate viral entry into target cells [16].

Consistent with other reports, the most common laboratory abnormalities among the COVID-19 patients were depressed total lymphocyte and lymphocyte subset counts, as well as an elevated CRP level[2, 3, 17]. Compared with the virus-negative patients, patients with persistent SARS-CoV-2 positivity had lower lymphocyte count and T-cell subset count but higher CRP and PCT level. This suggested more severe cellular immune deficiency and inflammatory activation in SARS-CoV-2 virus-positive patients. Decreases in T cell counts were strongly associated with the severity and progression of SARS and MERS in patients[18, 19]. The underlying reasons for lymphopenia may be the direct infection of lymphocytes by SARS-CoV-2, lymphocyte sequestration in the lung or cytokine-mediated lymphocyte trafficking[20, 21]. All of which may lead to extend the virus clearance time.

Although various drugs have been tested for COVID-19, there is still an urgent need for a clinically proven, effective antiviral treatment for SARS-CoV-2 infection. In this study, the antiviral therapy mainly included LPV/RTV and arbidol. Nevertheless, we

observed that there was no apparent effect on viral shedding from the use of these antiviral drugs or their combined use. Previous researches on the efficacy of LPV/RTV in SARS and MERS showed disparate results. Compared with those using ribavirin alone, improved outcomes for SARS and decreased viral loads were reported for patients using a combination of LPV/RTV and ribavirin[22], whereas no obvious antiviral effect against MERS-CoV with lopinavir was observed in vitro[23]. Another study revealed that LPV/RTV could improve pulmonary function without inhibiting virus replication in MERS-CoV infection [24]. Current evidence regarding the effect of arbidol against SARS-CoV-2 was rare. In vitro experiment have indicated that arbidol showed some antiviral activity against SARS[25]. It was previously demonstrated that remdesivir had excellent antiviral activity against MERS-CoV in vitro and in vivo[24]. Moreover, remdesivir exhibited a beneficial effect in the first case in America[26]. This novel nucleotide analogue had not been put into use in our center yet, while its therapeutic effect in illness improvement is worth attention. Overall, further studies are warranted to uncover the exact efficacy of these different antiviral agents on viral shedding.

Corticosteroids have been used frequently for the treatment of severe patients infected with coronavirus by reducing inflammatory-induced lung injury[3]. Several studies have reported that the use of corticosteroids is associated with delayed viral RNA clearance in MERS or SARS patients and even with higher mortality in influenza pneumonia[27-29]. In this study, corticosteroid usage did not independently increase

the risks for viral shedding regardless of the initial dosage or an earlier implementation. However, the percentage of corticosteroid use in our center reached up to 77%, and the dosages prescribed were relatively lower for these COVID-19 patients than for MERS and SARS patients in previous studies. It was reported that high dosages of corticosteroids (>150 mg/d) showed a similar influence on H7N9 infection with prolonged viral shedding time, whereas proper use of corticosteroids was associated with clinical improvement in SARS[30, 31]. Thus, more clinical data are urgently needed to indicate the effects of corticosteroids and the optimal scheme of their usage. In addition, the beneficial effect of immunoglobulin has also been observed in viral infection[32]. Our study found immunoglobulin application was associated with delayed viral clearance. In fact, patients using immunoglobulin were more likely to be in the more severe condition. Thus, more studies should be done to confirm the association between immunoglobulin treatment and viral shedding.

We identified the predictive value of the APACHE II score in SARS-CoV-2 infection. Moreover, patients transferred to the ICU showed delayed SARS-CoV-2 RNA clearance. The result indicated that SARS-CoV-2 viral shedding was associated with disease severity. Similarly, MERS-CoV sub-genomic mRNA was detected more frequently in the severe group than in the mild group[5]. Severe patients had more prolonged MERS-CoV shedding, up to 18-27 days following the onset of symptoms[33]. SARS patients with APACHE II scores ≥ 20 were associated with an increased risk of transmission of SARS-CoV[34]. It can be explained by the higher

APACHE II score, more severe immunosuppression and higher viral load among patients in the ICU.

Fecal-oral transmission was considered as an additional route for SARS-CoV-2 spread. In this study, we found a high percentage of COVID-19 patients with positive fecal virus results, who needed more time to clear SARS-CoV-2 than those with negative fecal virus results. Fecal viral RNA can remain positive even after respiratory specimens test results return negative. Consistent with our results, stool specimens tested positive in up to 53% of COVID-19 patients, and the positive staining of ACE2 and SARS-CoV-2 nucleocapsid protein was observed in gastrointestinal epithelium as well as isolated infectious SARS-CoV-2 from feces[4]. The duration of SARS fecal viral excretion could last for more than 100 days[6]. Positive fecal viral RNA was highly suggestive of virus intestinal infection; thus, a longer isolated observation time should be considered in these COVID-19 patients.

Our study has some notable limitations. First, we only observed the hospitalization of COVID-19 patients for 28 days, and many patients remained in hospital with SARS-CoV-2 RNA positivity. The duration of viral shedding may be longer than this cut-off time point. Second, some patients were transferred to our hospital after the disease had progressed and become worse; hence, our cohort might represent a more severe range of COVID-19.

In conclusion, we found that male sex, immunoglobulin use, APACHE II score, and lymphopenia were independent risk factors associated with the duration of SARS-CoV-2 RNA shedding, whereas ARV combination therapy and corticosteroid treatment were not. A prolonged SARS-CoV-2 RNA clearance time was associated with disease severity and fecal viral RNA positivity. Our findings suggest that more suitable antiviral agents and low-to-moderate corticosteroids rather than high dosages should be considered in the treatment of severe COVID-19, while seeking solutions to enhance immune function and to reduce the inflammatory response were also of great importance for viral clearance. In addition, more data based on randomized and controlled multicenter studies are needed to rigorously assess the clinical relevance of our proposed indicators.

Acknowledgments

This study was funded by Science and Technology Department of Zhejiang Province [2020c03123-1]; Zhejiang Provincial Natural Science Foundation of China [LQ20H030010]; the National Key Research and Development Program of China [2018YFC2000500]. No authors report conflicts of interest. We thank all patients involved in the study and all the front-line medical staff in the First Affiliated Hospital, School of Medicine, Zhejiang University.

References:

1. Zhu N, Zhang D, Wang W, et al. A Novel Coronavirus from Patients with Pneumonia in China, 2019. *The New England journal of medicine* **2020**; 382:727-33.
2. Guan W-J, Ni Z-Y, Hu Y, et al. Clinical Characteristics of Coronavirus Disease 2019 in China. *The New England journal of medicine* **2020**.
3. Huang C, Wang Y, Li X, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet (London, England)* **2020**; 395:497-506.
4. Xiao F, Tang M, Zheng X, Liu Y, Li X, Shan H. Evidence for gastrointestinal infection of SARS-CoV-2. *Gastroenterology* **2020**.
5. Park WB, Poon LLM, Choi S-J, et al. Replicative virus shedding in the respiratory tract of patients with Middle East respiratory syndrome coronavirus infection. *Int J Infect Dis* **2018**; 72.
6. Liu W, Tang F, Fontanet A, et al. Long-term SARS coronavirus excretion from patient cohort, China. *Emerging Infect Dis* **2004**; 10:1841-3.
7. Corman VM, Albarrak AM, Omrani AS, et al. Viral Shedding and Antibody Response in 37 Patients With Middle East Respiratory Syndrome Coronavirus Infection. *Clin Infect Dis* **2016**; 62:477-83.
8. McGeer A, Green KA, Plevneshi A, et al. Antiviral therapy and outcomes of influenza requiring hospitalization in Ontario, Canada. *Clin Infect Dis* **2007**; 45:1568-75.
9. WHO. Clinical management of severe acute respiratory infection when Novel coronavirus (nCoV) infection is suspected: interim guidance.
10. Zhou X-Y, Ben S-Q, Chen H-L, Ni S-S. A comparison of APACHE II and CPIS scores for the prediction of 30-day mortality in patients with ventilator-associated pneumonia. *Int J Infect*

Dis **2015**; 30:144-7.

11. Li Q, Guan X, Wu P, et al. Early Transmission Dynamics in Wuhan, China, of Novel Coronavirus-Infected Pneumonia. *The New England journal of medicine* **2020**.

12. Wang D, Hu B, Hu C, et al. Clinical Characteristics of 138 Hospitalized Patients With 2019 Novel Coronavirus-Infected Pneumonia in Wuhan, China. *JAMA* **2020**.

13. Leong HN, Earnest A, Lim HH, et al. SARS in Singapore--predictors of disease severity. *Annals of the Academy of Medicine, Singapore* **2006**; 35:326-31.

14. Clotet-Freixas S, Soler MJ, Palau V, et al. Sex dimorphism in ANGII-mediated crosstalk between ACE2 and ACE in diabetic nephropathy. *Laboratory investigation; a journal of technical methods and pathology* **2018**; 98:1237-49.

15. Wan Y, Shang J, Graham R, Baric RS, Li F. Receptor recognition by novel coronavirus from Wuhan: An analysis based on decade-long structural studies of SARS. *Journal of virology* **2020**.

16. Zhang H, Penninger JM, Li Y, Zhong N, Slutsky AS. Angiotensin-converting enzyme 2 (ACE2) as a SARS-CoV-2 receptor: molecular mechanisms and potential therapeutic target. *Intensive Care Med* **2020**.

17. Wan S, Yi Q, Fan S, et al. Characteristics of lymphocyte subsets and cytokines in peripheral blood of 123 hospitalized patients with 2019 novel coronavirus pneumonia (NCP). *medRxiv* **2020**:2020.02.10.20021832.

18. Li T, Qiu Z, Zhang L, et al. Significant changes of peripheral T lymphocyte subsets in patients with severe acute respiratory syndrome. *The Journal of infectious diseases* **2004**; 189:648-51.

19. Ko J-H, Park GE, Lee JY, et al. Predictive factors for pneumonia development and

progression to respiratory failure in MERS-CoV infected patients. *J Infect* **2016**; 73:468-75.

20. Lang ZW, Zhang LJ, Zhang SJ, et al. A clinicopathological study of three cases of severe acute respiratory syndrome (SARS). *Pathology* **2003**; 35:526-31.

21. Collins AR. In vitro detection of apoptosis in monocytes/macrophages infected with human coronavirus. *Clinical and diagnostic laboratory immunology* **2002**; 9:1392-5.

22. Chu CM, Cheng VCC, Hung IFN, et al. Role of lopinavir/ritonavir in the treatment of SARS: initial virological and clinical findings. *Thorax* **2004**; 59:252-6.

23. Chan JFW, Chan K-H, Kao RYT, et al. Broad-spectrum antivirals for the emerging Middle East respiratory syndrome coronavirus. *J Infect* **2013**; 67:606-16.

24. Sheahan TP, Sims AC, Leist SR, et al. Comparative therapeutic efficacy of remdesivir and combination lopinavir, ritonavir, and interferon beta against MERS-CoV. *Nat Commun* **2020**; 11:222.

25. Khamitov RA, Loginova SI, Shchukina VN, Borisevich SV, Maksimov VA, Shuster AM. [Antiviral activity of arbidol and its derivatives against the pathogen of severe acute respiratory syndrome in the cell cultures]. *Vopr Virusol* **2008**; 53.

26. Holshue ML, DeBolt C, Lindquist S, et al. First Case of 2019 Novel Coronavirus in the United States. *The New England journal of medicine* **2020**; 382:929-36.

27. Arabi YM, Mandourah Y, Al-Hameed F, et al. Corticosteroid Therapy for Critically Ill Patients with Middle East Respiratory Syndrome. *Am J Respir Crit Care Med* **2018**; 197:757-67.

28. Lee N, Allen Chan KC, Hui DS, et al. Effects of early corticosteroid treatment on plasma SARS-associated Coronavirus RNA concentrations in adult patients. *J Clin Virol* **2004**; 31:304-9.

29. Ni Y-N, Chen G, Sun J, Liang B-M, Liang Z-A. The effect of corticosteroids on mortality of patients with influenza pneumonia: a systematic review and meta-analysis. *Crit Care* **2019**; 23:99.
30. Chen R-C, Tang X-P, Tan S-Y, et al. Treatment of severe acute respiratory syndrome with glucosteroids: the Guangzhou experience. *Chest* **2006**; 129:1441-52.
31. Sung JJY, Wu A, Joynt GM, et al. Severe acute respiratory syndrome: report of treatment and outcome after a major outbreak. *Thorax* **2004**; 59:414-20.
32. Mofenson LM, Moye J, Jr., Bethel J, Hirschhorn R, Jordan C, Nugent R. Prophylactic intravenous immunoglobulin in HIV-infected children with CD4+ counts of 0.20 x 10⁹/L or more. Effect on viral, opportunistic, and bacterial infections. The National Institute of Child Health and Human Development Intravenous Immunoglobulin Clinical Trial Study Group. *Jama* **1992**; 268:483-8.
33. Min C-K, Cheon S, Ha N-Y, et al. Comparative and kinetic analysis of viral shedding and immunological responses in MERS patients representing a broad spectrum of disease severity. *Scientific reports* **2016**; 6:25359.
34. Raboud J, Shigayeva A, McGeer A, et al. Risk factors for SARS transmission from patients requiring intubation: a multicentre investigation in Toronto, Canada. *PloS one* **2010**; 5:e10717.

Figure legends

Figure 1 A, Cumulative proportion of patients with detectable SARS-CoV-2 RNA among patients who were transferred to ICU and those who were in general ward by day after illness onset. B, Cumulative proportion of patients with detectable SARS-CoV-2 RNA among those whose APACHE II score >10 and those whose APACHE II ≤ 10 by day after illness onset. C, Cumulative proportion of patients with detectable SARS-CoV-2 RNA among those whose lymphocyte count $\leq 0.5 \times 10^9/L$ and those whose lymphocyte count $> 0.5 \times 10^9/L$ by day after illness onset. D, Cumulative proportion of patients with detectable SARS-CoV-2 RNA among those with positive fecal viral RNA and those with negative fecal viral RNA by day after illness onset.

Figure 1




Table 1 Clinical Characteristics of 99 Hospitalized Patients With SARS-CoV-2 Infection.

Characteristic	Total (N=99)	Virus negative(N=61)	Virus positive (N=38)	P Value
Demographics, n (%)				
Age, median (IQR)	54 (39-64)	50 (38.5-59)	61.5 (50.3-72.3)	0.001
>65 years old	22 (22.2)	8 (13.1)	14 (36.8)	0.006
BMI, kg/m ²	24.2 (21.9-26)	24.2 (21.7-26.0)	24.0 (22.1-26.2)	0.84
Male sex	61 (61.6)	35 (57.4)	26 (68.4)	0.20
Current smoking	13 (13.1)	8 (13.1)	5 (13.2)	1.00
Comorbidity, n(%)				
Hypertension	36 (36.4)	19 (31.1)	17 (44.7)	0.17
Diabetes	16 (16.2)	8 (13.1)	8 (21.1)	0.29
Chronic lung disease ^a	5 (5.1)	2 (3.3)	3 (7.9)	0.37
Cardiac disease ^b	7 (7.1)	2 (3.3)	5 (13.2)	0.10
Immunosuppression ^c	2 (2.0)	1 (1.6)	1 (2.6)	1.00
Pregnancy	3 (3.0)	3 (4.9)	0	0.28
Initial laboratory findings, median (IQR)				
PaO ₂ /FiO ₂ , mmHg	274.6 (187.4-383)	288.3 (228.4-398.4)	250.3 (149.8-370.4)	0.03
Leukocyte count,10 ⁹ /L	5.9 (4.1-10.0)	5.2 (3.6-8.4)	7.2 (4.8-10.8)	0.07
Lymphocyte count,10 ⁹ /L	0.8 (0.5-1.2)	0.9 (0.6-1.3)	0.5 (0.3-0.9)	0.001
ALB, g/L	38 (33.9-43.4)	40.6 (35.3-43.6)	35.8 (32.4-40.8)	0.03
CRP, mg/L	23.9 (9.1-51.1)	16.1 (7.8-33.1)	45 (12.9-81.8)	0.01
PCT, ng/mL	0.06 (0.03-0.09)	0.06 (0.03-0.08)	0.07 (0.03-0.25)	0.01

CD4+T, / μ L	149 (79-324)	199 (108-476.3)	94 (67-210)	0.01
CD8+T, / μ L	133 (77-278)	152.5 (99.5-374.5)	79 (49-159)	0.02
B cell, / μ L	112 (66-178)	146.5 (79.3-188.8)	70 (32-123)	0.01
NK cell, / μ L	106 (65-187)	133 (69.3-207.5)	97 (53-158)	0.06

Abbreviations: IQR, interquartile range; BMI, body mass index; ALB, albumin; CRP, C-reactive protein; PCT, procalcitonin; NK cell, natural killer cell.

^a Chronic lung disease includes chronic obstructive pulmonary disease and interstitial lung disease

^b Cardiac disease includes congestive heart disease and coronary atherosclerotic heart disease.

^c Immunosuppression is defined as the receipt of anti-rejection drugs, or Acquired Immune Deficiency Syndrome.

Table2 Treatment and Outcome of 99 Hospitalized Patients With SARS-CoV-2 Infection.

Variable	Total	Virus negative	Virus positive	P Value
	(N=99)	(N=61)	(N=38)	
Virological result, d				
Days of viral persistent positive	16 (13-21)	15 (12-19)	19 (15-24)	0.002
Days from illness onset to diagnosis	5 (2-7)	5 (2.5-7.5)	4 (1-6)	0.05
Fecal viral positive	21 (21.2)	9 (14.8)	12 (31.6)	0.000
Treatment, n (%)				
ARV treatment	97 (97.9)	61 (100)	36 (94.7)	0.15
Days from illness onset to ARV start	6 (4-8.5)	6 (4-9.5)	5.5 (3-7.8)	0.15
Two ARV combination therapy	81 (81.8)	50 (82)	31 (81.6)	0.96
Corticosteroid treatment	77 (77.8)	47 (77)	30 (78.9)	0.82
Days from illness onset to corticosteroid start	8 (6-10)	8 (6-10)	7.5 (5-10)	0.35
Initial dosage of corticosteroid, mg/d	60 (40-80)	40 (40-80)	60 (40-80)	0.47
Antibiotic treatment	49 (49.5)	27 (44.3)	22 (57.9)	0.19
Days from illness onset to antibiotic start	9 (6-13)	9 (6-14)	9.5 (5.5-11)	0.29
Intravenous immunoglobulin	43 (43.4)	19 (31.1)	24 (63.2)	0.002
Days from illness onset to immunoglobulin use	8 (6-11)	8 (5-9)	8 (6-12)	0.44
Severity score and outcome				
APACHE II	6 (3-11)	5 (2-8)	9.5 (5-15)	0.000
ICU admission	30 (30.3)	10 (16.4)	20 (52.6)	0.000
ICU length of stay, d	7.5 (4-11)	4 (3-5.8)	8.5 (6.3-11)	0.000

Mechanical ventilation	12 (12.1)	2 (3.3)	10 (26.3)	0.001
ECMO	6 (6.1)	0	6 (15.8)	0.000

Abbreviations: ARV, antiviral treatment; APACHE II, Acute Physiology and Chronic Health Evaluation II; ICU,

Intensive Care Unit; ECOMO, extracorporeal membrane oxygenation.

Accepted Manuscript

Table 3 Multivariable Analyses of Factors Associated With Duration of SARS-COV-2 RNA Detection in 99 Hospitalized Patients.

Variable	Unadjusted HR (95% CI)	P value	Adjusted HR (95% CI)	P ^a Value
Age	0.99 (0.97-1.00)	0.067	0.99 (0.97-1.00)	0.098
Male sex	0.56 (0.34-0.94)	0.029	0.58 (0.35-0.98)	0.042
Current smoking	0.98 (0.46-2.07)	0.982	1.48 (0.65-3.35)	0.345
Hypertension	0.58 (0.33-1.02)	0.058	0.63 (0.36-1.10)	0.104
Diabetes	0.87 (0.41-1.82)	0.865	1.18 (0.53-2.62)	0.680
Days from illness onset to ARV start	0.96 (0.90-1.02)	0.221	0.95 (0.89-1.01)	0.105
Two ARV combination therapy	0.72 (0.38-1.39)	0.331	0.90 (0.45-1.81)	0.899
Corticosteroid treatment	0.85 (0.46-1.57)	0.599	1.00 (0.53-1.89)	0.990
Days from illness onset to corticosteroid start	0.94 (0.87-1.02)	0.127	0.94 (0.86-1.02)	0.121
Immunoglobulin use	0.38 (0.22-0.66)	0.001	0.42 (0.24-0.76)	0.004
Days from illness onset to immunoglobulin start	0.91 (0.81-1.03)	0.136	0.93 (0.82-1.05)	0.231
Antibiotic treatment	0.61 (0.37-1.03)	0.614	0.70 (0.42-1.19)	0.187
APACHE II score	0.89 (0.85-0.95)	0.000	0.89 (0.84-0.96)	0.002
Lymphocyte count	1.86 (1.16-2.99)	0.010	1.81 (1.05-3.10)	0.033

^aResults from the time-dependent Cox proportional hazards model. A hazard ratio (HR) of <1 indicates that the variable increases the duration of SARS-CoV-2 shedding. HRs in multivariable analyses were adjusted for age and sex.