

Committee: Infectious

Title: National Surveillance for Paralytic Poliomyelitis and Nonparalytic Poliovirus Infection

I. Statement of the Problem

CSTE position statement 07-EC-02 recognized the need to develop an official list of nationally notifiable conditions and a standardized reporting definition for each condition on the official list. The position statement also specified that each definition had to comply with American Health Information Community recommended standards to support “automated case reporting from electronic health records or other clinical care information systems.” In July 2008, CSTE identified sixty-eight conditions warranting inclusion on the official list, each of which now requires a standardized reporting definition.

II. Background and Justification

Poliomyelitis is characterized by the acute onset of flaccid paralysis caused by one of the 3 types of poliovirus. The paralysis is typically asymmetric, most often affecting the lower limbs. The great majority of poliovirus infections are asymptomatic or subclinical. Fewer than 1% are paralytic. The onset of paralysis is rapid and does not progress after 3 days. Transmission is primarily by the fecal-oral route. Vaccine-associated paralytic poliomyelitis can occur in both oral poliovirus vaccine recipients and their contacts. Vaccine-derived polioviruses may circulate and cause disease. Starting in 1988, the World Health Organization targeted poliomyelitis for eradication. Polio is currently endemic in only 6 countries in Africa and South Asia. Because of the risks associated with oral poliovirus vaccine, the US and most industrialized countries are now using only inactivated poliovirus vaccine. Non-immune persons traveling outside industrialized countries are at risk of being infected with either wild-type or vaccine-derived polioviruses. Until polio is eradicated and oral poliovirus vaccine use has been stopped worldwide, there is a risk that imported polioviruses may circulate and cause disease in populations with low immunization coverage.

Wild poliovirus transmission has been eliminated from the Americas since 1991. For this reason, two conditions are nationally notifiable: “paralytic poliomyelitis,” in which clinical criteria for the disease have been met; and “non-paralytic poliovirus infection,” in which the case-patient is asymptomatic or mildly ill without paralysis. Furthermore, because oral polio vaccine has not been recommended in the United States since 2000 and has been unavailable in the United States since about 2002, and because vaccine-derived polioviruses can circulate in unvaccinated communities and revert to wild poliovirus phenotype, suspected cases of paralytic poliomyelitis or non-paralytic poliovirus infection should be reported and notified immediately, regardless of whether the virus is suspected to be wild poliovirus or vaccine-derived poliovirus.

Paralytic poliomyelitis and non-paralytic poliovirus infection meet the definition of a nationally and **immediately** notifiable condition—as specified in CSTE position statement 08-EC-02—for the following reason(s):

- The conditions have special importance for international health regulations (IHR)¹.
- The conditions or diseases have been declared eliminated (absence of endemic disease transmission) in the United States.
- A majority of state and territorial jurisdictions—or jurisdictions comprising a majority of the US population—have laws or regulations requiring **immediate** reporting of the conditions to public health authorities; the Centers for Disease Control and Prevention (CDC) requests **immediate** notification of the conditions; and the CDC has condition-specific policies and practices concerning its response to and use of notifications.

III. Statement of the desired action(s) to be taken

CSTE requests that states adopt this standardized reporting definition and CDC adopt these standard notification criteria for paralytic poliomyelitis and nonparalytic poliovirus infection to facilitate more timely, complete, and standardized local and national reporting of this condition.

IV. Goals of Surveillance

To rapidly identify and contain any importations into the United States of wild or vaccine-derived poliovirus.

V. Methods for Surveillance

Surveillance for paralytic poliomyelitis and nonparalytic poliovirus infection should use the sources of data and the extent of coverage listed in Table V.

Table V. Recommended sources of data and extent of coverage for ascertaining cases of Paralytic Poliomyelitis and Nonparalytic Poliovirus Infection

Source of data for case ascertainment	Coverage	
	Population-wide	Sentinel sites
clinician reporting	x	
laboratory reporting	x	
reporting by other entities (e.g., hospitals, veterinarians, pharmacies)	x	
death certificates	x	
hospital discharge or outpatient records	x	
extracts from electronic medical records	x	
telephone survey		
school-based survey		
other _____		

VI. Criteria for Reporting

Reporting refers to the process of healthcare providers or institutions (e.g., clinicians, clinical laboratories, hospitals) submitting basic information to governmental public health agencies about cases of illness that meet certain reporting requirements or criteria. The purpose of this section is to provide those criteria that should be used by humans and machines to determine whether a specific illness should be reported.²

A. Narrative description of criteria to determine whether a case should be reported to public health authorities

Report any illness to public health authorities that meets any of the following criteria:

For Paralytic Poliomyelitis

- Acute onset of flaccid paralysis in a person
 - for whom any of the indicated lab tests listed in Table VI-B have been ordered; or
 - who has any of the epidemiologic risk factors listed in Table VI-B.

For Nonparalytic Poliovirus Infections

- Any case in a person
 - from whom culture of a clinical specimen for poliovirus has been ordered; or
 - from whom poliovirus has been isolated.

Other recommended reporting procedures

- All cases of paralytic poliomyelitis and nonparalytic poliovirus infection should be reported.
- Reporting should be immediate.
- All suspected cases of paralytic poliomyelitis should be reviewed by a panel of expert consultants before final classification occurs.

B. Table of criteria to be used by machines to determine whether a case should be reported to public health authorities

Table VI-B. Proposed Table of criteria to determine whether a case should be reported to public health authorities. Note: The following criteria are proposed for evaluation before general implementation. For purposes of currently implementing reporting the narrative description in VI-A, should be used.

Criterion	Reporting		
<i>Clinical Presentation</i>			
Acute onset, flaccid paralysis	N	N	
Trauma to affected limb			
Spinal cord injury			
<i>Laboratory findings</i>			
Isolation of poliovirus from a clinical specimen			S
Order of a culture for poliovirus	O		S
Order for acute and convalescent serum anti-polio IgG antibodies	O		
<i>Epidemiological risk factors</i>			
Resident of or international travel to country using OPV in past 30 days		O	
Receipt of oral polio vaccine in last 30 days		O	
Contact with person who has received oral polio vaccine in the last 75 days		O	
0 doses of polio vaccine (IPV or OPV)		O	

Notes:

¹ “Human-based” criteria (described below under “A. Narrative”) can be applied by medical care providers and laboratory staff based on clinical judgment and clinical diagnosis. Machine-based

criteria (described below under “B. Table”) can be applied using computerized algorithms that operate in electronic health record systems, including computerized records of laboratory test orders and laboratory test results.

A= This criterion can NOT be present for the case to meet the case definition.

S = This criterion alone is sufficient to report or confirm a case

N = This criterion in conjunction with all other “N” and any “O” criteria in the same column is required to report or confirm a case. A number following an “N” indicates that this criterion is only required for a specific clinical presentation.

O = At least one of these “O” criteria in each category (e.g., clinical presentation and laboratory findings)—in conjunction with all other “N” criteria in the same column—is required to report or confirm a case. A number following an “O” indicates that this criterion is only required for a specific clinical presentation.

* A requisition for any of the “S” or “N” laboratory tests is sufficient to meet the reporting criteria

+ International travel excludes countries that are free of polio and use only inactivated polio vaccine.

C. Disease Specific Data Elements:

Disease-specific data elements to be included in the initial report are listed below.

Clinical Criteria

Paralysis, date of onset

Asymmetric paralysis

Ascending paralysis

Immune deficiency, any

EMG test result

Nerve conduction velocity test results

Epidemiological Risk Factors

Countries visited in last 30 days

Number of OPV doses received

 Date of last OPV dose

Number of IPV doses received

Contact with a person who has received OPV in last 75 days

Contact with a person diagnosed with polio or poliovirus infection

VII. Case Definition for Case Classification

A. Narrative description of criteria to determine whether a case should be classified as confirmed or probable paralytic poliomyelitis or non-paralytic poliovirus infection:

1. Paralytic poliomyelitis

Probable: Acute onset of a flaccid paralysis of one or more limbs with decreased or absent tendon reflexes in the affected limbs, without other apparent cause, and without sensory or cognitive loss.

Confirmed: Acute onset of a flaccid paralysis of one or more limbs with decreased or absent tendon reflexes in the affected limbs, without other apparent cause, and without sensory or cognitive loss; AND in which the patient

- has a neurologic deficit 60 days after onset of initial symptoms; or
- has died; or
- has unknown follow-up status.

2. Nonparalytic poliovirus infection

Confirmed: Any person without symptoms of paralytic poliomyelitis in whom a poliovirus isolate identified in an appropriate clinical specimen (e.g., stool, cerebrospinal fluid, oropharyngeal secretions), with confirmatory typing and sequencing performed by the CDC Poliovirus Laboratory, as needed.

Comment

All suspected cases of paralytic poliomyelitis are reviewed by a panel of expert consultants before final classification occurs. Confirmed cases are then further classified based on epidemiologic and laboratory criteria (11). Only confirmed cases are included in Table I in the *MMWR*. Suspected cases are enumerated in a footnote to the *MMWR* table.

B. Classification Tables

Table VII-B lists the criteria that must be met for a case to be classified as confirmed or probable (presumptive) poliomyelitis or a non-paralytic poliovirus infection.

Table VII-B. Proposed table of criteria to determine whether a case is classified. Note: The following criteria are proposed for evaluation before general implementation. For purposes of current notification, the narrative description in VII-A, should be used.

Criterion	Case Definition			
				Non-paralytic Poliovirus Infection Confirmed
	Paralytic Poliomyelitis			
	Confirmed	Probable		
<i>Clinical Presentation</i>				
Acute onset, flaccid paralysis	N	N	N	A
Decreased tendon reflexes	N	N	N	
Sensory deficit in affected limb(s)	A	A	A	
Cognitive deficit	A	A	A	
Paralysis present 60 days after onset		O		
Death		O		
Follow-up status unknown		O		
Other apparent cause of paralysis (e.g., trauma to affected limb, spinal cord injury)	A	A	A	
<i>Laboratory findings</i>				
Isolation of poliovirus from a clinical specimen				S

Notes:

S = This criterion alone is sufficient to classify a case

N = This criterion in conjunction with all other “N” and any “O” criteria in the same column is required to classify a case.

O = At least one of these “O” criteria in each category in the same column (e.g., clinical presentation and laboratory findings)—in conjunction with all other “N” criteria in the same column—is required to classify a case.

A = This criterion must be absent (i.e., NOT present) for the case to meet the case definition.

* A requisition or order for any of the “S” or “N” laboratory tests is sufficient to meet the reporting criteria. A requisition or order for any of the “O” laboratory tests—in conjunction with

all other “N” criteria in the same column AND at least one of any “O” criteria in the other non-laboratory categories in the same column—is sufficient to meet the reporting criteria.

VIII. Period of Surveillance

Surveillance should be on-going.

IX. Data sharing/release and print criteria

Notification to CDC for confirmed cases of polio is recommended.

- Data reported to NCIRD staff is summarized weekly internally via an NCIRD weekly surveillance report for vaccine preventable diseases. Electronic reports of cases of paralytic poliomyelitis and nonparalytic poliovirus Infection in NNDSS are also summarized weekly in the MMWR Tables. However, because of delays in data entry and data transmission via NNDSS, these two data sources may not correspond. Annual case data on polio is also summarized in the yearly Summary of Notifiable Diseases.
- State-specific compiled data will continue to be published in the weekly and annual MMWR. In addition to those reports, the frequency of reports/feedback to the states and territories will be dependent on the current epidemiologic situation in the country. Frequency of cases, epidemiologic distribution, importation status, transmission risk, and other factors will influence frequency and method of communication and information feedback.
- State-specific compiled data will continue to be published in the weekly reports and annual MMWR Surveillance Summaries. All cases are verified with the state (s) before publication. Data are also included in PAHO and WHO annual reports. The frequency of release of additional publication of this data will be dependent on the current epidemiologic situation in the country. These publications might include annual epidemiologic summaries in the MMWR or manuscripts in peer-reviewed journals.
- As part of an effort to document polio elimination in the Americas, we currently report data on all suspect cases of polio known to NCIRD to PAHO on a weekly basis. Information on sex, age, rash onset, vaccination status, genotype and source (import, import-associated etc.) is provided. No personal identifying or state specific information is re-released to PAHO or WHO.

X. References

Centers for Disease Control and Prevention (CDC). Case definitions for infectious conditions under public health surveillance. MMWR 1997;46(No. RR-10):1–57. Available from: <http://www.cdc.gov/mmwr/>

Centers for Disease Control and Prevention (CDC). National notifiable diseases surveillance system: case definitions. Atlanta: CDC. Available from: <http://www.cdc.gov/ncphi/disss/nndss/casedef/index.htm> Last updated: 2008 Jan 9. Accessed:

Council of State and Territorial Epidemiologists (CSTE). CSTE official list of nationally notifiable conditions. CSTE position statement 07-EC-02. Atlanta: CSTE; June 2007. Available from: <http://www.cste.org>.

Council of State and Territorial epidemiologists (CSTE). Revised Case Definitions for Public Health Surveillance: Infectious Disease 1996-18. Atlanta: CSTE; 1996. Available from <http://www.cste.org>.

Council of State and Territorial Epidemiologists (CSTE). Criteria for inclusion of conditions on CSTE nationally notifiable condition list and for categorization as immediately or routinely notifiable. CSTE position statement 08-EC-02. Atlanta: CSTE; June 2008. Available from: <http://www.cste.org>.

Council of State and Territorial Epidemiologists, Centers for Disease Control and Prevention. CDC-CSTE Intergovernmental Data Release Guidelines Working Group (DRGWG) Report: CDC-ATSDR Data Release Guidelines and Procedures for Re-release of State-Provided Data. Atlanta: CSTE; 2005. Available from: <http://www.cste.org/pdffiles/2005/drgwgreport.pdf> or <http://www.cdc.gov/od/foia/policies/drgwg.pdf>.

Heymann DL, editor. Control of communicable diseases manual. 18th edition. Washington: American Public Health Association; 2004.

Modlin JF. Poliovirus. In: Mandell GL, Bennett JE, Dolin R, editors. Principles and Practice of Infectious Diseases, 6th edition. Philadelphia: Churchill Livingstone; 2005.

XI. Coordination:

Agencies for Response:

- (1) Thomas R Frieden, MD, MPH
Director
Centers for Disease Control and Prevention
1600 Clifton Road, NE
Atlanta GA 30333
(404) 639-7000
txf2@cdc.gov

XII. Submitting Author:

- (1) Paul Cieslak, MD
Manager, Acute and Communicable Disease Prevention
Oregon Department of Human Services
800 NE Oregon St./Suite 772
Portland, OR 97232
(971) 673-1111
paul.r.cieslak@state.or.us

Co-Authors:

- (1) Associate Member
Harry F. Hull, Medical Epidemiologist
HF Hull & Associates, LLC
1140 St. Dennis Court
Saint Paul, MN 55116
(651) 695-8114
hullhf@msn.com
- (2) Associate Member
Cecil Lynch, Medical Informaticist
OntoReason
7292 Shady Woods Circle
Midvale, UT 84047
(916) 412.5504
clynch@ontoreason.com
- (3) Associate Member
R. Gibson Parrish, Medical Epidemiologist
P.O. Box 197
480 Bayley Hazen Road
Peacham, VT 05862
(802) 592-3357
gib.parrish@gmail.com