

**Table 10a. Numbers of Deaths from or with Pneumoconiosis, Ages 15 Years and Older, by State and U.S., 2003**

|                                      | CA  | CT | FL  | KY | LA | MA | ME | MI | NC | NJ | NM | NY | OK | OR | WA | WI | US    |
|--------------------------------------|-----|----|-----|----|----|----|----|----|----|----|----|----|----|----|----|----|-------|
| All pneumoconiosis                   | 138 | 22 | 101 | 67 | 45 | 36 | 12 | 45 | 62 | 81 | 13 | 73 | 12 | 14 | 71 | 32 | 2,639 |
| Coal workers' pneumoconiosis         | <5  | 0  | 17  | 49 | <5 | 0  | <5 | <5 | <5 | <5 | 5  | <5 | <5 | <5 | <5 | <5 | 773   |
| Asbestosis                           | 127 | 22 | 75  | 6  | 37 | 35 | 10 | 35 | 49 | 72 | <5 | 61 | 9  | 13 | 70 | 22 | 1,471 |
| Silicosis                            | 6   | <5 | 8   | <5 | <5 | 0  | <5 | <5 | <5 | 5  | <5 | 6  | <5 | <5 | <5 | 8  | 179   |
| Other and unspecified pneumoconiosis | <5  | 0  | <5  | 10 | <5 | <5 | <5 | <5 | 6  | <5 | <5 | <5 | <5 | <5 | <5 | 0  | 241   |


The sum of particular types of pneumoconiosis deaths may be greater than the total because more than one type could be listed for an individual death.

**Table 10b. Age-Standardized Mortality Rates from or with Select Types of Pneumoconiosis, Ages 15 Years and Older, by State and U.S., 2003**

|                                      | CA  | CT | FL  | KY   | LA | MA | ME | MI | NC  | NJ  | NM  | NY  | OK | OR | WA | WI  | US  |
|--------------------------------------|-----|----|-----|------|----|----|----|----|-----|-----|-----|-----|----|----|----|-----|-----|
| Coal workers' pneumoconiosis         | —   | —  | 0.9 | 15.4 | —  | —  | —  | —  | —   | —   | 3.8 | —   | —  | —  | —  | —   | 3.4 |
| Silicosis                            | 0.3 | —  | 0.3 | —    | —  | —  | —  | —  | —   | 0.7 | —   | 0.4 | —  | —  | —  | 1.8 | 0.8 |
| Other and unspecified pneumoconiosis | —   | —  | —   | 3.1  | —  | —  | —  | —  | 1.0 | —   | —   | —   | —  | —  | —  | —   | 1.1 |

Rates are expressed as deaths per one million residents.

NOTE: Dashes indicate that rates could not be calculated for pneumoconiosis types with less than five deaths.

**Figure 10. Age-Standardized Mortality Rates from or with Total Pneumoconiosis and Asbestosis, Ages 15 Years and Older, by State and U.S., 2003**

Data Sources: Numbers of deaths: State Vital Records. Numbers of deaths in the U.S.: [National Center for Health Statistics](https://www.cdc.gov/nchs/data/tables/multiple-cause-of-death-file) multiple cause of death file.

Population statistics used to calculate rates: [U.S. Census Bureau](https://www.census.gov).