

Table 10a. Numbers of Deaths from or with Pneumoconiosis Ages 15 Years and Older, by State and U.S., 2005

	CA	CT	FL	IA	KY	LA	MA	ME	MI	NJ	NM	NY	OK	OR	TX	WA	WI	US
All pneumoconiosis	102	21	116	11	69	40	37	18	56	85	17	94	14	14	138	65	23	2430
Coal workers' pneumoconiosis	<5	<5	17	<5	47	<5	<5	<5	8	<5	7	<5	<5	<5	<5	<5	<5	653
Asbestosis	89	<5	91	5	9	32	34	18	38	75	5	83	8	12	127	62	14	1423
Silicosis	8	<5	<5	<5	<5	<5	<5	<5	7	<5	<5	7	<5	<5	7	<5	10	161
Other and unspecified pneumoconiosis	<5	<5	6	<5	10	<5	<5	<5	<5	<5	<5	<5	<5	<5	<5	<5	<5	227

The sum of particular types of pneumoconiosis deaths may be greater than the total because more than one type could be listed for an individual death.

Table 10b. Age-Standardized Mortality Rate from or with Pneumoconiosis Ages 15 Years and Older, by State and U.S., 2005

	CA	CT	FL	IA	KY	LA	MA	ME	MI	NJ	NM	NY	OK	OR	TX	WA	WI	US
Coal workers' pneumoconiosis	N/A	N/A	0.8	N/A	14.3	N/A	N/A	N/A	0.9	N/A	4.7	N/A	N/A	N/A	N/A	N/A	N/A	2.7
Silicosis	0.3	N/A	N/A	N/A	N/A	N/A	N/A	N/A	0.9	N/A	N/A	0.5	N/A	N/A	0.5	N/A	2.2	0.7
Other and unspecified pneumoconiosis	N/A	N/A	0.3	N/A	3.0	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	0.9

Rates are expressed as deaths per one million residents.

"N/A" indicates that rates could not be calculated for pneumoconiosis types with less than five deaths.

Data Sources: Number of deaths: State Vital Records. Numbers of deaths in the U.S.: National Center for Health Statistics multiple cause of death file. Population statistics used to calculate rates: U.S. Census Bureau.

Figure 10. Age-Standardized Mortality Rate from or with Total Pneumoconiosis and Asbestosis Ages 15 and Older, by State and U.S., 2005