

**Table 10a. Numbers of Deaths<sup>1</sup> from or with Pneumoconiosis by State and U.S., 2000**

|                                      | CA  | CT | MA | ME | MI | NC | NE | NJ | NM <sup>1</sup> | NY | OR | WA | WI | US    |
|--------------------------------------|-----|----|----|----|----|----|----|----|-----------------|----|----|----|----|-------|
| All pneumoconiosis                   | 143 | 24 | 42 | <5 | 41 | 63 | 5  | 98 | 16              | 79 | 10 | 70 | 21 | 2,864 |
| Coal workers' pneumoconiosis deaths  | 8   | <5 | 0  | 0  | 9  | 9  | 0  | <5 | 6               | <5 | <5 | <5 | <5 | 950   |
| Asbestosis                           | 122 | 20 | 40 | <5 | 25 | 41 | 5  | 88 | 6               | 61 | 9  | 68 | 12 | 1,493 |
| Silicosis                            | 8   | <5 | <5 | 0  | 8  | 6  | 0  | <5 | <5              | 9  | <5 | <5 | 6  | 152   |
| Other and unspecified pneumoconiosis | 5   | <5 | <5 | 0  | 0  | 8  | 0  | 6  | <5              | 5  | <5 | <5 | <5 | 307   |

<sup>1</sup>The sum of particular types of pneumoconiosis deaths may be greater than the total because more than one type could be listed for an individual death..

**Table 10b. Age-Standardized Mortality Rate<sup>1</sup> from or with Selected Pneumoconiosis by State and U.S., 2000**

|                                      | CA  | CT              | MA | MI  | NC  | NE | NJ  | NM  | NY  | OR | WA | WI  | US  |
|--------------------------------------|-----|-----------------|----|-----|-----|----|-----|-----|-----|----|----|-----|-----|
| Coal workers' pneumoconiosis         | 0.3 | -- <sup>2</sup> |    | 1.2 | 1.6 |    |     | 4.8 |     |    |    |     | 4.4 |
| Silicosis                            | 0.3 |                 |    | 1.1 | 1.0 |    |     |     | 0.6 |    |    | 1.4 | 0.7 |
| Other and unspecified pneumoconiosis | 0.2 |                 |    |     | 1.4 |    | 0.9 |     | 0.3 |    |    |     | 1.4 |

<sup>1</sup>Rates are expressed as deaths per one million residents

<sup>2</sup> NOTE: Dashes indicate that mortality rates were not calculated for states with less than five deaths.

**Figure 10. Age-Standardized Mortality Rate from or with Total Pneumoconiosis and Asbestosis by State and U.S., 2000**