

Table 10a. Numbers of Deaths from or with Pneumoconiosis, Ages 15 Years and Older, by State and U.S., 2000

	CA	CT	MA	ME	MI	NC	NE	NJ	NM ¹	NY	OR	WA	WI	US
All pneumoconiosis	143	24	42	<5	41	63	5	98	16	79	31	70	21	2,864
Coal workers' pneumoconiosis deaths	8	<5	0	0	9	9	0	<5	6	<5	<5	<5	<5	950
Asbestosis	122	20	40	<5	25	41	5	88	6	61	28	68	12	1,493
Silicosis	8	<5	<5	0	8	6	0	<5	<5	9	<5	<5	6	152
Other and unspecified pneumoconiosis	5	<5	<5	0	0	8	0	6	<5	5	<5	<5	<5	307


The sum of particular types of pneumoconiosis deaths may be greater than the total because more than one type could be listed for an individual death.

Table 10b. Age-Standardized Mortality Rates from or with Selected Pneumoconiosis, Ages 15 Years and Older, by State and U.S., 2000

	CA	CT	MA	ME	MI	NC	NE	NJ	NM	NY	OR	WA	WI	US
Coal workers' pneumoconiosis	0.3	--	--	--	1.2	1.6	--	--	4.8	--	--	--	--	4.4
Silicosis	0.3	--	--	--	1.1	1.0	--	--	--	0.6	--	--	1.4	0.7
Other and unspecified pneumoconiosis	0.2	--	--	--	--	1.4	--	0.9	--	0.3	--	--	--	1.4

Rates are expressed as deaths per one million residents

NOTE: Dashes indicate that mortality rates were not calculated for pneumoconiosis types with less than five deaths.

Figure 10. Age-Standardized Mortality Rate from or with Total Pneumoconiosis and Asbestosis by State and U.S., 2000

Rates are not illustrated for Maine because it had less than five total pneumoconiosis deaths.

Data Sources: Numbers of deaths: State Vital Records. Numbers of deaths in the U.S.: [National Center for Health Statistics](#) multiple cause of death file. Population statistics used to calculate rates: [U.S. Census Bureau](#)