

Influenza
Virologic Surveillance
Right Size Roadmap

DRAFT 3

Distributed to reviewers: April 29, 2013

FOR REVIEW ONLY, NOT FOR BROAD DISTRIBUTION.

Contents

Forward (<i>to be added</i>)	3
Introduction	3
Virologic Surveillance Requirements	6
How to Use the Roadmap	9
Virologic Surveillance Roadmap.....	11
Objectives: Thresholds and Representativeness	11
Requirements.....	16
Sampling.....	16
Laboratory Testing	21
Data Management	25
Partnerships and Communication	29
Quality Management Systems	32
Surge Capacity for Influenza Surveillance, Novel Event Investigation and Outbreak Events.....	33
Financial Resources.....	36
Implementation Guidance	40
Sampling.....	41
Laboratory Testing	55
Data Management	64
Partnerships & Communication.....	67
Quality Management System.....	71
Surge Capacity for Influenza Surveillance, Novel Event Investigation and Outbreak Events.....	74
Financial Resources.....	76
Appendices.....	80
Appendix A: Surveillance Sampling Process Map	81
Appendix B: Sample Size Calculator Instructions for Use	82
Appendix C: Laboratory methods	83
Appendix D: Funding Fact Sheet Template.....	85
Appendix E: Glossary of Terms and Acronyms	87
Appendix F: Additional Resources and References.....	90
References	92

Forward *(to be added)*

Promote importance of the recommendations, describe the iterative and inclusive process of development, and establish this as 1st Edition so users recognize there are still opportunities for improvements.

Introduction

How much surveillance is really needed? Do we need more or less laboratory testing? How do we know the surveillance data we have is an accurate picture of what's really happening? These are frequent questions of public health decision makers in times of fiscal constraints, and escalate when the threat of novel viruses with pandemic potential seems imminent. The 2009 H1N1 events heightened awareness of these issues, demonstrating the need for a more strategic and scientific approach to virologic surveillance.

Public health laboratories (PHL's) and the Centers for Disease Control (CDC) serve as the backbone of state and national virologic surveillance programs. The amount of virologic surveillance testing performed both at CDC and in PHLs has largely been determined by the capacity of the laboratory.

The CDC-APHL Influenza Virologic Surveillance Right-Size project was launched in 2010 to systematically define the rationale, vital capabilities, and optimal "right-size" for influenza virologic surveillance. The resulting Roadmap consolidates requirements for all components of virologic surveillance in one document, and provides tools to assess and improve the precision of the system to support disease surveillance, response and control efforts and policy decisions. The requirements provide scientific justification for program and laboratory resources to support virologic surveillance policy decisions. Implementation of the right-size virologic surveillance guidelines will assist CDC and PHLs maximize available resources, redirect and build new capacity as needed for optimal surveillance.

Benefits of Right Sizing Influenza Virologic Surveillance:

- *Efficiency*
- *Standardization*
- *Data confidence*

Background

A comprehensive system for influenza surveillance is important to confirm when and where influenza viruses are circulating each year and identify changes in the circulating viruses which may impact vaccine or treatment decisions, or signal the emergence of a new virus with pandemic potential.

In the U.S., the influenza surveillance system is a collaborative effort between the Centers for Disease Control and Prevention (CDC) and its many partners in state, local, and territorial health departments, public health and clinical laboratories, vital statistics offices, healthcare providers, clinics, and emergency departments. The goals for national influenza surveillance include:

- Detect the onset, duration and spread of influenza activity in a geographic area.
- Measure and describe the severity of influenza during a season.
- Determine the populations affected and identify special risk groups.

Requirements

- Monitor the prevalence of circulating virus types and subtypes and match to annual vaccine strains.
- Monitor genetic, phenotypic, and functional changes to circulating influenza viruses and evaluate their potential risk to public health and the need for changes to the annual vaccine composition.
- Identify and monitor novel subtypes that might signal a pandemic.
- Provide data to guide interventions in clinical, individual, and public health control measures.
- Provide information to key partners, including policy makers, emergency response officials, the public and media, and clinical decision makers.

“This is a lot of information that comes from a lot of different people — physicians, people at state health departments and state labs and in hospitals and vital statistics offices,” Brammer said. “Sometimes you step back and look at it and think it’s pretty amazing that this system keeps running week after week, and it always does.”

--Lynnette Brammer, The Washington Post, March 11, 2013 (Service, 2013)

Virologic surveillance is a key and complex component of the influenza surveillance system, informed by a variety of independent but related elements. Specific objectives of virologic surveillance include seasonal situational awareness and determination of virus strain prevalence, early detection of novel viruses or novel events, annual vaccine strain selection, and antiviral resistance monitoring. The 2011 Right Size Influenza Virologic Surveillance Landscape survey provides the most recent and comprehensive summary of influenza testing and surveillance practices employed at both state and local public health entities (APHL, 2012).

All state, and many local, public health laboratories are participants in the U.S. World Health Organization (WHO) Influenza Collaborating Laboratories Network. Influenza virologic surveillance is an essential function of all state health departments and requires a partnership between the state public health laboratory, epidemiologists and influenza surveillance coordinators.

At a minimum, virologic surveillance includes the ability to:

- Access a representative sample of clinical specimens from Outpatient Influenza-like Illness Surveillance Network (ILINet) providers, other clinical primary care sources and clinical laboratories.
- Detect, type and subtype influenza viruses from clinical specimens in a timely manner using standard laboratory methods.
- Report results to providers, epidemiologists and CDC using standard electronic data systems.
- Rapidly refer unsubtypeable influenza viruses to CDC to identify or rule out novel viruses.
- Routinely refer a subset of specimens and viruses to CDC or a CDC-designated laboratory for genetic and antigenic characterization.

Requirements

- Maintain the expertise, warm base (a minimum level of readiness or capacity), and surge capabilities necessary for pandemic response.

This document is a “road map” to achieving an effective virologic surveillance system; it describes the system requirements, and provides options and tools for decision-making processes and system implementation. **In this context, a requirement is an essential component of virologic surveillance that is needed to produce reliable results to achieve state and national surveillance goals. Requirements can be used to focus resources and efficiencies, improve systems approaches, inform policymakers, and provide criteria for funding.** Each state will need to determine how to achieve these goals to meet both national and state needs, including considering options for shared resources.

The Roadmap recommendations were developed over three years based on extensive stakeholder input obtained through meetings, teleconferences, webinars, and a table-top exercise held in December 2012. Stakeholders and exercise participants have identified numerous benefits to implementing the requirements. The approach:

- Standardizes the approach to virologic surveillance and surveillance practices;
- Aids in the development and definition of public health surveillance priorities;
- Provides requirements, resources and statistical calculators to aid in planning and justifying budget and resource requests,
- Increase understanding and support of political leaders and the public;
- Allows epidemiologists and laboratorians to more systematically establish virologic sample-sizes for different surveillance objectives and scenarios based on minimum thresholds of detection and acceptable confidence levels;
- Establishes common language between the laboratorians and epidemiologists, resulting in improved communication between the two groups and better understanding of each other’s needs;
- Provides information to assist decision makers in analyzing the impacts of budget cutbacks on national surveillance objectives (e.g., decreased confidence levels, reduced pandemic preparedness capacity, inability to perform secondary testing such as virus culture).

Moves virologic surveillance from art to science
--Michael Pentella, Ph.D., Right Size Pilot Study Participant

Virologic Surveillance Requirements

The requirements listed here are the essential components needed for effective, efficient and economical influenza

A requirement is an essential component of virologic surveillance that is needed to produce reliable results to achieve state and national surveillance goals. Requirements can be used to focus resources and efficiencies, improve systems approaches, inform policymakers, and provide criteria for funding.

Sampling: Provide year-round access to clinical specimens from ILINet providers and/or other primary care providers and clinical laboratories.

1. Establish a system that ensures efficient collection and timely flow of high quality specimens from the patient management tier of influenza surveillance to the CDC tier throughout the year.
2. Establish a representative network of specimen submitters using ILINet providers and/or other clinical primary care sources. Also collect specimens from hospital/clinical laboratories to ensure that a subset of specimens represents hospitalized patients. Capture unsubtypable influenza positives from clinical and commercial laboratories performing PCR methods that subtype currently circulating viruses.
3. Utilize a statistical, systematic approach to collect an appropriate, adequate number of specimens for testing that will provide reliable data with acceptable confidence limits to meet surveillance objectives and recommended thresholds of detection, including timely detection of novel events. The sampling methodology should limit sampling bias where possible.
4. Utilize sampling approaches that ensure specimens submitted throughout the entire surveillance specimen submission and testing process are representative of:
 - virus types and subtypes
 - the entire year
 - geographic diversity of the population
 - age of ILI patients
 - disease severity
 - targeted populations when necessary for specific investigations
5. Send representative clinical specimens and/or virus isolates to CDC or a CDC-designated laboratory for national surveillance purposes, including annual vaccine virus selection, based on annual CDC criteria and guidance.

Testing: Ensure capability to detect type, subtype and characterize influenza viruses from clinical specimens in a timely manner using reliable laboratory methods.

1. Utilize molecular detection, typing and subtyping methods (e.g., rRT-PCR) for influenza virologic surveillance.

Requirements

2. Maintain instrumentation, personnel, expertise and adequate capacity to test the volume of specimens needed to achieve surveillance objectives.
3. Ensure that staff members are knowledgeable in general principles of virology, molecular biology, and surveillance, as well as appropriate specimen collection, handling, and transport methods.
4. Notify CDC immediately and ship unsubtypeable influenza A viruses to CDC within 24 hours of detection to rule-out novel viruses.
5. Routinely refer a representative subset of specimens and viruses to CDC or CDC-designated laboratory for genetic and antigenic characterization.
6. Maintain capability to rapidly adopt new molecular test methods or test modifications if a new influenza virus with pandemic potential emerges or when new technology provides improvements to virologic surveillance.
7. Maintain additional influenza testing capabilities (as defined in this document) as appropriate for your jurisdiction, or utilize shared testing services models to ensure access to testing.

Data Management: Report results to providers, epidemiologists and CDC.

1. Use electronic data systems that provide data in real time and comply with national standards (HL7, SNOMED, LOINC).
2. All data submitted should provide:
 - Specimen identifier and unique patient identifier
 - The state where specimen was collected
 - Date of Birth of patient and/or Age with unit (years, weeks, months, days)
 - Specimen collection date
 - Specimen Received Date
 - Test method performed
 - Test result
3. Laboratories that have established PHLIP capability should also provide the following data elements, if available:
 - Vaccination status
 - Antiviral treatment
 - Travel information
 - Patient death information
 - Additional geographic information (e.g., county, city, zip)
 - Patient location at time of testing (inpatient, outpatient, long-term care facility)
 - Whether specimen was related to an outbreak
 - Whether specimen was sent to CDC and if so, include specimen identifier
 - Date of illness onset
 - Submitter Information
 - Provider Identifier for the CDC Program (i.e., ILINet provider, EIP, other)
4. States should consider incorporating data from rapid test sites and/or clinical laboratories to supplement influenza surveillance data.

Partnerships and Communication

1. Establish and maintain partnerships and networks enabling communications that support routine surveillance and emergency preparedness and response, data sharing, and specimen sharing. Several interrelated partnerships are needed among the public health and healthcare communities for routine surveillance including:
 - CDC
 - State Epidemiologist/Surveillance Coordinator
 - Public Health Laboratory (PHL)
 - Clinical Laboratories
 - Commercial Laboratories
 - Clinicians
 - Rapid Influenza Diagnostic Testing (RIDT) sites

Quality Systems

1. Establish performance metrics, monitor performance and make improvements as needed to ensure national surveillance requirements are being met in an effective and efficient manner.

Surge

1. Maintain a year-round virologic surveillance system that is flexible and scalable for rapid, effective response to support diagnostic needs and case counts in novel event investigations, and enhance surveillance for outbreak and pandemic scenarios, and has criteria to determine when to scale up and ramp down.
2. Incorporate the role and resource needs of the public health laboratory in the state pandemic plan. PH laboratory representatives should be part of state pandemic planning processes.
3. Develop and maintain a laboratory pandemic surge plan that addresses criteria for specimen triage, algorithm changes to improve throughput, and resource needs (e.g., staff, equipment, space, reagents and supplies).

Financial Resources

1. State influenza surveillance programs and public health laboratories should have adequate funding to support virologic surveillance requirements.
2. State influenza surveillance programs and public health laboratories should coordinate planning and allocation of available funds (ELC, PHEP, EIP, state) to program and laboratory elements (staff, information technology, all supplies, reagents, and equipment maintenance).
3. National, state, and local programs and public health laboratories should have effective cost accounting practices to justify resource needs and efficiently allocate available funds.
4. CDC should have adequate funding to support CDC's national virologic surveillance activities as well as state/local surveillance activities that rely on federal funds.
5. CDC should collaborate internally with cross-cutting CDC programs that provide funding to support other state and local programs, such as ELC and PHEP, to ensure that changes in one program do not unintentionally impact other individual programs.

How to Use the Roadmap

The success of the influenza virologic surveillance system in any jurisdiction requires a strong partnership and collaboration between epidemiology and the public health laboratory, as well as active support of leadership and policy makers. The infrastructure, capabilities, and surveillance system of each state differ, requiring each state to independently evaluate its current surveillance system and determine how to incorporate elements of the Roadmap.

The primary audiences for this Roadmap are the state and local epidemiologists/surveillance coordinators and public health laboratory directors and laboratorians responsible for influenza virologic surveillance. The list of requirements and the descriptions of these essential elements in the Requirements Intent Section will also be useful to policymakers and leadership making resource and funding decisions. Guidance and information in this document will assist each state in identifying strengths and weaknesses in the existing virologic surveillance system, determining the optimal amount of surveillance required and identifying priority implementation activities. The Roadmap will also be a useful tool to assist in crisis management, whether the crisis is the result of detection of a novel virus, a large outbreak, or a crisis of resources due to fiscal constraints. This is not intended to be an SOP (standard operating procedure) manual, but rather a guide, or “roadmap” to assist states in achieving an effective influenza virologic surveillance system.

The most important partnership for effective virologic surveillance is the relationship between the public health laboratory and the epidemiologists/influenza surveillance coordinators. Collaboration to implement these guidelines will be more successful if there is broader understanding of each partner's role.

This document includes three major sections:

1. Virologic Surveillance Objectives: Thresholds and Representativeness which defines the key surveillance objectives, describes specific considerations to ensure that specimens are broadly representative of the population as a whole, and establishes national thresholds for detection. In this context, a threshold is defined as the level (proportion) which triggers some action, which may be as simple as defining a point in the influenza season, or initiating an investigation following detection of a novel virus. The threshold is necessary to “right size” the virologic system – this number is used to calculate the number of specimens that should be tested to ensure adequate confidence levels in surveillance data, and detection of novel viruses at an early enough point where intervention can be effective.
2. Requirements Intent, in which the essential elements for an effective national influenza virologic surveillance system are described, and the rationale for applying these requirements at the state local and national level is explained.
3. Implementation Guidelines, in which suggestions are provided to assist states in operationalizing the requirements. The calculator tools that can be used to estimate the appropriate sample size for key surveillance objectives are described and guidelines for using the on-line calculator tools are provided. The model practices provided in this section are

based on experience with the surveillance system since its inception in the late 1990's, a series of stakeholder meetings, a table-top exercise conducted in December 2012 testing the utility of the roadmap recommendations, and the data gathered through pilot projects conducted in four states during the 2012-2013 influenza season.

Recommended Steps for Implementation

Epidemiology/influenza coordinator and the public health laboratory should:

- Review the document in its entirety to become familiar with the content. Although some sections may seem more relevant to program or laboratory functions, collaboration to implement these guidelines will be more successful if there is broader understanding of each partner's role.
 - Individual sections may stand alone only when considered in context with the [Introduction](#) and the list of all [Requirements](#).
- Identify key partners who should be included in discussions on specific sections or surveillance decisions.
- Convene a meeting (preferably in-person) between program and laboratory staff to address all components of the roadmap document, including use of sample size calculators. Include external partners as needed to address relevant requirements.
- Identify existing practices that meet the roadmap requirements as well as gaps in your virologic surveillance system.
- Use the sample size calculators (or the pre-calculated sample size tables) to assess the reliability of data (confidence levels and error rates) obtained through current (typical) sampling practices and testing volumes.
- Determine which elements or practices will provide the most significant improvement to your system, i.e., the most “bang for your buck”. Consider a staged implementation, rather than an immediate redesign the entire system.
- Identify available funding and resources from all sources. Prioritize capabilities; ensure flexibility and capacity to respond to seasonal variations and emergence of a novel virus.
- Engage public health leaders and policymakers to garner support for implementation.

The pilot study was the first time epi and lab came together to discuss one specific program in detail, in-person. Using the roadmap document forced us to get together and wow, what we learned! A major outcome was learning the importance of being in the same room.

--University of Iowa State Hygienic Laboratory Pilot Study Outcome

Virologic Surveillance Roadmap

Objectives: Thresholds and Representativeness

In order to promote a more statistical, systematic approach to virologic surveillance, thresholds for the following key surveillance objectives have been established. The thresholds are necessary to “right-size” the virologic system – this number is used to calculate the number of specimens that should be tested to ensure adequate confidence in surveillance data as well as detection of novel viruses at a point where intervention can be effective. **In this context, a threshold is defined as the level which triggers some action.** The action may be as simple as defining a point in the influenza season, or initiating an investigation following detection of a novel virus as defined in the CDC’s Interim Guidance on Use of Intervals, Triggers, and Actions for Novel Influenza A (H1N1) Response (ref).

Routine surveillance includes situational awareness, novel event detection and antiviral resistance monitoring. At a minimum, the system should be sized to achieve state level situational awareness needs, and national novel event and antiviral resistance thresholds. Efficiencies in the total number of specimens that need to be tested can be achieved by using a single sampling approach to address as many surveillance objectives as possible. The surveillance program should also have the capability to establish targeted surveillance of specific populations if needed.

1. Situational Awareness: Virologic surveillance provides confirmation of when and where influenza viruses are circulating to inform clinical decision making and public health interventions.
 - a. Surveillance Objective: determine the beginning and end of the influenza season and monitor the prevalence and spread of influenza viruses throughout the year.
 - b. Threshold: 10% prevalence of influenza positive specimens among total ILI specimens submitted to a public health laboratory or the total national system over a two week consecutive time period.

While there is no specific threshold for action, the CDC has traditionally established the start of influenza season at a threshold of 10% positivity, calculated based on positivity of specimens submitted to the PHLs for testing by ILINet and other specimen providers, and the number of pre-screened positive influenza specimens received at PHLs. This value roughly corresponds to the CDC ILINet Seasonal Baseline where the percentage of outpatient visits for influenza like illness reaches 2.2%.

The 10% positivity threshold has been selected for use in the right size situational awareness sample size calculation based on this historical precedent. Calculation of the sample size is made using assumptions regarding medically attended ILI (MA-ILI) based on historical data. State and local surveillance programs may use alternate criteria for declaring the start or end of the influenza season. Additionally, jurisdictions may choose to alter the percent positive used in the sample size calculator to more accurately determine the amount of testing needed throughout the season or assess the confidence level of the data provided.

OBJECTIVES: THRESHOLDS AND REPRESENTATIVENESS

In the past, ILI specimens tested in state public health laboratories were largely unscreened, i.e., not tested by the provider. Today, a significant portion of specimens submitted to public health laboratories may be prescreened positive for influenza by the submitter, (i.e., tested positive using a commercially available influenza test), which can greatly alter the PHL positivity rate. The increased reliance on pre-screened specimens, and the higher sensitivity of PCR methods used more commonly in many clinical laboratories and in all state public health laboratories, may bias the influenza prevalence calculations, impacting the assessment of the scope or severity of the influenza season. Ideally, the percent positivity should be determined using specimens that have not been pre-screened to the greatest extent possible. If data from clinical laboratory testing is being used for situational awareness, at a minimum ensure that the data is coming from sites that are performing high quality testing, and using sensitive methods such as rRT-PCR. Future revisions to this threshold may be needed in the context of changing testing and specimen submission practices.

- c. Representativeness: specimens submitted for routine virologic surveillance to inform community, state and national situational awareness should be broadly representative of the population as a whole (age, geography, risk groups, disease severity).
2. Novel event detection: Virologic surveillance detects the emergence of reassortant, animal origin or completely novel virus subtypes in humans. The initial detection of a novel virus is always laboratory-dependent, and may occur anywhere in the U.S. The sensitivity of the system to detect a novel virus relies on all states contributing at a reasonable level proportionate with their population.
 - a. Surveillance Objective: Detect a novel influenza virus among influenza positive surveillance specimens tested in all states at a low enough threshold for effective intervention and control measures. Note: This objective relates to the initial detection of a novel virus, which generally occurs as part of routine surveillance. Investigation of a novel event (the deep-dive) after initial detection is a separate objective and is discussed in more detail in the Sampling Requirements and Implementation sections.
 - b. Threshold: 0.5% (1/200); one novel virus among 200 influenza virus positive specimens aggregated at the national surveillance level over a one week period. A minimum threshold of 1/165 may be used for determining the sample size in states with limited testing capacity; this approximates the prevalence at which the 2009H1N1 influenza virus was detected. Application of a less sensitive threshold for detection (e.g. 1/150, 1/100) would mean that more novel viruses are circulating prior to detection and impair disease prevention and control efforts.

Using the same detection threshold for identification of novel viruses at a state level (i.e., 1/200 among flu positive specimens tested in the state) would require a significantly larger sample size to achieve an adequate data confidence level. The resources and capacity are generally not adequate to test the number of specimens needed to generate statistically powerful novel event detection data at the state/local or even regional level.
 - c. Representativeness:

OBJECTIVES: THRESHOLDS AND REPRESENTATIVENESS

- i. Routine Surveillance: Novel event detection is a component of routine virologic surveillance, specimens should be broadly representative of the population as a whole (age, geography, risk groups, disease severity).
 - ii. Targeted Surveillance: Detection of a novel virus that emerges within the U.S. may be enhanced with more targeted surveillance in specific populations or risk groups, based on the most current information of risk for novel virus emergence (e.g., returning travelers from high risk areas with ILI).
3. Vaccine Virus Selection: Virologic surveillance provides specimens to CDC for antigenic and genetic characterization to determine whether the circulating strains match the seasonal vaccine strains in “real time”, and to inform annual vaccine virus selection. Submission of specimens should remain consistent throughout the season.
 - a. Thresholds for the degree of difference between circulating viruses and vaccine strains are not defined here, as these criteria are more appropriately established seasonally by the WHO vaccine virus selection experts. Due to seasonal variability in subtype prevalence and the data and virus needs for vaccine virus selection and vaccine candidate development, CDC will provide specific guidance on specimen submission requirements at the beginning of the season, and may adjust submission requirements throughout the season as needed. Every PHL participating in virologic surveillance should submit specimens to CDC (or CDC-designated laboratory) in accordance with annual guidelines.
 - b. Representativeness: Surveillance sampling strategies to ensure appropriate representativeness for vaccine virus selection should prioritize:
 - Timeliness – the most recent viruses.
 - Type and Subtype – viruses representing all circulating types and subtypes. Oversampling of less prevalent subtypes may be necessary to ensure an adequate number of viruses are available for antigenic and molecular characterization and vaccine candidate development.
 - Geographic – CDC should test viruses with sufficient diversity to be representative of the U.S. at a regional level; PHLs should ensure that specimens submitted to CDC are representative of the entire state.
 - Disease Severity –viruses representative of a range of disease severities (from outpatients to fatal cases). .
 - Age - age representativeness is not an important factor for vaccine virus selection.
4. Antiviral Resistance: Virologic surveillance testing to detect antiviral resistance is performed using molecular methods for detection of resistance markers AND phenotypic functional resistance testing, which requires viable virus. If surge antiviral resistance testing capacity is needed, genotypic testing (i.e., pyrosequencing) would be used to meet testing demand.
 - a. Surveillance Objective: Detect antiviral resistance virus among influenza positive surveillance specimens tested across all states at a low enough threshold for effective intervention and control measures. Currently the majority of antiviral resistance

surveillance testing is focused on oseltamivir and is performed at CDC on the same viruses that are submitted for vaccine virus selection. Some state PHLs perform pyrosequencing for molecular markers of antiviral resistance, these results should be reported to CDC for inclusion in national surveillance FluView reports. National “percent resistance” is determined using all sources of data.

b. Thresholds:

- National threshold: detect oseltamivir resistance at or below 5% prevalence among each influenza A subtype or influenza B positive specimens tested at the national level. Seasonal oversampling of certain subtypes may be necessary if increased transmissibility or increasing numbers of resistant viruses are detected. Detection of antiviral resistance approaching or exceeding defined thresholds will initiate investigations and increased testing. Models may be used to assess sample size needed at other prevalence levels. These recommendations or thresholds may change over time depending on resistance trends, or if new viruses with resistance markers emerge. A sustained increase or an unexplained jump in number of resistant viruses may trigger an investigation and expanded testing. Confirmed, substantial increases in resistance may result in changes to clinical treatment guidance, depending on the overall influenza prevalence, resistant virus prevalence, and geographic/temporal spread. If there is an increase in influenza antiviral resistance outside of the U.S., the right size virologic surveillance thresholds may be lowered, targeted surveillance may be implemented, or additional samples may be tested to increase the confidence and decrease the error in detecting a 5% prevalence of resistant viruses.
- State and local thresholds: Using the same antiviral resistance detection threshold at a regional or state/local level would require a significantly larger sample size to achieve an adequate data confidence level. Although some jurisdictions may wish to report antiviral resistance surveillance data at the local/state level to help inform local provider’s clinical management decisions, the resources and capacity are generally not adequate to test the number of specimens needed to generate statistically powerful antiviral resistance testing data at the state/local or even regional level. State and local laboratories choosing to perform antiviral resistance testing are encouraged to utilize sample size models to assess statistical confidence of prevalence rates generated from PHL testing. All PHLs performing pyrosequencing are required to report test results to CDC in a timely manner to be incorporated into national surveillance data.

c. Representativeness:

All surveillance samples submitted to CDC (or CDC-designated laboratory) for antigenic characterization are tested for antiviral surveillance. Surveillance sampling strategies to ensure appropriate representativeness for monitoring antiviral resistance should prioritize:

- Timeliness – recent specimens provide the most valuable data. Testing early and peak season specimens is especially important to monitor changes in antiviral resistance profiles. (Note: Surveillance testing is generally not sufficiently timely for individual

OBJECTIVES: THRESHOLDS AND REPRESENTATIVENESS

patient treatment decisions. Individual results are not reported. CDC does provide diagnostic testing on a case specific basis. Contact xxx for more information).

- Subtype – viruses representing all circulating subtypes should be tested. Oversampling of certain subtypes may be recommended based on seasonal criteria or emergence of resistant viruses.
- Geographic – sufficient diversity to be representative of the US.
- Disease Severity – submit a mix of both outpatient and inpatient specimens.
- Outbreaks/clusters – will be investigated to evaluate geographic spread and drug exposure.
- Age - Age representativeness is not considered to be an important factor for this surveillance objective.

Requirements

This section describes the essential elements for an effective national influenza virologic surveillance system and explains the rationale for applying these requirements at the state local and national level.

Sampling

Sampling Requirements: Provide year-round access to clinical specimens from ILINet providers and/or other primary care providers and clinical laboratories.

1. Establish a system that ensures efficient collection and timely flow of high quality specimens from the patient management tier of influenza surveillance to the CDC tier throughout the year.
2. Establish a representative network of specimen submitters using ILINet providers and/or other clinical primary care sources. Also collect specimens from hospital/clinical laboratories to ensure that a subset of specimens represents hospitalized patients. Capture unsubtypable influenza positives from clinical and commercial laboratories performing PCR methods that subtype currently circulating viruses.
3. Utilize a statistical, systematic approach to collect an appropriate, adequate number of specimens for testing that will provide reliable data with acceptable confidence limits to meet surveillance objectives and recommended thresholds of detection, including timely detection of novel events. The sampling methodology should limit sampling bias where possible.
4. Utilize sampling approaches that ensure specimens submitted throughout the entire surveillance specimen submission and testing process are representative of:
 - virus types and subtypes
 - the entire year
 - geographic diversity of the population
 - age of ILI patients
 - disease severity
 - targeted populations when necessary for specific investigations
5. Send representative clinical specimens and/or virus isolates to CDC or a CDC-designated laboratory for national surveillance purposes, including annual vaccine virus selection, based on annual CDC criteria and guidance.


Requirement Intent

The primary goals of influenza surveillance are to detect novel events, provide viruses for vaccine strain selection and gain a broad understanding of domestic influenza activity. An adequate number of specimens should be tested to provide reliable data to meet the surveillance objectives at the recommended thresholds of detection previously described. Specimen sampling should be designed to enhance detection of novel events, while at the same time collecting a representative sample of routine

influenza cases for overall situational awareness. Where possible, measures to limit sampling bias should be utilized.

Influenza testing occurs in a variety of settings, including physician office laboratories and primary care centers, hospital and commercial laboratories, and local and state public health laboratories. Human respiratory tract specimens and influenza test results data from all these groups contribute to the domestic U.S. influenza virus surveillance system. This complex virologic surveillance landscape can be organized into five major testing tiers based on where testing is performed, as shown in Figure 1 (and in Appendix A).

Figure 1: Surveillance specimen and data submission process



The five tiers of influenza virus surveillance reflect the temporal flow of specimens and fundamental activities performed within each setting. At each level within the five-tier surveillance system, specimens are collected and tested by varying methods to diagnose influenza disease, monitor virus spread, and characterize virus attributes. Since specimens are primarily obtained in the first tier, where they may or may not be tested, and then passed to subsequent tiers for diagnostic and/or surveillance testing, a sampling process takes place at each transfer point. As subsets of specimens flow from the patient management tier to the CDC tier, the number of specimens declines and testing becomes more complex. The system also contains inherent biases due to the complexity of the funnel effect of the sampling system and the use of different test methods in the different tiers. The successive selection of specimen subsets for testing can impact the overall representativeness of samples that are ultimately used to conduct virologic surveillance and select vaccine candidates. The fact that each state surveillance system may impose distinct sampling criteria introduces unanticipated biases that are not always easily understood, further complicating the aggregation of data. For instance, one state may

request only pre-screened rapid test positive specimens from surveillance partners, another state may request a combination of ILI unscreened and influenza pre-screened positive specimens from surveillance partners, impacting the percent positivity reported by the PHL each week.

Sample size and representativeness criteria should be established for sampling at each point in the system. Consistent compliance with sampling criteria will reduce the complexity of data analysis and interpretation at both state and national levels. Sources of bias should be considered and addressed if possible when selecting specimen providers, selecting test methods, and analyzing and interpreting data.

a. Specimen providers and representativeness

Specimens for routine surveillance during influenza season should be obtained from

- ILINet providers and other clinical primary care sources (Tier 1) who commit to regularly sending a subset of ILI patient specimens that have been randomly selected and are not pre-screened (or if tested, a random mix irrespective of test results) to state or local public health laboratories for testing, and
- clinical laboratories (Tier 2) who submit specimens that have tested positive and negative by PCR based on jurisdictional sampling and sample size criteria. Additionally, a subset of culture positive specimens or virus isolates from clinical laboratories that perform virus isolation should be obtained.

Outside of influenza season, specimen providers and clinical laboratories should send all specimens that test positive by RIDT or PCR to the PHL for confirmation and further characterization, as well as specimens from patients with unusual respiratory illness, travel history or risk of exposure to animal origin viruses.

Feasibility and representativeness are the most important factors to consider when choosing specimen submitters. Recruitment and specimen submission criteria should be established so that specimens obtained from primary care sources and clinical laboratories and submitted throughout the entire testing process (“funnel”) for virologic surveillance are representative of the population as a whole or of specific targeted populations as needed to meet surveillance objectives. More details on representativeness are provided in the [Objectives: Thresholds and Representativeness](#) section. The surveillance program should have the capability to establish targeted surveillance of specific populations if needed. Targeted surveillance (i.e., outbreaks, animal exposure, travelers outside the U.S.) may be useful to answer specific questions, especially if a novel event or new virus is detected.

Every public health laboratory (Tier 3) participating in virologic surveillance is responsible for submitting representative clinical specimens and/or virus isolates to CDC or CDC-designated laboratories for national surveillance purposes, including annual vaccine virus selection. Laboratories should submit specimens in a timely manner based on annual CDC criteria and

guidance. Unsubtypable specimens¹ require immediate action as they may reflect a novel virus with pandemic potential. These specimens are sent immediately to CDC for more comprehensive testing.

b. Sample Size

The number of specimens tested each week by state and local public health laboratories has typically been a function of the number of surveillance partners that participate in collection each week and the testing capacity of the PHL, in contrast to the number of specimens needed to meet the surveillance objectives at the recommended thresholds. To address these issues, statistical sample collection calculators have been created to estimate the quantity of specimens that should be tested to provide data with a defined confidence level for each of the surveillance objectives. These calculators can also be used to determine the confidence level of data derived from a particular sampling of ILI patient specimens.

The sample size calculations are based on population size, desired level of confidence, margin of error, and estimated or known prevalence or threshold for detection. More details on thresholds are provided in the Objectives: Thresholds and Representativeness section. State and local public health laboratories are encouraged to use sample size calculators or pre-calculated sample size tables to achieve a more scientific, statistically based sample size that supports surveillance objectives. Specimen sampling approaches should be established to enhance detection of novel events based on national thresholds, while at the same time collecting a representative sample of routine influenza cases for overall situational awareness at the state level. For many states, the number of samples to be tested for each of these objectives is very similar. For smaller states, however, the number of samples necessary to achieve high confidence in situational awareness state level data will be much higher than the number of samples needed to contribute to national novel event detection thresholds.

Outside of influenza season, achieving statistical confidence will not be possible; the focus of surveillance should shift to obtaining all specimens from clinical sites that have tested positive for influenza, or from patients with unusual respiratory illness or travel history or risk of exposure to animal-origin viruses. The Implementation Guidance section and Appendix B provide more information on using the sample size calculators.

c. Sample quality

Influenza surveillance programs and/or submitting laboratories should ensure proper collection, storage and transport of specimens. Proper specimen collection, handling and transport are critical to assuring the quality of results from any laboratory diagnostic test, including diagnostic testing in support of virologic surveillance. Respiratory specimens should be of high quality and properly collected; specimen providers need to be trained in proper collection technique. Timely

¹ Any influenza positive specimen that cannot be definitively typed and subtyped as a circulating seasonal influenza virus, influenza positive specimens producing non-standard or inconclusive results as defined in the CDC Human Influenza Virus Real-Time RT-PCR Diagnostic Panel Instructions for Use package insert.

and efficient transport of specimens is often quite costly, and must be adequately funded by the public health system for effective surveillance.

d. Bias

The influenza virologic surveillance system contains inherent biases due to the complexity of the sampling and submission selection processes (the funnel effect) of the sampling system and the use of different test methods in the different tiers. Sources of bias should be considered and addressed if possible when selecting specimen providers, selecting test methods, and analyzing and interpreting data.

REQUIREMENTS

Laboratory Testing

Laboratory Testing Requirements: Detect, type, subtype, and characterize influenza viruses from clinical specimens in a timely manner using reliable laboratory methods.

1. Utilize molecular detection and subtyping methods (e.g., rRT-PCR) for influenza virologic surveillance.
2. Maintain instrumentation, personnel, expertise and adequate capacity to test the volume of specimens needed to achieve national, state, and local surveillance objectives.
3. Ensure that staff members are knowledgeable in general principles of virology, molecular biology, and surveillance, as well as appropriate specimen collection, handling, and transport methods.
4. Notify CDC immediately and ship unsubtypeable influenza A viruses to CDC within 24 hours of detection to rule-out novel viruses.
5. Routinely refer a representative subset of specimens (and viruses) to CDC or CDC-designated laboratory for genetic and antigenic characterization.
6. Maintain capability to rapidly adopt new molecular test methods or test modifications if a new influenza virus with pandemic potential emerges or when new technology provides improvements to virologic surveillance.
7. Maintain additional influenza testing capabilities (as defined in this document) as appropriate for your jurisdiction, or utilize shared services models to ensure access to testing.

Requirement Intent

Influenza virologic surveillance, by definition, requires laboratories with the capability and capacity to detect, type, subtype and characterize circulating and emerging viruses. The introduction and widespread adoption of molecular methods has reduced the need to maintain classic virologic capabilities in every PHL. The essential components of laboratory testing described below takes into consideration the role of new technologies, the changing landscape of virology expertise in PHLs, and the expected availability of national, state, and local fiscal resources.

The roadmap classifies virologic testing components based on a stakeholder assessment of requirements that should be maintained and available at all PHLs involved in influenza surveillance (primary testing); or as additional surveillance testing capabilities that may be maintained based on jurisdictional needs and resources, or provided through a shared services model. PHLs engaged in influenza virologic surveillance should utilize molecular methods such as rRT-PCR as their primary testing method due to their superior sensitivity and specificity.

Primary Testing Method:

PHLs performing virologic surveillance are expected to utilize molecular methods, such as rRT-PCR, as the primary method for influenza detection and subtyping. This is an ELC benchmark. Influenza rRT-PCR testing provides rapid, sensitive and accurate detection and identification of influenza viruses for routine influenza surveillance, outbreak detection, and pandemic response. The CDC Human Influenza Virus Real-Time RT-PCR Diagnostic Panel (CDC Flu rRT-PCR Dx Panel) is an FDA-cleared *in vitro* diagnostic assay that is manufactured and distributed by CDC to all state and qualified local PHLs engaged in influenza virologic surveillance testing. Although PHLs have the option to use commercial rRT-PCR assays, there are specific benefits to utilizing the CDC assay and CDC supplied reagents. This is a nationally recognized reference method, and allows for standardization of influenza testing across all PHLs. The assay detects current influenza strains, is continually assessed and updated as needed to detect strain variations, and should detect novel viruses. The assay also allows for higher throughput testing algorithms to support outbreak and pandemic response. This method is consistent with the laboratory workforce's increasing proficiency in molecular testing methods. However, all PHLs engaged in influenza surveillance should have staff knowledgeable in general principles of virology and surveillance, and appropriate specimen collection, handling, and transport methods.

Additional Testing Methods:

There are additional testing methods that are used to support influenza virologic surveillance. These include influenza virus culture, antiviral resistance testing of influenza viruses, influenza hemagglutination inhibition (HAI), immunofluorescence identification of influenza viruses, and serology testing. Each of these methods has distinct purposes, advantages and disadvantages for both national and state surveillance. The determination to use any of these methods in PHLs should be made based on state and jurisdictional needs, and include a cost analysis and identification of sustainable funding sources.

Virus isolation has the advantage of producing quantities of virus sufficient for full antigenic characterization for determining vaccine match, and conducting antiviral resistance testing². However, influenza virus culture is less sensitive and specific than rRT-PCR and there are vast variations in the sensitivity of different cell culture lines, the growth characteristics of influenza virus strains, and PHL practices and expertise. In addition, influenza virus culture is less rapid than influenza rRT-PCR and less adaptable to sudden surges in specimen numbers. Influenza virus culture must be maintained at CDC and a subset of PHLs.

Antiviral resistance testing is necessary to monitor the presence and level of antiviral resistance in circulating influenza viruses. These data inform patient management and treatment recommendations as well as national antiviral stockpile policies. Definitive antiviral resistance testing requires both phenotypic resistance testing done using an NAI assay (requires cultured viruses), and detection of genetic changes (drug resistance markers) by pyrosequencing and/or sequencing. Both of these test methods are available at CDC and a subset of PHLs. Performing pyrosequencing at a subset of PHLs provides a cost-effective and efficient approach to expanded surveillance screening. Supporting a

² Manual for the Surveillance of Vaccine-Preventable Diseases, 5th edition, 2012;
<http://www.cdc.gov/vaccines/pubs/surv-manual/index/html>

limited number of testing sites allows the system and methods to be more readily updated and adapted for viral mutations, limits costs, and allows for effective training to develop extensive expertise required to perform and interpret the test results.

Influenza hemagglutination inhibition (HAI) testing remains a cornerstone of antigenic characterization for influenza vaccine strain selection. CDC maintains the expertise to test influenza virus culture isolates by HAI for antigenic characterization using specialized antisera. HAI tests using the WHO kit reagents that are provided to PHLs are a reliable indicator of influenza type and influenza A subtype, but are not a reliable indicator of influenza strain or strain changes; results of influenza HAI tests using WHO kit reagents may be over-interpreted as indicators of vaccine effectiveness and circulating strains. In addition, influenza HAI testing requires frequent practice to maintain expertise and proficiency in test performance and result interpretation.

Immunofluorescent antibody identification of influenza viruses are used in some PHL's for confirmation and identification of influenza viruses that have grown in cell cultures. Immunofluorescence (IF) testing is not as specific as rRT-PCR testing and requires a fluorescent microscope and staff with specialized expertise. However, influenza IF testing has the potential to identify novel strains of influenza grown in culture that evade detection by rRT-PCR.

Serologic testing is neither a routine surveillance nor diagnostic tool; it is currently used primarily by CDC or academic institutions for research purposes or retrospective seroprevalence studies. Testing requires staff with a high level of expertise and specialized reagents, there are currently no FDA cleared influenza serology tests.

Rapid influenza diagnostic tests (RIDTs) for the detection and identification of influenza viruses are used by many sentinel surveillance providers, other primary care sites, emergency departments as well as physician office and clinical laboratories. However, these tests are significantly less sensitive and specific than molecular assays³. RIDTs may also be less reliable when new virus strains emerge⁴. If PHLs choose to use RIDTs to serve diagnostic needs for selected patient populations, they should be aware of the limitations of these tests (e.g., the considerable variability of the positive and negative predictive values depending upon the prevalence of influenza in the community) and should follow guidelines provided by CDC (currently at www.cdc.gov/flu/professionals/diagnosis/rapidlab.htm). If an important clinical decision will be affected by either a positive or negative rapid test result, the rapid test result should be confirmed by another test, such as rRT-PCR. Even if the PHL does not perform

³ Uyeki TM, Prasad R, Vukotich C, et al. Low sensitivity of rapid diagnostic test for influenza. Clin Infect Dis 2009;48:e89.

CDC. Evaluation of rapid influenza diagnostic tests for detection of novel influenza A (H1N1) virus—United States, 2009. MMWR 2009;58:826–9.

⁴ Balish A, Garten R, Klimov A, Villanueva J. Analytical detection of influenza A(H3N2)v and other A variant viruses from the USA by rapid influenza diagnostic tests. Influenza Other Respi Viruses 2012;September 18 [Epub ahead of print].

RIDTs, virology staff should maintain awareness of the performance characteristics of currently available RIDT's that may be used within their jurisdiction in order to provide seasonal guidance to clinicians, clinical laboratories, and surveillance coordinators. PHLs or surveillance coordinators may also collect data from clinical sites using RIDTs to provide additional seasonal situational awareness data.

Direct specimen immunofluorescence ("DFA") involves testing clinical material taken directly from the patient and using fluorescent-labeled antibodies to detect influenza antigens that may be present. DFA can be performed quickly, but is not as accurate as rRT-PCR testing and requires significant expertise and expensive fluorescent microscopes. In addition, results can be very subjective and are dependent on the individual reader's expertise.

Expanded testing for other respiratory viruses using molecular respiratory virus panels (RVPs) has become common in many clinical laboratories and is increasingly used in PHLs to provide jurisdictional and national information about circulating viruses that are associated with acute respiratory illness. While detection and identification of non-influenza respiratory viruses is not considered an essential component of national influenza virologic surveillance, these assays data can aid in identifying the cause of non-influenza community illnesses or outbreaks. If surveillance for other respiratory viruses is performed to meet jurisdictional needs, it is recommended that PHLs consider adopting molecular RVPs to replace less sensitive viral culture, if resources are available.

REQUIREMENTS

Data Management

Data Management Requirements: Report results to providers, epidemiologists and CDC

1. Use electronic data systems that provide data in real time and comply with national standards (HL7, SNOMED, LOINC).
2. All data submitted should provide:
 - Specimen identifier and unique patient identifier
 - The state where specimen was collected
 - Date of Birth of patient and/or Age with unit (years, weeks, months, days)
 - Specimen collection date
 - Specimen Received Date
 - Test method performed
 - Test result
3. Laboratories that have established PHLIP capability should also provide the following data elements, if available:
 - Vaccination status
 - Antiviral treatment
 - Travel information
 - Patient death information
 - Additional geographic information (ex. county, city, zip)
 - Patient location at time of testing (inpatient, outpatient, long-term care facility)
 - Whether specimen was related to an outbreak
 - Whether specimen was sent to CDC and if so, include specimen identifier
 - Date of illness onset
 - Submitter Information
 - Provider Identifier for the CDC Program (i.e., ILINet provider, EIP, other)
4. States should consider incorporating data from rapid test sites and/or clinical laboratories to supplement influenza surveillance data.

Requirement Intent:


Virologic surveillance in the U.S. relies on a combination of data and specimens: data from laboratory tests performed at U.S. WHO Collaborating Laboratories and National Respiratory and Enteric Virus Surveillance System (NREVSS) laboratories, and specimens from patients with ILI that tested positive for influenza at PHLs and are submitted to CDC for further characterization. There are approximately 85 WHO collaborating laboratories. The WHO collaborating laboratories include 65 state and city/county PHLs supported by CDC, as well as several large tertiary care or academic medical centers⁵. These laboratories have the capability to test for and report the type of influenza virus (A or B) and subtype of

⁵ <http://www.cdc.gov/flu/pdf/weekly/overview.pdf>

influenza A viruses (2009H1, H3, and some novel subtypes), and provide both data and specimens/viruses to CDC.

The NREVSS system includes some PHLs and clinical, commercial and academic medical center laboratories. The NREVSS laboratories that are not WHO collaborating centers provide data on influenza laboratory test results only as “influenza A” and “influenza B”, and those laboratories participating as a WHO collaborating center submit non influenza respiratory pathogen data into NREVSS system according to the 2011 Right Size Virologic Surveillance Survey. The influenza data from the WHO collaborating laboratories and NREVSS play a large role in the weekly national reports generated by CDC in FluView, www.cdc.gov/flu/weekly/.

Additional influenza testing data from rapid influenza diagnostic testing sites and/or other clinical laboratories may be available to states and can help to provide a fuller representation of influenza activity at the local level. These clinical sites have the capability to test for and report the type of influenza (A or B), some also utilize methods that can subtype influenza A (2009H1, H3, H1).



Source: Centers for Disease Control and Prevention Unpublished data.

Currently, influenza surveillance data is obtained both as aggregated data, using web entry methods, and as patient level data, using HL7 electronic laboratory reporting and comma delimited files. This dichotomy makes the data management, aggregation, and linking of virologic and epidemiologic data challenging. However, this spectrum of reporting formats reflects a long history of virologic surveillance with varying technological solutions and varying capabilities of reporters.

The increasing availability of Laboratory Information Management Systems (LIMS) in PHLs makes it possible to establish automated electronic laboratory messaging of influenza test results to other public health entities (e.g., state epidemiology offices, other public health laboratories, and CDC). APHL's Public Health Laboratory Interoperability Project (PHLIP) provides PHLs with an electronic method to report laboratory test results using national standards such as HL7 Version 2 messaging, SNOMED vocabulary, and LOINC codes for laboratory tests. The PHLIP vision is to provide each SPHL with a viable option for electronic transmission of laboratory test data, in order to achieve interoperability between different systems and to exchange information in a useful and meaningful way. The APHL PHLIP effort began in 2008, and the majority of state PHLs are now reporting influenza results electronically using PHLIP.

PHLIP is the preferred reporting mechanism for influenza and is considered a requirement. It is understood that not all SPHLs have the same capabilities or resources to participate in PHLIP at present; however, PHLIP implementation should be the goal for all SPHLs and for county and local PHLs that participate in virologic surveillance. PHLIP offers many advantages, which include:

- Standardized patient level reporting, improving data quality and simplifying data aggregation.
- Individual reports in near "real-time".
- PHLIP messaging is compatible with other national electronic messaging solutions
- PHLIP can be expanded to report laboratory results for other pathogens using the same mechanisms for messaging.
- Laboratory staff time is not required to collect and report laboratory results.
- Additional identifiers for type of specimen submitter (i.e., ILINet provider, EIP site, etc.) can be reported.
- Utilization of PHLIP to facilitate a shared services approach among PHLs (i.e., PHLs specializing in influenza virus culture or antiviral resistance testing).

Electronic data messaging of specimen level data allows for more detailed analysis of the data and fuller understanding of potential biases in the data. Biases may include lack of population representativeness in the selection of patients for testing, prescreening of specimens before submission to public health laboratories and the quality of tests used for initial screening (Refer to [Sampling Section](#) for more information).

Virologic Surveillance Data from Non-Public Health Laboratory Sources

The focus of virologic surveillance has been on data generated by the PHLs participating as WHO-CLs in the U.S. However, influenza testing performed by clinical and commercial diagnostic laboratories may provide useful supplementary data, increasing the overall volume of testing and geographic

representation. Existing virologic data from non-PHL sources, notably NREVSS, and some options for new sources of virologic data are discussed below.

NREVSS is managed by the CDC's Division of Viral Diseases. The system is the main source of national surveillance data for non-influenza respiratory viruses. Influenza laboratory testing data is also collected in NREVSS and shared with the Influenza Division for reporting as part of the national virologic surveillance report. Due to a rapid expansion of the reporting provider network in recent years, there were approximately 500 laboratories enrolled as NREVSS sites that reported influenza in the 2011-12 season. However, concerns about the sustainability of the expansion, along with a desire to maintain the historical number of reporters used in influenza virologic surveillance has resulted in use of only the data from 60 original participants. The reports submitted to NREVSS include laboratory-level information about the following, 1) number of specimens tested for influenza, 2) test method used (i.e., RIDT, culture, PCR), and 3) number of influenza positive specimens by influenza type and if available, by subtype.

Additional sources of influenza (NREVSS-like) clinical/hospital and/or commercial laboratories data can also be utilized and developed. For the same reasons that NREVSS data is useful at the national-level, building a network of non-PHLs within a state is beneficial. These state laboratory networks are frequently used as a source of additional local level data for situational awareness. Data from these sites may be transmitted electronically at the specimen level or in aggregate by a simpler method. Regardless of whether specimen level or aggregate data is received, necessary data elements would include:

- a. Date or week of specimen collection, receipt, or test
- b. Total number of tests performed and influenza positives by:
 - Type
 - Subtype (if available)
 - Age group

REQUIREMENTS

Partnerships and Communication

Partnerships and Communication Requirements

1. Establish and maintain partnerships and networks enabling communications that support routine surveillance data and specimen sharing, and emergency preparedness and response. Several interrelated partnerships are needed among the public health and healthcare communities for routine surveillance including:
 - CDC
 - State Epidemiologist/Surveillance Coordinator
 - Public Health Laboratory (PHL)
 - Clinical Laboratories
 - Commercial Laboratories
 - Clinicians
 - Rapid Influenza Diagnostic Testing (RIDT) sites

Requirement Intent

The U.S. influenza surveillance system, which includes virologic, morbidity and mortality components, relies heavily on partnerships across the local, state and national levels. As shown in Figure 3, these partnerships and networks are critical to communications that support routine surveillance, emergency response, data sharing, and specimen sharing. The role and value of partnerships was very apparent in the highly effective public health response to the 2009 influenza A(H1N1) pandemic and has been documented in APHL's [Lessons from a Virus](#) (APHL, 2011).

The most important partnership for effective virologic surveillance is the relationship between the **public health laboratory staff and epidemiology/Influenza Coordinators**. Data from the 2011 Right Size Influenza Virologic Surveillance Landscape survey to assess influenza-related activities at PHLs highlighted the value of this collaboration for jointly developing surveillance policy, strategies, and resource allocation (APHL, 2012). This partnership also serves to improve communication, education and outreach to specimen submitters, data sharing, and outbreak investigations. The roles and responsibilities of the laboratorians and epidemiology/influenza coordinators will vary across jurisdictions. Therefore, it is important that both parties have an understanding of each other's roles, and agreement on the best approach to address each surveillance component.

Building and maintaining relationships with external partners has been identified as a pivotal contributor to the success of public health surveillance efforts. A strong **PHL/Epidemiology /Clinical-Commercial-Academic laboratory** partnership will support the formation of an effective specimen submitter network, enhance information sharing and outbreak response. Strong relationships among state epidemiology, public health laboratory, and clinical partners are crucial to ensuring quality and consistent data and specimens for influenza virologic surveillance.

Additional key PHL relationships are outlined in several documents, including APHL's "[Core Functions of Public Health Laboratories](#)", [Definition of a State Public Health Laboratory System](#), and CDC's "[Public Health Preparedness Capabilities: National Standards for State and Local Planning](#)" (APHL, 2010; APHL, 2007; CDC, 2011). These relationships have also been included as elements in Public Health Emergency

Response planning. Efforts to create state-based laboratory networks that interconnect to form a cohesive national system have been promoted in the context of APHL's [Lab System Improvement Program \(L-SIP\)](#), [Laboratory Efficiency Initiative \(LEI\)](#), [All-Hazards Public Health Emergency Preparedness \(PHEP\)](#) initiatives, the CDC/CSTE [Competencies for Applied Epidemiologists in Governmental Public Health Agencies \(AECs\)](#) and the [Laboratory Response Network \(LRN\)](#) for more than a decade (APHL, 2013; APHL, 2013; CDC, 2013; CDC, 2012; CDC & CSTE, 2008). The LRN structure for bioterrorism is represented by a pyramid, with clinical ("Sentinel") laboratories as the foundation, PHLs as the primary members of the reference laboratory level, and CDC and other national laboratories at the peak of the structure (see Figure 2). The LRN pyramid demonstrates the interrelatedness of various partners in responding to potential bioterrorism threats and sets a foundation for partnerships and communication for other surveillance systems such as influenza virologic surveillance.

Partnerships between **CDC and PHLs** have also resulted in a number of important collaborative efforts including, but not limited to, informational teleconferences for PHLs, development of a "warm base" of diagnostics capabilities in PHLs for rapid deployment of tests (e.g., 2009 influenza A H1N1) and ongoing reagent and equipment support facilitated by the CDC, APHL and private industry, and others included in Figure 3. Similar relationships exist between **CDC and state based influenza surveillance coordinators**. Monthly conference calls and annual meetings allow for discussions about influenza circulation and potential areas of concern. Annual communications have been established between CDC, PHLs and epidemiology staff to ensure that all stakeholders are receiving relevant information at the beginning of each season, and working collaboratively toward common surveillance goals.

While partnerships between influenza surveillance programs and PHLs have been established to some degree in most states, maintaining these partnerships in the future may present a challenge as state resources dwindle and funding becomes more uncertain. Gaps in effective partnerships can result in significant but often poorly recognized negative impacts on virologic surveillance.





Figure 2. LRN pyramid showing the partnership relations between sentinel, reference, and national laboratories Laboratory Response Network (LRN)

Figure 3. Essential Influenza Virologic Surveillance Partnerships and Communication Web



REQUIREMENTS

Quality Management Systems

Quality Management Systems Requirements

1. Establish performance metrics, monitor performance and make improvements as needed to ensure national (and state/local) surveillance requirements are being met in an effective and efficient manner.

Requirement Intent

A quality management system can be defined as a coordinated activity to monitor and control organizational processes and resources. Both national and state/local virologic surveillance systems require monitoring and management of the various components that inform surveillance. Establishing and applying performance metrics encourages continuous improvement, demonstrates return on investments, and helps to justify funding. It is important that CDC and state/local jurisdictions monitor activities related roadmap requirements as well as compliance with ELC and PHEP benchmarks.

State and local influenza programs and PHLs should monitor quality and consistency of specimen submissions throughout the system, data confidence in relation to sample sizes and representativeness of surveillance data, and laboratory testing quality assurance parameters.

CDC should monitor reporting and specimen submissions to ensure national surveillance data are representative of influenza activity, meet current national needs across the entire country, and that specimens are being submitted in a timely manner throughout the year to help inform annual vaccine virus selection. Additionally, monitoring resource allocation and usage in the context of surveillance test activities allows CDC to identify areas for improvement and justify funding for national and state/local surveillance systems. National quality monitoring efforts may include timeliness and consistency of testing data reported to CDC, influenza virologic surveillance specimen submissions to CDC and CDC-designated laboratories, and utilization rates of Influenza Reagent Resource (IRR) reagents.

CDC, public health laboratories and state/local surveillance programs should use data gathered through quality monitoring practices to identify and implement improvements and efficiencies as appropriate.

REQUIREMENTS

Surge Capacity for Influenza Surveillance, Novel Event Investigation and Outbreak Events

Surge Capacity Requirements

1. Maintain a year-round virologic surveillance system that is flexible and scalable for rapid, effective response to support diagnostic needs and case counts in novel event investigations, and enhance surveillance for outbreak and pandemic scenarios, and has criteria to determine when to scale up and ramp down.
2. Incorporate the role and resource needs of the public health laboratory in the state pandemic plan. PH laboratory representatives should be part of state pandemic planning processes.
3. Develop and maintain a laboratory pandemic surge plan that addresses criteria for specimen triage, algorithm changes to improve throughput, and resource needs (e.g., staff, equipment, space, reagents and supplies).

Requirements Intent

Virologic surveillance is vital to support novel event and outbreak investigations and pandemic response. Pre-event and during an event, communication and coordination between epidemiology and laboratory leadership is essential to develop, refine and change the strategy for virologic surge sampling and testing. However, the term “surge capacity” has many different meanings, which can result in unrealistic expectations of the virologic surveillance system. While seasonal surveillance provides the warm base of expertise and infrastructure necessary to provide surge capacity, the response needed for a local outbreak investigation, emergence of a novel influenza virus, or a pandemic response are qualitatively and quantitatively very different. The Institute of Medicine Medical Surge Capacity Workshop report grossly defines surge capacity as the ability to rapidly accommodate a large number of patients from a defined mass-casualty incident or pandemic, and considers surge capacity on a continuum with three distinct stages: Conventional capacity, Contingency capacity, and Crisis capacity (Institute of Medicine (US) Forum on Medical and Public Health Preparedness for Catastrophic Events, 2010).⁶ These medical surge definitions are adapted here to provide standardized terminology to improve planning and response:

- Conventional capacity: Routine virologic surveillance, capacity to test adequate sample size to produce data with reasonable confidence levels.
- Contingency capacity: Minor adaptations are made that generally have limited impact on routine operations. This “spare” capacity must be maintained to deal with fluctuations in testing that may be necessary during a bad influenza season (e.g., increased hospitalizations, rapid transmission

⁶ Institute of Medicine (US) Forum on Medical and Public Health Preparedness for Catastrophic Events. Medical Surge Capacity: Workshop Summary. Washington (DC): National Academies Press (US); 2010. The Communications Challenge: Definitions, Standards, and Metrics. Available from: <http://www.ncbi.nlm.nih.gov/books/NBK32856/>

within the community, drifted virus), a local outbreak investigation, or a novel event investigation. Efficient use of contingency capacity may require emphasis on targeted testing based on event specific criteria.

- **Crisis capacity:** A fundamental, systematic change into a system in which standards of operation are significantly altered. When crisis capacity is reached, the focus will shift to expanded hours of operation, utilizing staff from other programs or areas of the laboratory, and most importantly significantly limiting testing based on event specific governance criteria.

These definitions of capacity relate to the equipment and supplies available, and even more importantly to the staff needed to provide all the tasks required for specimen accessioning, processing, testing, data management and analysis. Therefore each jurisdiction may have different triggers that will cause a shift from one stage to another.

Novel event/outbreak Investigations:

Following identification of a potential outbreak, a novel virus or other rare event, populations that will be targeted for testing will be determined based on:

- Epidemiologic criteria, e.g., exposure , geographic location, event specific risk factors
- Clinical criteria, e.g., severe or fatal illness
- Specimen sources, e.g., ILINet or other primary care providers, clinical laboratories using high performing assays

Although epidemiologists will serve as gate-keepers for PHL testing, collaborative epidemiology-laboratory pre-event planning and event response is needed to ensure common understanding and expectation of contingency and crisis capacity. PHL influenza testing capacity models (ref) and sample size calculators can be used to identify system efficiencies and limitations. The APHL Emerging Infections Response Framework is another useful tool for planning (ref).

Pandemic Surge:

In the event a novel virus emerges that is highly transmissible, the public health laboratory will likely be the only resource for diagnostic testing at the start of the event, particularly if the commercially available tests do not reliably detect or differentiate the virus. As demonstrated in the 2009 H1N1 pandemic response, PHL testing capacity will be stretched by testing demands, rapidly reaching unsustainable crisis capacity. Effective governance for triage of cases eligible for testing at the public health laboratory will be necessary. Epidemiologists, in collaboration with the PHL leadership, will need to manage the demand for diagnostic testing, and ensure that surveillance testing

H1N1 Response: Lessons of the Virus

Public health labs opened their doors to let in specimens from clinical labs because they had the best test and were eager to fulfill their surveillance mission.....

You can't just hire a molecular biologist off the street. The layers of quality assurance involved in validation add critical steps to the process. Specific, sophisticated instruments and expensive reagents are required. Regulatory clearance can add time to the process. The subtleties don't always translate to the general public or political representatives. For them, the gold-standard is an immediate yes-no answer at the point of care.

that is representative of the relevant populations is prioritized so that effective response and control measures can be effectively implemented.

Even if diagnostic testing demand can be met by the clinical laboratory sector, the PHL will be the primary resource for virologic surveillance data. Therefore, the PHL should be represented in state pandemic planning activities. The role and resource needs of the PHL should be addressed in the state pandemic plan. Expectations for state and local epidemiologists to serve as gate-keepers for specimen testing demands should be coordinated in advance and defined in the plan.

All PHLs should develop and maintain an internal pandemic surge plan that addresses criteria for specimen triage, algorithm changes to improve testing and reporting throughput, and resource needs (e.g., staff, equipment, space, safety, reagents and supplies).

REQUIREMENTS

Financial Resources

Financial Resources Requirements

1. State influenza surveillance programs and public health laboratories should have adequate funding to support virologic surveillance requirements.
2. State influenza surveillance programs and public health laboratories should coordinate planning and allocation of available funds (ELC, PHEP, EIP, state) to program and laboratory elements (staff, information technology, all supplies, reagents, and equipment maintenance).
3. National, state, and local programs and public health laboratories should have effective cost accounting practices to justify resource needs and efficiently allocate available funds.
4. CDC should have adequate funding to support CDC's national virologic surveillance activities as well as state/local surveillance activities that rely on federal funds.
5. CDC should collaborate internally with cross-cutting CDC programs that provide funding to support other state and local programs, such as ELC and PHEP, to ensure that changes in one program do not unintentionally impact other individual programs.

Requirement Intent

An optimal influenza surveillance system requires adequate resources to support all essential elements defined in this roadmap document. Sustaining the national, state and local components of this system is increasingly threatened by the decline in annual and pandemic response funding. Implementation of the right size virologic surveillance guidelines will help CDC, PHLs, and surveillance programs maximize available resources, redirect resources as necessary, and build new capacity as needed for optimal surveillance. Accurate assessments of the cost of virologic surveillance activities are critical to justify and prioritize funding.

Federal funding provided since 2005 to support pandemic planning, and supplemental funding made available during the 2009 H1N1 response have resulted in many improvements to the US influenza surveillance system. It is important to use available resources now to optimize systems for the future. Collaborative planning, grant proposal development and funding allocation between influenza surveillance programs and PHLs is essential to ensure all involved parties have an understanding of the costs associated with all aspects of influenza surveillance and that all virologic surveillance requirements are adequately resourced.

Surveillance is supported by several different funding streams, distributed at different times depending on source. Additionally, the cost of surveillance, and the availability and allocation of funds for the different components of virologic surveillance varies across jurisdictions; these challenges can impact the overall effectiveness of the surveillance system. While funding is often cited as a key limiting factor, the true costs of virologic surveillance are not well defined, likely because of the complexity of the system. Optimizing resources and justifying funding requests will require better cost accounting at the national, state and local level.

Federal Funding Sources

Epidemiology and Laboratory Capacity (ELC) Cooperative Agreement

Currently, state/local influenza virologic surveillance systems rely heavily on CDC funding resources. In particular, the Epidemiology and Laboratory Capacity (ELC) cooperative agreement program has been the primary funding source for surveillance at the state level, especially for supporting programmatic and laboratory personnel. All 50 states and several large cities receive funding from CDC to support US influenza surveillance goals via the ELC Cooperative Agreement program. The primary goals of the influenza component of the ELC program include establishing and supporting ILI sentinel provider networks; providing timely ILINet data to CDC, and maintaining laboratory testing and reporting capability and capacity for year round virologic surveillance. CDC supports public health influenza virologic surveillance supported through the ELC because the work of the PHL's contributes to national and global disease prevention efforts. Specimens submitted to CDC and CDC-designated laboratories contributes the viruses used to assess antigenic changes that impact vaccine effectiveness, these viruses are also frequently selected as seed strains for the manufacturer for seasonal vaccines.

While routine annual influenza surveillance principally relies on ELC and state funding, these federal funding programs also contribute to national virologic surveillance goals:

- **Emerging Infections Program (EIP):** active population-based surveillance in ten states for laboratory confirmed influenza-related hospitalizations. EIP sites also conduct influenza vaccine effectiveness evaluations among groups for which the Advisory Committee on Immunization Practices (ACIP) recommends annual vaccination.
- **Public Health Emergency Preparedness Cooperative Agreements (PHEP):** Provides some funding for certain pandemic planning and response activities, including partner and clinical laboratory outreach, the purchase of laboratory equipment or support for specimen courier/transport systems.
- **Other special projects:** As resources permit, CDC supports additional studies and special projects such as the Influenza Incidence Surveillance Program (IISP). These programs help increase capacity for participants and provide valuable data for national surveillance.

Additional Federal Resources

In addition to funding, other resources are provided to states by CDC to help minimize the financial and resource burden on each jurisdiction. Listed below are some of the key non-financial resources that help

Sustainable Funding is Critical

At least once a year, the influenza virus changes slightly. It stays ahead of testing and research—and ahead of funding for testing and research. The changing virus is one side of the equation. On the other side is the changing levels of funding. Labs experience “roller-coaster” funding levels—a surge of money in response to a crisis and cuts when a crisis is behind.

“People think of a lab as a building—you build it and you walk away. But you need people who are trained, you need new equipment, you need to stay up-to-date with disease pathogens.”

Excerpted from Lessons from a Virus

CDC and state/local jurisdictions meet the surveillance requirements outlined in this roadmap document.

- **Influenza Reagent Resource (IRR):** Since 2009 CDC's Influenza Reagent Resource provides rRT-PCR reagents to qualified PHLs to help sustain rapid virus detection and subtyping capacity. This is a critical resource to states that significantly reduces the financial burden for state/local jurisdictions and ensures the timely availability of molecular testing reagents intended for virologic surveillance. Financial support for ancillary reagents through the IRR is assessed on an annual basis and is based on the availability of funds. The direct material costs to CDC for each full IVD CDC Human Influenza Virus Real-Time RT-PCR Diagnostic Panel, including Influenza A/B typing and subtyping reagents, superscript, extraction kits and plastics, is approximately \$14,000 (i.e. \$70 for each specimen tested with the 5 target assay by the PHL).
- **CDC-Designated National Surveillance Laboratories:** CDC, in collaboration with APHL, has established advanced capacity in several PHLs to provide antiviral susceptibility testing and influenza virus isolation that serves all PHLs. The virus culture capability supports expanded availability of viruses for antigenic and sequence-based characterization at CDC, providing data and viruses for annual vaccine virus selection.
- **PHLIP and Information Technology Support:** Resources to assist PHL's implement PHLIP have come through CDC Pandemic Influenza funds provided to APHL, although recently additional support has been made available from the CDC Laboratory Response Network (LRN), Laboratory Efficiency Initiative (LEI), and Vaccine Preventable Disease (VPD) initiatives. The broad applicability of PHLIP to other programs and to other efforts to support a "shared services" model among PHLs provides a path to sustainability for PHLIP; however, this will require ongoing focus and effort to ensure continued funding and technical support. Technical assistance teams provide training, on-site assistance, and follow-up consultation.
- **Technical Support and Training:** CDC subject matter experts are readily available to PHL and program staff to address clinical, operational and technical questions. Additionally CDC conducts diagnostic testing of unsubtypeable and other specimens of clinical interest such as suspect antiviral resistant specimens. CDC, in collaboration with APHL, have provided a variety of in-person technical training courses (i.e. PCR, pyrosequencing) for state and local jurisdictions at little to no cost to states to ensure the necessary expertise is readily available at PHLs.

It is critical that CDC be adequately funded to continue supporting state and national activities which ensure an effective national surveillance system. In return, states should ensure that they are meeting ELC, PHEP, and other federal grant benchmarks to be good stewards of these resources.

As funding is always a limiting factor, every state will need to determine how to achieve influenza surveillance goals to meet national and state needs. Federal resources (funding, reagents) distributed to states need to be directed to activities that support overall national priorities. State/local capabilities beyond those recommended as essential to meet national virologic surveillance goals will require financial support from the state.

State Funding Sources

In addition to federal funding sources, many states also receive financial support from their state and/or local jurisdiction. These additional funds can be a critical funding stream for supporting state surveillance activities. The actual mechanisms and level of support varies across states. As data indicated in the 2011 Right Size Influenza Virologic Surveillance Landscape Survey, the amount of state funds expended in support of influenza surveillance varies greatly across states (APHL, 2012). As previously described, it will become increasingly important for state funds to supplement federal funds for testing not deemed essential to meet national virologic surveillance goals. It is important that each state determine which testing services are essential for their state needs.

Implementation Guidance

Influenza virologic surveillance is an essential function of all state health departments and requires partnership between the state public health laboratory representatives, epidemiologists and influenza surveillance coordinators. At a minimum, virologic surveillance includes the ability to:

- Access a representative sample of clinical specimens from ILINet (influenza like illness network) providers or other clinical primary care sources and clinical laboratories.
- Detect type and subtype influenza viruses from clinical specimens in a timely manner using standard laboratory methods.
- Report results to providers, epidemiologists, and CDC using standard electronic data systems.
- Rapidly refer unsubtypeable influenza viruses to CDC to rule out novel viruses.
- Refer a subset of specimens (and viruses) to CDC or CDC-designated laboratory for genetic and antigenic characterization routinely.
- Maintain the expertise, warm base (a minimum level of readiness or capacity), and surge capabilities necessary for pandemic response.

Detailed descriptions of the essential components for virologic surveillance were presented in the [Requirements Section](#). In this section, suggestions and model practices are provided to assist state and local PH directors and laboratorians, epidemiologists and influenza surveillance coordinators operationalize the requirements. The model practices provided in this section are based on experience with the system since its inception in the late 1990's, a series of stakeholder meetings, a table-top exercise conducted in December 2012 testing the utility of the roadmap recommendations, and the data gathered through pilot projects conducted in four states during the 2012-2013 influenza season.

Sampling

A virologic surveillance sampling strategy should be implemented that will ensure year round access to an adequate number of representative clinical specimens to meet key surveillance objectives. Specimens should be obtained from ILINet providers and/or other clinical primary care sources and clinical laboratories. Feasibility and representativeness are the most important factors to consider when choosing specimen providers.

As discussed in the Requirements section, the virologic surveillance landscape can be organized into five tiers based on where sampling and testing is performed. The five tiers of influenza surveillance are described in the Process Model in Appendix A. Since specimens are primarily obtained in the first tier and passed to subsequent tiers for testing, a sampling process takes place at each transfer point. The variation in sampling criteria throughout this sequential process complicates extrapolating the data from one testing tier and applying it to the population of another tier. The variability in sampling can greatly challenge national surveillance efforts where the data is aggregated from multiple states. The Roadmap sampling requirements are intended to apply a more consistent, standardized collection/sampling process to improve overall data confidence and representativeness.

State and Local Implementation Steps

1. Establish a specimen provider network

The primary care tier provides data and specimens for the influenza surveillance system. Selected ILINet providers and/or other primary care sites send respiratory specimens to a public health laboratory according to jurisdictional sampling criteria. ILINet and other clinicians may elect to test specimens using a point of care rapid influenza diagnostic test (RIDT) if one is available in their clinical setting. Hospitalized patients, as well as patients seen by non-ILINet health care providers may have specimens tested by fluorescent antibody, virus culture, or molecular methods in a clinical/commercial laboratory. State and local public health laboratories make up the third tier of influenza surveillance. These laboratories typically perform rRT-PCR testing to type and subtype influenza viruses in clinical specimens.

A. ILINet or other specimen providers (Tier 1)

- Specimen submitters may be ILINet sites or other primary care health care providers. The selected specimen submitters should be committed to collecting high quality specimens, and submitting the required number of samples in a timely manner and in accordance with jurisdictional criteria throughout the entire year. A number of states report that ILINet sites are generally not a good source of specimens, so alternate primary care sites have been recruited as specimen submitters.
- Specimen provider recruitment and submission criteria should be established so that specimens submitted for virologic surveillance are representative of the diversity of the population as a whole or of specific targeted population as needed. The collective group of selected specimen providers should cover all age groups.

- Unscreened specimens are preferred for routine seasonal surveillance. If primary care submitters are using RIDT's for diagnostic purposes, a random mix of specimens, irrespective of the test result, should be sent to the PHL for surveillance purposes. This provides a better assessment of true positivity in the community, and reduces potential bias introduced by pre-screening with tests that have variable sensitivity, and may not detect novel or drifted viruses (i.e., give false negative result)⁷. Outside of influenza season, all RIDT positives should be submitted to the PHL. In a novel event investigation, oversampling pre-screened positives may be appropriate if the tests used are high performing with demonstrated reliability for detection of the virus of interest.
- Provider compliance with specimen submission criteria may be enhanced by providing:
 - clear instructions and submission forms customized for their site,
 - cost-free specimen collection kits and shipping,
 - feedback and data to submitters, including influenza test results and/or aggregate results of testing for other respiratory pathogens if performed,
 - training.
 - Some PHLs and/or surveillance coordinators have also elected to provide RIDT kits to incentivize specimen submission.

RIDT data and specimens contribute to Influenza Surveillance

The Iowa statewide influenza surveillance program collects data on the number of RIDTs performed and the percentage positive each week using a survey monkey tool. Additionally, during times of low prevalence, laboratories submit rapid test positive specimens to the State Hygienic Laboratory (SHL) for confirmatory testing using the CDC's real-time RT-PCR test panel. The RIDT survey data and the results of confirmatory testing are incorporated into the weekly influenza surveillance report compiled by the Iowa Department of Public Health. This report is widely distributed to public health officials, infection control practitioners, health care providers and others to improve awareness about seasonal influenza activity and reliability of RIDT results.

B. Clinical laboratory providers (Tier 2)

In addition to the ILI/primary care provider network, virologic surveillance should include specimens from hospital/clinical laboratories to ensure that a subset of specimens represent more severe illness (inpatients, mortality, unusual cases) and outbreak sources. Many clinical labs also serve as reference labs for outpatient satellite clinics, and therefore may be a good source of ILI specimens for routine surveillance. Clinical laboratories will also be essential partners when responding to large scale outbreaks or a pandemic. The influenza surveillance

⁷ Balish A, Garten R, Klimov A, Villanueva J. Analytical detection of influenza A(H3N2)v and other A variant viruses from the USA by rapid influenza diagnostic tests. *Influenza Other Respi Viruses* 2012;September 18 [Epub ahead of print].

coordinator, in collaboration with the PHL, should develop and disseminate policies and establish mechanisms to ensure submission of a subset of positive specimens and all unsubtypable influenza positives (if subtyping assays are used) from hospital/clinical laboratories performing influenza testing. If clinical laboratories are the primary resource for surveillance specimens, the specimens sent to the PHL may be overly representative of hospitalized patients (i.e., more severe cases). Specimens from clinical laboratories should include both influenza positive and negative samples. PHL testing of negative specimens will be useful to monitor the performance of test methods used in clinical laboratories and enhance likelihood of identifying novel viruses that may not be detected by commercial influenza assays.

C. Public Health Laboratories (Tier 3):

Every public health laboratory participating in virologic surveillance is responsible for testing clinical specimens submitted for surveillance purposes or epidemiologic investigations, and reporting data to CDC in a timely manner. PHLs are also required to submit representative clinical specimens and/or virus isolates to CDC (Tier 5) or CDC-designated laboratory (Tier 4) for national surveillance purposes, including annual vaccine virus selection. Viruses from these specimens may also be used to manufacture seasonal influenza vaccines. The 2013-2014 vaccine will contain A/California/7/2009pdm09-like(2009 H1N1) virus, A/Texas/50/2012(H3N2) and B/Massachusetts/2/2012-like (B/Yamagata lineage) virus.

Laboratories should submit specimens/viruses based on annual CDC criteria and guidance which is sent to state PHL Directors and disseminated by APHL. In collaboration with APHL and CSTE, CDC also convenes teleconferences pre-season and throughout the season as needed to update surveillance guidance. Participation in these teleconferences is strongly encouraged.

PHLs performing virus culture should send both the original clinical material and the virus isolate to CDC (or CDC-designated laboratory). Providing the virus isolate along with the original clinical material allows for more rapid antigenic characterization at CDC. Maintaining virus culture requires specialized cell lines and well trained staff with sufficient expertise to perform high quality testing. PHLs opting not to perform virus culture should submit original clinical specimens with a minimum volume of xx mls.

To enhance CDC's vaccine virus selection efforts, it is important to send recently collected specimens. Specimens submitted to CDC should be representative of the circulating influenza types/subtypes, geography, disease severity and age. Oversampling of low prevalence subtypes may be necessary to ensure that all circulating subtypes are represented in the samples sent to CDC. When available, viruses from particularly severe or unusual cases, and a sample of viruses isolated from outbreak investigations should also be represented in submissions to CDC. The two examples below demonstrate the criteria that should be considered by the PHL when selecting the specimens that will be sent to CDC for routine national surveillance purposes.

- In a two week period, the PHL testing yields 50 A/H3, 1 A/2009H1, and 5 Influenza B positive specimens. Send to CDC or the CDC-designated laboratory: the A/H1 specimen, one Flu B, and 3 A/H3 viruses that are representative of state geography and patient ages.

- In a two week period, the PHL testing yields 20 A/H3, one of which is from a patient who died, 12 A/2009H1 viruses, and 5 Influenza B viruses. Send to CDC or the CDC-designated laboratory: the A/H3 specimen from the patient who died, 2 other representative A/H3 specimens, 2 A/2009 H 1 and one Flu B specimen.

CDC may request additional viruses/specimens depending on circulating virus trends, vaccine virus selection and vaccine candidate development needs. CDC strongly recommends that PHLs subtype all, and at least 90% of Influenza A positives. Unsubtypable⁸ viruses that may represent a novel subtype should be submitted to CDC within 24 hours of detection.

2. Determine appropriate sample sizes for each surveillance objective

The need to characterize and improve the precision of the data that is provided through virologic surveillance was one of the principal drivers of the Right Size Virologic Surveillance project. Implementing a statistical, systematic approach to determine the appropriate number of specimens to be collected and tested can be achieved by using sample size calculators. The calculators developed as right-size virologic surveillance tools provide a statistical basis to estimate the number of specimens to be tested in order to provide a specific level of data confidence for situational awareness, novel event detection and novel event investigation. In addition, these calculators can also be used to determine the confidence level of data derived from an existing sample of ILI patient specimens.

Some state and local public health laboratories may need to test more or fewer specimens to achieve the same level of data confidence as another state or local jurisdiction with a larger or smaller population. Alternately, influenza surveillance coordinators may need to accept a lower confidence level or higher margin of error if the system does not have the capacity to collect or test the number of samples estimated by the calculators.

Efficiencies in the total number of specimens that need to be tested can be achieved by using a single sampling approach to address as many surveillance objectives as possible. For example, situational awareness and novel event detection rely on samples collected and tested for routine surveillance. At the national level, routine antiviral resistance surveillance testing currently uses the same samples as vaccine strain selection testing. Where differences are important, they should be addressed by the sampling strategy.

The sample size calculations are based on population size, desired level of confidence, margin of error, and estimated or known prevalence or threshold for detection. More details on thresholds are provided in the Objectives: Thresholds and Representativeness section. State and local public health laboratories are encouraged to use sample size calculators or pre-calculated sample size tables to achieve a more scientific, statistically based sample size that supports surveillance objectives. Specimen sampling approaches should be established to prioritize collecting an

⁸ Any influenza positive specimen that cannot be definitively typed and subtyped as a circulating seasonal influenza virus. Influenza positive specimens producing non-standard or inconclusive results as defined in the CDC Human Influenza Virus Real-Time RT-PCR Diagnostic Panel Instructions for Use package insert.

adequate number of specimens for detection of novel events based on national thresholds, while at the same time providing sufficient number of representative for overall situational awareness at the state level. For many states, the number of samples to be tested for each of these objectives is very similar. However, for smaller states, the number of samples necessary to achieve high confidence in situational awareness state level data will be much higher than the number of samples needed to contribute to national novel event detection thresholds. Outside of influenza season, achieving statistical confidence will not be possible in most states; the focus of surveillance should shift to obtaining all specimens from clinical sites that have tested positive for influenza, or from patients with unusual respiratory illness, travel history or risk of exposure to animal-origin viruses.

Targeted surveillance may be useful to answer specific questions, especially when conducting an investigation if a novel event or new virus is detected. Therefore the surveillance program should have the capability to establish targeted surveillance of specific populations if needed. CDC will provide guidance to state epidemiologists and PHLs on the specific risk factors and need for enhanced surveillance (e.g., HPAI H5N1 risk factors, swine exposure). However, the current version of the novel event investigation calculator may not be useful in these situations, future editions of the Roadmap are expected to provide more options for targeted surveillance, addressing intentional and unintentional bias.

a. Calculator Inputs and Outputs:

The key variables in calculating sample size are described in the table below. Understanding how these variables affect sample size and data confidence levels is important for generating valuable surveillance data.

	Relationship to Sample Size
Confidence Level	The confidence level is the amount of uncertainty that the true prevalence is equivalent to the estimated prevalence. As this value increases the sample size also increases.
Margin of Error	This is the amount of error that can be tolerated. A 2% error would mean that the calculated prevalence may be plus or minus 2% from the true answer. As this value decreases the sample size also increases.
Population	This is the population under surveillance. For routine influenza surveillance, this is the number of people in your state with ILI. As the population size increases the sample size increases.
Medically attended ILI (MA-ILI)	The population of individuals with influenza-like illness who seek medical care. This is the subset of the population available for surveillance testing. This number is determined based on estimates that each person in the US visits an emergency room or generalist physician 2 times per year, and that the percentage of outpatient visits that are for influenza like illness is 2.2% at CDC ILINet Seasonal Baseline).

Expected Prevalence	In the calculators, this is the prevalence that you expect to calculate or the level of detection you wish to achieve. For the purposes of calculating sample size, the expected prevalence refers to the prevalence of influenza positive specimens among the number of ILI specimens tested. This is NOT the prevalence of disease in the community. Note that as the expected prevalence decreases, the sample size becomes smaller when the margin of error is held constant. This seems counter intuitive, but when you scale your margin of error to align with your expected prevalence, the sample size should increase. For instance, a 5% margin of error is more appropriate for a predicted prevalence of 50% than a predicted prevalence of 1%. A more appropriate margin of error for a predicted prevalence of 1% may be 0.5%. Thus, it is important to scale your margin of error to your predicted prevalence.
----------------------------	---

b. Choosing an appropriate confidence level and error rate:

The number of samples to be tested will vary depending on the Confidence Level and Margin of Error used in the calculators. Selecting a lower confidence level increases the amount of uncertainty in the calculated prevalence. A higher margin of error means that more error can be tolerated. Stakeholders and exercise participants identified optimal, mid-range and minimal confidence levels and error rates or thresholds for the two objectives that comprise routine surveillance.

	Situational Awareness		Novel event Detection	
	Confidence Level	Margin of Error	Confidence Level	Threshold
Optimal	95	5	95	1/200
Mid-range	90	5	95	1/165
Minimum	85	5	90	1/165

Pre-calculated sample sizes for each of the objectives using these confidence levels are provided in Appendix B. Depending on the surveillance priorities and capacity of the system overall, jurisdictions may choose to use the on-line calculators to vary the inputs to see how sample size is impacted.

- c. **Assumptions:** As is typical with most models, the sample size calculators rely on certain assumptions regarding the population, or the expected prevalence (positivity rate). The assumptions used in these calculators are based on existing and/or historical ambulatory care and seasonal influenza data.
- d. **Using the Sample Size Calculators:** The virologic surveillance sample size calculators are available as a web-based tool at www.to.be.determined.org. Basic information on the intent of

the calculator for each of the key surveillance objectives, along with inputs and output examples is provided here. A more detailed “user guide” with instructions for using the web-based tools is provided in Appendix B. The appendix also includes tables with pre-calculated sample sizes for situational awareness and novel event detection covering a range of population sizes. States may opt to use these tables for quick reference or as an alternative to the on-line tool.

i. **Situational Awareness- Flu⁺/medically attended ILI (MA-ILI)**

- **Surveillance Objective:** determine the beginning and end of the influenza season and monitor the prevalence and spread of influenza viruses throughout the year.
- **Threshold:** The beginning of the influenza season is defined as the time when the prevalence of Flu⁺ among MA-ILI is at or above 10% over two consecutive weeks. This value roughly corresponds to the CDC ILINet Seasonal Baseline where the percentage of outpatient visits for influenza like illness reaches 2.2%. Jurisdictions may choose to alter the percent positive used in the sample size calculator to more accurately determine the amount of testing needed throughout the season or assess the confidence level of the data provided.
- **Surveillance Question:** How many specimens from MA-ILI patients does the laboratory need to test (in a given period, usually one week) to determine that the prevalence of Flu⁺ specimens among MA-ILI persons tested is x% at a specified confidence level and error rate?
- **Assumptions:** Each person in the US visits an emergency room or generalist physician 2 times per year. 2.2% of medical visits are for ILI outside of influenza season ILINet baseline) (ref). The providers are randomly selecting patients with ILI for surveillance testing. The specimens tested were either not pre-screened, or submitted randomly irrespective of test result.
- **User Inputs:**
 - Population-: The input is the total population under surveillance (e.g., state population). The calculator uses this number and the assumptions above to estimate the weekly number of MA-ILI cases.
 - Estimated prevalence of MA-ILI: Input based on ILINet data during the season. The default is 2.2%, which is the estimated ILINet seasonal baseline for the percentage of outpatient visits that are for influenza like illness.
 - Expected prevalence of Flu⁺/MA-ILI: Input your surveillance target. The default value is 10% for the beginning and end of the influenza season. Other percent positive values may be used based on jurisdictional preferences or seasonal variability in the prevalence of ILI.
 - Specified confidence level: The optimal level of confidence for situational awareness is 95%, the minimum should be no less than 85%.
 - Acceptable margin of error: An acceptable margin of error should be no greater than 5%.
- **Output example:** A sample size of 132 unscreened MA-ILI specimens is needed in order to be 95% confident that the true prevalence of Flu⁺/MA-ILI is 10% +/- 5%.

- **Alternate calculation (sample power):** Determine the level of confidence and margin of error associated with the measured prevalence of flu positives, given the sample size tested, i.e., what are the confidence and error rates associated with current sample size?
- **Alternate Output example:** If 100 MA-ILI specimens were tested and the estimated prevalence is 10%, you can be 70 % +/- 3% that the true prevalence is 10%.

ii. **Detecting a Rare/Novel Event**

- **Surveillance Objective:** Detect a rare event/novel influenza virus among influenza positive surveillance specimens tested in all states at a low enough threshold for effective intervention and control measures.
- **Threshold:** State surveillance systems should collectively test a sufficient number of specimens to detect a novel influenza virus at a prevalence threshold of 0.5% (1/200) Novel Flu+/Total Flu+ specimens at the aggregated national level. At a minimum, a threshold of 1/165 may be used for determining the sample size in states with limited resources.
- **Surveillance Question:** How many specimens does the PHL need to test to allow the national surveillance system to detect a rare/novel virus at 0.5% prevalence with 95% confidence (aggregating testing data from all states)?
- **Assumptions used in the calculator:**
 - Specimens are collected randomly.
 - There is no correction for finite population size – this is a conservative assumption to prevent undersampling. Correcting for finite sample size requires accurately characterizing the surveillance population. In the case of a novel event, the size of the relevant population may be largely unknown. The sample size determined without correcting for a finite population size is always correct. If a sample size correction factor is improperly applied, the target population will be under-sampled, resulting in an overestimate of the confidence level and underestimate of the error.
- **Options:** The novel event detection sample size calculation can be made based on a) the number of positives already identified as Flu+ by a clinical laboratory or the PHL, b) the number of MA-ILI specimens, or c) a combination of both. Although testing prescreened Flu+ specimens decreases the total number of specimens needed to meet the recommended threshold and confidence level, using only specimens that are Flu+ received from clinical laboratories may reduce the sensitivity of the system to detect novel events because of the unknown sensitivity of commercial systems to detect novel or drifted viruses. Using a combination of Flu+ and MA-ILI specimens will moderate the potential loss in sensitivity, and allow PHLs with large populations to achieve statistical confidence with reasonable specimen numbers.
- **User Inputs:**
 - **Population-:** The input is the total population under surveillance (e.g., state population).

- Surveillance scale: The default is national, meaning that all states are contributing to a national surveillance effort proportional to their population size. The number of samples that a state needs to tests is apportioned based on population size. The calculator also provides the option for states to calculate the number of specimens to test for detection of a novel event at a specific threshold *within their state, however, the sample size for an individual state at the same 1/200 threshold will be significantly larger than that needed for the national threshold.*
- Specified confidence level: The optimal level of confidence for novel event detection is 95% at 1/200, the minimum should be no less than 90% at 1/165.
- Expected Flu+/MA-ILI: This is an input when calculating the number of MA-ILI specimens needed, or the number of combined MA-ILI and Flu+ specimens. Use 10% positivity for seasonal baseline, other percent positivity as needed throughout the year.
- **Output examples:**
 - Number of Flu+ specimens: To be 95% confident of detecting 1 or more rare events at a prevalence of 1/200 at a national level, the PHL must test 11 Flu+ specimens per week.
 - Number of MA-ILI specimens: To be 95% confident of detecting 1 or more rare events at a prevalence of 1/200 at a national level, the PHL must test 108 MA-ILI specimens per week.
 - Combined number of Flu+ and MA-ILI specimens: To be 95% confident of detecting 1 or more rare events at a prevalence of 1/200 at a national level, the PHL must test 58 MA-ILI and 5 Flu+ specimens per week.
 - Combined number of Flu+ and MA-ILI specimens (state level): To be 95% confident of detecting 1 or more rare events at a prevalence of 1/200 at a state level, the PHL must test 257 MA-ILI and 411 Flu+ specimens per week.
- **Alternate calculation (sample power):** Determine the level of confidence that a novel event can be detected at a given threshold, given the sample size tested.
 - **User Input:** the number of Flu+ specimens tested and the number of MA-ILI specimens tested.
- **Alternate output:** If the laboratory tested x Flu+ and y MA-ILI specimens, you can be 70% confident that the novel virus would be detected at a prevalence of 1/200.

iii. **Rare/Novel Event Investigation**

- **Surveillance Objective:** Determine the prevalence of the novel virus (Novel Flu+/Total Flu+) within a state following detection of a novel influenza virus; *confirm that the prevalence of a rare event does not* exceed a specific percent positivity. Investigation of a novel event is typically performed using enhanced, targeted surveillance.
- **Threshold:** There are no defined thresholds for novel event investigation, as specific situations and jurisdictional considerations may warrant different thresholds. In

general, if the novel event was detected at 1/200, the investigation threshold should be between 1-5%.

- **Surveillance Question:** Once a novel virus is detected, how many ILI specimens does the PHL need to test to determine that the true prevalence does not exceed a specified percent of Flu+?
- **Assumptions:**
 - Specimens are collected randomly.
 - There is no correction for finite population size – this is a conservative assumption to prevent under sampling. Correcting for finite sample size requires accurately characterizing the surveillance population. In the case of a novel event investigation, the size of the relevant population may be largely unknown. The sample size determined without correcting for a finite population size is always correct. If a sample size correction factor is improperly applied, the target population will be under-sampled, resulting in an overestimate of the confidence level and underestimate of the error.
 - This calculator would be most relevant in an H1N1 type event, where the at-risk population group is unknown. This assumption, however, results in very high sample sizes. This calculator may not be appropriate when targeted surveillance is a more efficient initial approach, such as the 2012 H3N2v summer surveillance scenario targeting visitors to state/county fairs.
 - Asymmetrical distribution
- **User Inputs:**
 - Population: The input is the total population under surveillance (e.g., state population).
 - Surveillance scale: STATE vs national. The default is state because the investigation of the novel event would occur locally.
 - Expected prevalence of Rare+/Flu+: This is the percent positivity of the rare event that you want to confirm has not been exceeded.
 - Specified confidence level: The optimal level of confidence is 95%, the minimum should be no less than 85%.
 - Expected Flu+/MA-ILI: This is an input only for when calculating the number of MA-ILI needed, or the number of combined MA-ILI and Flu+ specimens. Use 10% positivity for seasonal baseline, other percent positivity as needed throughout the year.
- **Output examples:**
 - Number of Flu+ specimens: To be 95% confident that the true prevalence of the novel virus does not exceed 1% +/- 2% of Flu+ specimens, the PHL must test 478 Flu+ specimens per week.
 - Number of MA-ILI specimens: To be 95% confident that the true prevalence of the novel virus does not exceed 1% +/- 2% of Flu+ specimens, the PHL must test 4778 MA-ILI specimens per week.

- Combined number of Flu+ and MA-ILI specimens: To be 95% confident that the true prevalence of the novel virus does not exceed 1% +/- 2% of Flu+ specimens, the PHL must test 533 MA-ILI and 359 Flu+ specimens per week.
- **Alternate calculation:** Determine the level of confidence given the sample size tested.
- **Alternate output:** If a combination of **75** Flu+ specimens and **300** unscreened MA-ILI specimens were tested, and the estimated prevalence of the novel virus among all flu positive specimens (Rare+/Flu+) is **1%**, you can be 75% confidence that the true prevalence does not exceed 1% +/-1%. (This assumes that **10%** of MA-ILI+ patients are Flu+).

iv. **Detecting/Monitoring Antiviral resistance**

- **Surveillance Objective:** Detect antiviral resistance virus among influenza positive surveillance specimens tested in all states at a low enough threshold for effective intervention and control measures.
- **Threshold:** 5% prevalence of oseltamivir resistant viruses among positive specimens for each influenza A subtype or influenza B at the national level.
- **Surveillance Question:** How many of each influenza A subtype or influenza B Flu+ specimens does the laboratory need to test to allow the national surveillance system to detect antiviral resistant viruses at or below a 5% prevalence with 95% confidence (aggregating testing data from all states)?
- **User Inputs:** The Flu+ tab of the novel event detection calculator can be used to determine sample size for this objective.
 - **Population-:** The input is the total population under surveillance (e.g., state population).
 - **Surveillance scale:** The default is national, representing the number of specimens that need to be tested by the state to detect antiviral resistance at a national aggregated threshold. The number of samples the state needs to test is apportioned based on population size. States wishing to calculate the number of specimens to test for detection of antiviral resistance at a specific threshold within their state can select their state, note that the sample size of an individual state will be significantly larger than that needed for the national threshold.
 - **Specified confidence level:** The optimal level of confidence for antiviral resistance is 95%, the minimum should be no less than 85%.
 - **Expected Flu+/MA-ILI:** This is an input only for when calculating the number of MA-ILI needed, or the number of combined MA-ILI and Flu+ specimens. Use 10% positivity for seasonal baseline, other percent positivity as needed throughout the year.
- **Output example:**
 - Number of Flu+ specimens: To be 95% confident of detecting antiviral resistant Influenza A 2009H1N1 viruses at a prevalence of 5% among influenza A 2009H1N1

positive specimens tested at the national level, the PHL must test x Influenza A H1N1 Flu+ specimens per week.

- **Alternate calculation (sample power):** Determine the level of confidence that antiviral resistant viruses can be detected at a given threshold, given the sample size tested.
 - o **User Input:** the number of Flu+ specimens tested and the number of MA-ILI specimens tested.
- **Alternate output:** If the laboratory tested x Flu+ and y MA-ILI specimens, you can be 70% confident that the antiviral resistant viruses would be detected at a prevalence of 5%.

3. Establish policy for frequency of submissions

a. Primary Care and clinical laboratory specimen submissions to PHL

The frequency of specimen submission for routine surveillance will vary depending on jurisdictional needs, and PHL capacity for specimen intake and processing. During influenza season it may be most convenient to ask providers to send specimens from the first few ILI patients they see each week. If the PHL prefers to receive specimens throughout the week, each provider may be asked to collect and send specimens on a different day. Specimens/viruses need to be submitted and tested in real time, not batched, in order to inform timely clinical management guidelines. If specimens are being sent to the PHL for diagnostic testing (patient with high risk travel history, or unusual case presentation), these specimens should be transported promptly and not batched with surveillance specimens. Clinical laboratories that perform PCR testing with subtyping should immediately submit any specimens that produce unsubtypeable test results.

b. PH Labs submission to CDC or CDC-designated laboratory

Laboratories should submit specimens/viruses for routine surveillance year round based on annual CDC criteria and guidance provided to state PHL Directors and disseminated by APHL. Routine surveillance specimens should be forwarded to CDC or a CDC-designated laboratory in a timely manner to provide real-time surveillance information. Ship routine surveillance specimens at least once every two weeks, this ensures that CDC can perform further characterization in time to guide international and domestic annual vaccine virus selection. Unsubtypeable specimens, as defined in the RT-PCR package insert, require immediate action as they may reflect a novel virus with pandemic potential. These specimens are to be sent immediately to CDC for more comprehensive testing to ensure that appropriate interventions can be implemented if needed, and that CDC meets WHO international health regulations⁹ for novel virus reporting.

⁹ IHR Regulations: <http://www.who.int/ihr/en/>. State Parties to the IHR (2005) are required to immediately notify WHO of any laboratory confirmed case of a recent human infection caused by an influenza A virus with the potential to cause a pandemic.

4. Ensure samples are of acceptable quality

Influenza surveillance coordinators and PHLs should provide instructions and training to specimen submitters to ensure that respiratory specimens are of high quality, properly collected, stored and transported.

a. Specimen collection

Respiratory tract specimens required for influenza diagnosis and identification are well-defined and include nasopharyngeal swabs and throat swabs, submitted separately or combined, nasopharyngeal aspirates, nasal washes, bronchoalveolar lavages, tracheal aspirates, bronchial washes and, following autopsy, respiratory tract tissues. The most appropriate specimen to collect depends upon the diagnostic test employed. This information will be provided by the test/reagent manufacturer and the laboratory performing the test. Additional resources can be found in clinical microbiology textbooks, and at the CDC website www.cdc.gov/flu/professionals/diagnosis/index.htm.

Diagnostic test results are only as good as the quality of the specimen. Specimen quality depends on proper collection technique and the amount of virus present at the source. The amount of virus shed in the upper respiratory tract declines over the course of the illness, therefore collecting specimens as close to symptom onset as possible is recommended. Optimally, specimens for virologic surveillance should be collected within 24-72 hours of symptom onset, and no later than 5 days post onset of symptoms.

Specimen providers need to be trained in proper collection technique. It is ultimately the responsibility of the laboratory to ensure that specimens are properly collected. Descriptions of proper methods for specimen collection can be found in clinical textbooks, in product inserts, and online. The most effective method, however, is demonstration by someone skilled in the collection technique, followed by practice under observation. The Joint Commission Strategies for Improving Rapid Influenza Testing in Ambulatory Settings (SIRAS) website www.jointcommission.org/siras.aspx offers two free on-line courses one for health care providers in ambulatory settings and one for specimen collectors.

b. Specimen Handling

Specimen quality also depends on proper handling of the specimen after collection. The laboratory, in coordination with the influenza surveillance coordinator, is responsible for providing information on proper specimen handling to specimen providers.

Specimens should be placed immediately into an acceptable viral transport medium in accordance with standard testing protocols or kit manufacturer recommendations and held at 4-8°C until testing is performed. Testing ideally should be performed as soon as possible. If a delay of more than 2-3 days until specimens are tested is anticipated, specimens can be frozen at -

An influenza A virus is considered to have the potential to cause a pandemic if the virus has demonstrated the capacity to infect a human and if the hemagglutinin gene (or protein) is not a variant or mutated form of those, i.e. A/H1 or A/H3, circulating widely in the human population.

70C. However multiple freezing and thawing of specimens can adversely affect the test result and should be avoided whenever possible. Virus isolates and nucleic acid extracts also require special handling.

5. Establish and support specimen transport systems

Specimen transport is another critical component of influenza virologic surveillance. Timely and efficient transport of specimens is often quite costly, and must be adequately funded by the public health system for effective surveillance. Specimen collection and regulation compliant transport supplies, as well as courier/carrier costs, need to be covered. It is not reasonable to expect the providers and clinical labs to assume these costs for surveillance testing.

Specimen integrity must be maintained during transit. An effective and efficient process for specimen submission must account for the reliable and timely transport of specimens from clinical sites (providers) and clinical labs to the PHL and from the PHL to CDC or CDC-designated laboratories. Specimen transport must comply with a variety of federal (DOT, USPS) and international (IATA) regulations that ensure that specimens and infectious materials are properly packaged and safely shipped (ref). In-state commercial couriers, healthcare system couriers, PHL-provided couriers or national carriers can be employed to transport specimens to the PHL. Redundancy in transport options is important to cover disruption of any particular method of transport and to provide maximum daily service. An interstate carrier is most often used for transport to CDC or the CDC-designated laboratories.

In special circumstances, direct shipment from the health care provider or clinical laboratory to the CDC may be warranted; however, this should be facilitated whenever possible by the PHL to ensure proper handling and state epidemiologist engagement if case investigation is needed.

6. Recognize and Address Sampling Biases

The influenza virologic surveillance system contains inherent biases due to the complexity of the sampling system and the use of different test methods in the different tiers. Sources of bias should be considered and addressed if possible when selecting specimen providers, selecting test methods, analyzing data and interpreting results.

- a. Specimen providers: should represent the entire population under surveillance. Choose a mix of primary care health care providers representing all age groups (pediatrics, family practice, internal medicine and geriatrics). Specimen providers should be selected representing areas of diverse population density (urban, suburban, and rural).
- b. Unscreened vs. prescreened specimens: efforts should be made to limit sampling of prescreened (influenza positive) specimens. As previously discussed, unscreened specimens are preferred. If submitters are using RIDT's for diagnostic purposes, a random mix of positive and negative specimens, irrespective of RIDT results, should be submitted to the PHL for surveillance purposes. At a minimum, data should differentiate pre-screened from unscreened specimens. If prescreened specimens from clinical laboratories are the primary source of surveillance specimens, these may be overly representative of hospitalized patients, i.e., bias to severe cases). Data may not be representative of true prevalence of virus subtypes in the community.

Laboratory Testing

Laboratory testing requires staff with expertise in the methods performed, adequate space with appropriate biosafety controls, well-maintained equipment, appropriate testing reagents and supplies, and reliable results reporting mechanisms. Essential equipment, materials, and reagents necessary to perform influenza virologic surveillance are listed in Appendix C: Laboratory Methods.

Influenza virologic testing methods are classified in the roadmap based on a stakeholder assessment of requirements that should be maintained and available at all PHLs involved in influenza surveillance (primary testing); or as additional influenza surveillance laboratory capabilities that may be maintained based on state and/or jurisdictional needs or provided through a shared services model. The laboratory must ensure quality for all influenza testing methodologies performed for influenza surveillance.

Primary Testing Method:

PHLs performing virologic surveillance are expected to utilize molecular methods, such as reverse transcriptase real-time PCR (rRT-PCR), as the primary testing method for influenza detection and subtyping. This is an ELC benchmark. During rRT-PCR test processes, viral RNA is extracted from patient specimens, transcribed into DNA and then amplified. The product of the test reaction is detected in “real time” using labeled probes. Reverse transcriptase real-time PCR can rapidly identify influenza A and B, distinguish influenza A subtypes, and offers the best performance characteristics (i.e. sensitivity, specificity) of all currently available testing methods. CDC, through the Influenza Reagent Resource (IRR), manufactures and distributes the CDC Human Influenza Virus Real-Time RT-PCR Diagnostic Panel (CDC Flu rRT-PCR Dx Panel) to all state and qualified local PHLs engaged in influenza surveillance testing to ensure widespread use of a nationally standardized protocol. The CDC also has the capability to rapidly adapt the assay to detect newly emerging viruses. CDC Influenza diagnostic panel is distributed under FDA special controls. The special controls require that all users of the diagnostic device be trained to perform and interpret the results by a competent instructor prior to use and that CDC limit distribution to only those users who have successfully completed a training course provided or approved by CDC.

If the PHL decides not to use the CDC Flu rRT-PCR Dx Panel for influenza surveillance testing but instead chooses an alternative molecular influenza testing methodology or platform, the PHL must ensure that the manufacturer of the assay monitors and updates assay performance as circulating viruses change over time and that the assay being utilized within the laboratory has been optimized to identify all currently circulating influenza viruses.

Influenza testing algorithms should be adopted in order to optimize testing efficiency and throughput. Laboratories should choose an algorithm that best fits their laboratory test flow, surveillance needs as the influenza prevalence changes throughout the year, and ensures judicious use of CDC provided reagents. These should be continually reviewed by PHLs throughout the influenza season to ensure that the most efficient algorithm is being utilized. The direct material costs to CDC for each full IVD CDC Human Influenza Virus Real-Time RT-PCR Diagnostic Panel, including Influenza A/B typing and subtyping reagents, superscript, extraction kits and plastics) is approximately \$14,000 (equals \$70 for each specimen tested with the 5 target assay by the PHL).

Below are three common testing algorithms to consider.

- Using the 7-target algorithm, influenza typing and subtyping are performed on a single plate at the same time. This algorithm tests simultaneously for influenza A, B, A/H1, A/H3, A/pdmH12009, and RNaseP (internal control).
- The Influenza A/B Reflex algorithm is a two-step process in which Influenza A/B typing (A, B, RNaseP) is performed first and the Influenza A subtyping (H1, H3, H1pdm2009) is performed only on Influenza A positive specimens.
- Surge algorithms can also be implemented if needed to accommodate high volume testing demand in an outbreak investigation or pandemic. This may include first screening all specimens associated with outbreak for the outbreak strain (e.g., A, ApdmH1), followed by reflex testing of negative samples to detect other circulating influenza viruses.

The public health laboratory is responsible for the timely referral of representative specimens (and viruses, if culture is performed) to CDC or a CDC-designated PHL for genetic and antigenic characterization throughout the year. To enhance CDC's vaccine virus selection efforts, it is important to routinely send recently collected specimens. Specimens submitted to CDC should be representative of the circulating influenza types/subtypes, geography, disease severity and age. Oversampling of low prevalence subtypes may be necessary to ensure that all circulating subtypes are represented in the samples sent to CDC. When available, viruses from particularly severe or unusual cases, and a sample of viruses isolated from outbreak investigations should also be represented in submissions to CDC. The two examples below illustrate the criteria that should be considered by the PHL when selecting the specimens that will be sent to CDC for routine national surveillance purposes.

- In a two week period, the PHL testing yields 50 A/H3, 1 A/2009H1, and 5 Influenza B positive specimens. Send to CDC or the CDC-designated laboratory: the A/H1 specimen, one Flu B, and 3 A/H3 viruses that are representative of state geography and patient ages.
- In a two week period, the PHL testing yields 20 A/H3, one of which is from a patient who died, 12 A/2009H1 viruses, and 5 Influenza B viruses. Send to CDC or the CDC-designated laboratory: the A/H3 specimen from the patient who died, 2 other representative A/H3 specimens, 2 A/2009 H 1 and one Flu B specimen.

Outside of influenza season, PHLs should follow CDC summer submission guidance and send all specimens that test positive by PCR to CDC or CDC-designated laboratory for further characterization.

IMPORTANT REPORTING AND REFERRAL REQUIREMENT: At any time through the year, if the PHL identifies a specimen as unsubtypable following testing using all available targets (influenza A, B, A/H1, A/H3, RNP, etc), the laboratory must notify CDC immediately and refer the specimen to CDC within 24 hours of detection. Specific information about interpretation of inconclusive test results and referral requirements can be found in the CDC Human Influenza Virus Real-Time RT-PCR Diagnostic Panel package insert.

To ensure national and state coordination on virologic surveillance priorities, "Influenza Seasonal Kick off Teleconferences" are convened for PHLs, influenza coordinators, and state epidemiologists in the

early fall each year. The main purpose of the teleconferences is to provide a situational update and surveillance guidance for the upcoming season, including expectations for submission of specimens to CDC. The guidance will identify how many specimens to submit throughout the season, where to submit the specimens, and the specimen submission form. Announcements with teleconference information along with annual electronic forms are sent to all PHL directors via email at the beginning of each season.

Additional Testing Methods:

Additional testing methods include influenza virus culture, antiviral resistance testing, influenza hemagglutination inhibition (HAI), immunofluorescence testing, and serology. These methodology recommendations focus specifically on advantages and disadvantages for influenza and may not apply to other virus assays. Each testing methodology listed has many advantages and disadvantages that will factor into jurisdictional test methodology decisions. Some of the advantages or disadvantages listed here are more applicable to national surveillance needs, but are included to help inform jurisdictional decisions.

Influenza Virus Culture

PHLs are encouraged to perform virus culture if they can sustain the appropriate level of staff expertise and the necessary resources. When PHLs submit virus isolates along with clinical material to CDC or the CDC-designated public health laboratory, this improves the efficiency of the antigenic characterization process at CDC, so that information regarding influenza vaccine match or virus drift can be disseminated back to public health laboratories more quickly.

At minimum this capability must be maintained at CDC, and at a subset of state PHLs. PHLs that do not have the capability or resources to perform high quality virus culture using consensus protocols may opt to utilize a shared services model instead, relying on the CDC or state PHLs that have been designated by CDC as national surveillance laboratories to provide virus culture.

Advantages of Maintaining or Implementing Influenza Virus Culture:**PHL:**

- Provides isolates for validation and verification of new or modified assays, and troubleshooting investigations.
- Provides a back-up method to PCR.
- Detects other respiratory viruses.
- Provides viruses required for phenotypic antiviral resistance testing.

CDC:

- Provides viruses required for phenotypic and antigenic characterization. These are critical components of surveillance for vaccine virus strain selection, and development of the annual WHO kit distributed to domestic and international laboratories.
- Provides viruses that can be used to develop vaccine candidates.

Disadvantages of Influenza Virus Culture:

- Less rapid and sensitive than rRT-PCR and is not efficient in times of surge.

- Requires specialized expertise and capability to maintain high quality cell lines.
- Variable growth characteristics and sensitivity with influenza strains in different cell lines.

Minimal Implementation Considerations:

- Utilize standard reference methods such as those described in the “WHO Manual for the Laboratory Diagnosis and Virological Surveillance of Influenza”¹⁰.
- Ensure sustainable expertise; each laboratory must have a training plan in place for virus isolation methods and troubleshooting.
- Establish a backup plan for times when commercially available cells are not available.
- Perform an accurate cost analysis, and identify sustainable sources of funding.

Antiviral Resistance Testing (pyrosequencing, neuraminidase inhibition):

Definitive antiviral resistance testing requires both phenotypic/functional resistance testing of the virus using a neuraminidase inhibition (NAI) assay, and detection of genetic markers of drug resistance by pyrosequencing and/or sequencing. Both of these test methods are performed at CDC.

Pyrosequencing is also performed at a subset of PHLs; this provides a cost-effective and efficient approach to expand antiviral resistance surveillance, providing data that can be used to inform patient management and treatment recommendations at the regional and national level.

Advantages of Maintaining or Implementing Pyrosequencing:

- Detects known genetic markers of resistance and rapidly detects cases of potential resistance.
- Provides more rapid turn-around-time than NAI.

Disadvantages of Pyrosequencing:

- Requires multiple assays to detect different genetic markers for each influenza subtype (eg H275Y is the marker for only A/H1pdm 2009).
- Requires supplemental testing by NAI for definitive confirmation of antiviral resistance.
- Requires specialized expertise to perform and interpret results.
- Requires specialized equipment and expensive reagents.
- Requires periodic revalidation of assays due to frequent genetic changes.

Advantages of Maintaining or Implementing NAI:

PHL and CDC:

- Provides definitive characterization of functional antiviral resistance testing.
- Detects resistance that is not caused by previously identified genetic markers of resistance.
- Detects resistance in a novel influenza virus that may be due to previously unrecognized genetic mutations.

¹⁰ Manual for the laboratory diagnosis and virological surveillance of influenza, WHO Global Influenza Surveillance Network http://whqlibdoc.who.int/publications/2011/9789241548090_eng.pdf

Disadvantages of NAI:

- Requires specialized expertise to perform and interpret results.
- Requires specialized equipment and expensive reagents.
- Requires antiviral drugs obtained from each drug manufacturer.

Minimal Implementation Considerations:

- Perform an accurate cost analysis, identify sustainable sources of funding.
- Laboratory must have the capability to:
 - Interpret and report test results appropriately for either virologic surveillance, or patient management.
 - Report all results to CDC in a timely manner for inclusion in national antiviral data.
 - Revalidate assays when mutations or changes to the protocols occur.
 - Perform virus isolation if using NAI assay.
 - Submit all influenza viruses positive for any resistance marker to CDC in a timely manner for confirmatory testing.

Respiratory Pathogen Assays

Molecular testing for other respiratory viruses, such as commercially available RVP assays or laboratory developed tests (LDTs), have become common in many clinical laboratories and are increasingly used in PHLs to provide jurisdictional and national information about other circulating respiratory pathogens. These data can aid in identifying other agents that cause flu-like community illnesses or outbreaks, and provide reassurance that surveillance is not “missing” influenza during periods when flu activity is low but other flu-like illnesses are prevalent.

Advantages of Maintaining or Implementing Molecular Respiratory Pathogen Assays:**PHL:**

- Provides data on circulating respiratory viruses that can be used for state, regional or national (NREVSS) surveillance or outbreak investigations.
- Provides a more rapid, sensitive and specific alternative to respiratory virus culture.
- Provides influenza surveillance partners with additional information about circulating viruses that may cause ILI, this may also help to incentivize partners to consistently contribute to flu surveillance.

Disadvantages of Respiratory Pathogen Assays:

- Expensive
- Variable sensitivity and specificity across pathogens and commercial assays.
- May lose sensitivity overtime, especially for influenza, due to changes in the viruses.

Minimal Implementation Considerations:

- Perform an accurate cost analysis, identify sustainable sources of funding.
- Determine how respiratory pathogen data will be used to supplement influenza surveillance data and/or diagnostic testing.

Influenza Hemagglutination

Influenza hemagglutination inhibition (HI, HAI) testing performed at CDC, using specialized antisera, remains a cornerstone of antigenic characterization to monitor changes in circulating influenza viruses and inform influenza vaccine virus selection. HAI tests using the WHO kit reagents that are provided by CDC to the PHLs can be used to detect and identify influenza viruses, but are not a reliable indicator of antigenic drift or strain changes.

Advantages for Maintaining or Implementing Influenza Hemagglutination:

PHL:

- o Identifies influenza type, influenza A subtype, and influenza B lineage using the Influenza WHO kit reagents provided by CDC.
- o Facilitates and helps to ensure more efficient detection of antigenic variants.

CDC:

- o Provides information on antigenic changes that may impact vaccine effectiveness, these data are critical to inform annual vaccine virus selection.

Disadvantages of Influenza Hemagglutination:

- o Requires significant expertise and specialized reagents (e.g., turkey red blood cells).
- o Requires virus culture.
- o Time consuming.
- o Influenza type and subtype can more rapidly be determined by rRT-PCR.

Minimal Implementation Considerations:

- o Laboratory must have the capability to:
 - Interpret and report test results appropriately for virologic surveillance.
 - Perform virus isolation.
 - Obtain red blood cells from appropriate species (e.g., turkey).
- o Perform an accurate cost analysis, identify sustainable sources of funding.

Influenza Serologic testing:

Serologic testing is neither a surveillance nor diagnostic testing tool, but is currently used primarily by CDC or academic institutions for research purposes or retrospective seroprevalence studies. Serologic testing to detect influenza virus antibodies may be performed using a variety of methods, including hemagglutination inhibition, enzyme-linked immunosorbent assay, and neutralization.

Advantages of Maintaining or Implementing Influenza Serologic Testing:

PHL and CDC:

- o Provides retrospective seroprevalence data.

Disadvantages of Influenza Serologic Testing:

- o Requires capability to implement assay as a laboratory-developed test, there are no FDA cleared methods for influenza serology.
- o Requires both a high level of expertise and specialized reagents.

- Patient serologic responses may be cross-reactive and difficult to interpret.

Minimal Implementation Considerations:

- Perform an accurate cost analysis, identify sustainable sources of funding.
- Maintain staff with capability to interpret serologic test results in the context of surveillance.

Direct Specimen Immunofluorescence:

Direct specimen immunofluorescence (DFA) testing methods involve testing clinical material taken directly from the patient. These cell preparations are tested using fluorescent-labeled antibodies, which are visible when examined using an immunofluorescence microscope. DFA is primarily a diagnostic tool.

Advantages of Maintaining and Implementing DFA:

- Provides the ability to produce results within an hour of specimen receipt.

Disadvantages of DFA:

- Requires specialized expertise and fluorescent microscopes.
- Results are subjective and dependent on the individual reader's expertise
- Not as accurate as rRT-PCR.
- Sensitivity and specificity of reagents may vary in response to current strains.

Minimal Implementation Considerations:

- Assess the necessity of maintaining this as a diagnostic tool.
- Perform an accurate cost analysis, identify sustainable sources of funding.

Rapid Influenza Diagnostic Tests (RIDTs):

Currently available RIDTs for the detection of influenza viruses employ a variety of methods, including enzyme-linked immunosorbent assays, immunochromatographic lateral flow immunoassays, and membrane-based immunoassays. In addition to differences in methodologies, these tests also have varying requirements for specimen collection and handling.

Advantages of Maintaining and Implementing RIDTs:

PHL

- Produce rapid results for patient care (often within fifteen to thirty minutes).
- Have widespread commercial availability.

Performance accuracy of RIDTs

RIDTs may be used to help with diagnostic and treatment decisions for patients in clinical settings, such as whether to prescribe antiviral medications. However, due to the [limited sensitivities and predictive values of RIDTs](#), negative results of RIDTs do not exclude influenza virus infection and influenza should still be considered in a patient if clinical suspicion is high based upon history, signs, symptoms and clinical examination.

Reported sensitivities of RIDTs range from 10-80% compared to viral culture or RT-PCR. Specificities of RIDTs are approximately 90-95% (range 85-100%). Thus false negative results occur more commonly than false positive results.

RT-PCR to confirm results of an RIDT are recommended when:

- A patient tests negative by RIDT when community influenza activity is high and laboratory confirmation of influenza is desired.
- A patient tests positive by RIDT and the community prevalence of influenza is low, and a false positive result is a consideration.
- A patient has had recent close exposure to pigs or poultry or other animals and novel influenza A virus infection is possible.

- Require minimal training, can be performed in non-laboratory settings.

Disadvantages of RIDTs:

- Significantly less sensitive and specific than rRT-PCR assays.
- Potentially less reliable when new virus strains emerge.

Minimal Implementation Considerations:

- Determine how respiratory pathogen data will be used to supplement influenza surveillance data and/or diagnostic testing.
- Establish policies for reflex/confirmatory testing.
- Perform an accurate cost analysis, identify sustainable sources of funding.

Considerations for Maintaining or Implementing Influenza Testing Methodologies

In addition to considering the advantages and disadvantages of the various test methods, below is a set of questions that can be used as a decision tool in deliberations among the laboratory director, epidemiologist/influenza surveillance coordinator, and clinical laboratory partners.

- Which influenza tests are most important to maintain?
- For what purpose is the influenza test(s) needed?
- How will the test results contribute to influenza virologic surveillance data within the state?
- How is the influenza test methodology in question currently funded? Is this funding mechanism sustainable?
- Is there an expectation, either through official policy or relations with the local clinical community, that the laboratory maintain capacity to perform some or all of the “additional” influenza tests?
 - If yes, what are the reasons for this expectation (e.g., is there a perceived need to maintain “traditional methods” in the PHL as the capability to perform culture and HAI declines in the clinical laboratory sector)?
- Is the necessary expertise available in the PHL and is it sustainable? Does your laboratory have access to and utilize appropriate consensus protocols and testing materials (e.g., MDCK cell lines, antiviral drugs, monoclonal fluorescent antibodies)?
- Does the laboratory have an experienced microbiologist/virologist that understands the basic virology, epidemiology, pathogenesis of influenza and other respiratory pathogens to enable test result and surveillance data interpretation, and surveillance capability decisions?
- If an influenza test method, such as culture, is eliminated, do decision makers understand that implementing testing on an as needed basis would be extremely difficult due to loss of expertise, length of time needed to acquire testing supplies, and time to perform CLIA required validation studies?
- Is the influenza methodology that is deemed necessary available from another source, such as CDC, a PHL shared services site, or a hospital/academic laboratory? What are the concerns with and road blocks to accessing alternate sources of testing?

- Do the resources (funding, staffing, supply costs) needed to maintain secondary testing adversely impact the capacity to perform rRT-PCR and test the number of specimens necessary for routine surveillance, or other essential laboratory functions?
- If the IRR was unable to continue providing either some (i.e. ancillary) or all of reagents needed for RT-PCR, how would this affect the decision to maintain “additional” testing methods?
- Does the laboratory meet the minimal considerations listed for each test method that will be implemented or maintained?

Data Management

Public Health Laboratory Data for Surveillance

The data generated through PHL reporting is published each week through CDC's FluView, www.cdc.gov/flu/weekly/. Therefore, timely and accurate reporting of laboratory surveillance data using electronic data systems is of the utmost importance. For public health laboratories, the main method of electronic reporting is through PHLIP. PHLIP achieves interoperability between different types of systems in order to exchange information in a useful and meaningful way. PHLIP utilizes an HL7 messaging standard to facilitate data exchange, allowing for harmonization of laboratory test results using standard vocabularies and terminology, like LOINC and SNOMED. PHLs can find information regarding implementation of HL7 messaging for CDC FLU rRT-PCR panels, including applicable LOINC test codes and SNOMED result codes at www.cdc.gov/flu/professionals/diagnosis/rtpcr-test-kits.htm.

Additional information about LOINC, SNOMED and HL7 can be found at:

- LOINC – www.loinc.org
- SNOMED – www.ihtsdo.org
- HL7 – www.hl7.org

The use of electronic data systems that provide data in real time and comply with national standards is a requirement to achieve right size virologic surveillance. The real time data elements (as found in the [HL7 PHLIP message guide](#)) that are available at each PHL may vary and some PHLs receive considerably more epidemiologic and specimen collection information than other PHLs. The minimum the PHLs should provide is:

- Specimen identifier and unique patient identifier
- State where specimen was collected
- Date of Birth of patient and/or Age plus unit (years, weeks, months, days)
- Specimen collection date
- Specimen Received Date
- Performed Test method
- Test result

The PHLs that already have PHLIP capability should consider reporting additional information. The additional information to include, if available:

- Vaccination status
- Antiviral treatment
- Travel information
- Patient death information
- Additional geographic information (ex. county, city, zip)
- Patient location at time of testing (inpatient, outpatient, long-term care facility)
- Whether specimen was related to an outbreak
- Whether specimen was sent to CDC and if so, ID included with CDC specimen
- Date of illness onset

APHL's Informatics program has a Technical Assistance Team available to assist public health laboratories with PHLIP implementation. The Technical Assistance Team provides tools and human resources to assist PHLs, public health agencies, and other data exchange partners in understanding, navigating and accomplishing the task of sending electronic data using simple, effective, standards-based methods. For more information about the Technical Assistant Team, visit www.aphl.org/aphlprograms/informatics/collaborations/phlip/Pages/default.aspx and refer to frequently asked questions about technical assistance teams.

Non-Public Health Laboratory Data for Surveillance

As resources for PHL testing shrink, the value of alternate data sources becomes increasingly important. State or local public health departments and laboratories are encouraged to explore options to collect and incorporate influenza testing data from non-public health laboratory sources. The supplemental data can help increase the confidence in the surveillance data within the state. This data could include rapid influenza diagnostic testing (RIDT) data and/or data from clinical/commercial laboratories within a jurisdiction. A number of other sources of virologic surveillance data may be available to augment both state and national surveillance. Data from these sites may be transmitted electronically as specimen level records or in aggregate by a simpler method. Regardless of whether specimen level or aggregate data is received, necessary data elements include:

- a. Date or week of specimen collection, receipt, or test
- b. Total number of tests performed and influenza positives by:
 - Type
 - Subtype (if available)
 - Age group

Potential Alternate Data Sources of local, state, and national data include:

- Clinical Sites including RIDT Sites and Clinical Laboratories – A number of influenza coordinators have initiated laboratory test reporting from selected facilities in their states. These data are used by the influenza coordinators to monitor influenza activity. Providing the supplemental surveillance data to clinicians is a useful resource to guide patient management decisions.
- Commercial Laboratories – A significant number of non-hospital laboratory testing is performed by a small number of commercial laboratories (e.g., Quest Diagnostics, LabCorp, ARUP Laboratories, and Mayo Medical Laboratories). Surveillance programs could obtain data from these laboratories on specimens tested from within their jurisdiction.
- Electronic Laboratory Reporting (ELR) for Meaningful Use – As part of the Affordable Care Act and related activities, inpatient and outpatient healthcare facilities have been provided monetary incentives to implement electronic health records. Along with these incentives, facilities are required to report notifiable diseases, syndromic surveillance, and vaccine registry data using automated electronic messaging standards (e.g., HL7, LOINC, SNOMED). For states that have required reporting of influenza laboratory results, health departments may be able to use these electronic messages to augment virologic surveillance.

Alternate data sources for CDC to supplement national surveillance data:

- Commercial Laboratories – A significant number of non-hospital laboratory tests are performed by a small number of commercial laboratories (e.g., Quest Diagnostics, LabCorp, ARUP Laboratories, and Mayo Medical Laboratories). These laboratories provide a significant amount of influenza testing for hospitals, and physician offices around the country.
- Federally Qualified Health Centers (FQHCs) –As of 2010, over 1100 of these clinics operate under the supervision of the Health Resources and Services Administration (HRSA) to provide care to the medically underserved¹¹. With the Affordable Care Act (ACA), the numbers and patient volumes are anticipated to increase.
- Department of Veterans' Affairs (VA) – There are 1,766 VA facilities¹², although influenza laboratory testing is only available at a subset of these.
- Department of Defense (DoD) – Data available from DoD includes virologic data collected among military personnel around the globe and laboratory test information from military and non-military facilities that care for military dependents, notably through TriCare Insurance.
- Centers for Medicare and Medicaid Services (CMS) – Since 2009, data on influenza testing has been made available on a more real-time basis utilizing the CMS reimbursement exchange data repository.
- Vendors of Electronic Health Records (EHR) and Laboratory Information Management Systems (LIMS) – Vendors and large users of EHRs and LIMS may be a source of influenza testing information. Examples of these include GE Healthcare, Cerner, Sunquest, HCA, and others.
- Diagnostic Device Manufacturers – An increasing trend in diagnostic testing is the use of mobile communications from test devices to cloud-based web services to allow ease of access of information to patients, doctors, and insurers. This capability also allows automated messaging of de-identified results from influenza test devices to a cloud where public health entities can access the information for monitoring influenza in their jurisdictions.

¹¹ <http://bphc.hrsa.gov/ftca/riskmanagement/healthcenterqualityimprovement.pdf>

¹² <http://www2.va.gov/directory/guide/home.asp?isflash=1>

Partnerships & Communication

It is important that states establish and maintain partnerships and networks among PHLs, clinicians, state epidemiologist/influenza surveillance coordinator, clinical laboratories, RIDT sites, CDC and manufacturers. Many states already have existing partnership and communication networks for both influenza surveillance and other activities such as [Laboratory Response Network \(LRN\)](#) and APHL's [Lab System Improvement Program \(L-SIP\)](#) (APHL, 2013; CDC, 2012). Improvements to influenza surveillance can be made by leveraging existing partnerships and communication networks for influenza surveillance, LRN, and other laboratory-based surveillance activities. For example contact databases that already exist for LRN can be enhanced to include laboratories that perform influenza testing without creating an entire new system. Many states have established courier services to transport specimens for LRN, newborn screening other programs. These may be leveraged to improve access to specimens for influenza surveillance. Additional examples related to key partnerships, provided through stakeholder input and pilot site activities are described below.

The most important partnership for effective virologic surveillance is the relationship between the **public health laboratory and epidemiology/Influenza Coordinators**. Examples of ways to optimize epi-lab partnerships:

- Conduct regular in-person epidemiology-laboratory meetings to establish seasonal virologic surveillance strategies, determine appropriate sample sizes, allocate funds, and regularly assess the effectiveness of the surveillance system.
- Collaborate in grant writing, monitor grant activities, identify and address problems and gaps, and coordinate outbreak response.
- Establish consensus protocols for sharing influenza testing data. Examples include releasing laboratory data to a secure portal that epidemiologists can access, or by providing epidemiologists access to selected views in the laboratory databases.

Collaboration between Epidemiologists and Laboratorians at the Washington State Department of Health

Every August, epidemiologists and laboratorians at the Washington State Department of Health (DOH) meet in person to discuss virologic surveillance plans for the upcoming influenza season. They discuss criteria for influenza testing at the Public Health Laboratories (PHL), plans for engaging sentinel providers and laboratories, and changes to specimen submission instructions. Written instructions for submitting specimens to PHL for influenza testing are revised collaboratively.

Healthcare providers and local health jurisdiction staff who want to submit a non-routine specimens for influenza testing, including specimens from patients with suspected novel influenza and those in outbreaks, are asked to contact a DOH influenza epidemiologist prior to submission. The epidemiologist reviews the request and informs the laboratory staff about the estimated arrival time and priority status of the specimens. If critical specimens do not arrive at PHL by the expected time, a laboratorian will contact the submitter to determine the whereabouts of the specimens. This system ensures that epi and lab partners have access to timely information regarding status of high priority specimens.

PHL-Epidemiology-Clinician-Academic Partnerships

- Provide strategic communication from state epidemiologists and PHLs to clinicians and clinical laboratories as needed to increase awareness when targeted surveillance is needed to identify emerging viruses or characterize outbreaks.
- Provide education programs to clinicians, especially on the utility of rapid point of care tests, and provide training on specimen collection, handling and transport to ILINet and other surveillance specimen submitters. Training may be achieved through on-site presentations, teleconferences, and mailings.
- Collaborate with clinicians and academic researchers on studies to increase understanding of influenza.

Value of Epidemiology, Public Health Laboratory and Clinician Partnerships

Successful influenza virologic surveillance programs are not built overnight and cannot be sustained without proper care. Strong relationships between state epidemiology, public health laboratory, and clinical partners are crucial to ensuring quality and consistent data and specimens for influenza virologic surveillance. Clinicians with a keen interest in public health who can help grow and foster surveillance efforts in the community and among their colleagues can be an enormous asset. Establishment of a strong network of providers who will submit timely and quality specimens requires dedicated resources that can provide encouragement, feedback, and guidance. When dedicated resources routinely work with submitters on appropriate reporting, specimen collection and submission, specimens are more likely to be of higher quality and improve the virus detection abilities at PHLs. Virologic surveillance efforts cannot be a one way street. Giving back to providers who participate serves as a reminder of the importance of their contributions. Providing incentives can be as simple as ensuring timely feedback of results and findings or as complex as offering additional testing of negative sample for other respiratory pathogens.

Source: Unpublished communications with Influenza Incidence Surveillance Project (IISP) participants

PHL-Clinical Laboratory/Testing Site –Influenza Surveillance Coordinator Partnerships

Strong partnerships and communications with clinical laboratories and influenza testing sites are important to obtaining quality and consistent data and specimens. Listed below are some examples for improving these relationships.

- Access alternate data sources to supplement influenza surveillance, as described in the [Data Management Section](#). Commercial, web-based survey instruments are available at little to no cost (e.g. SurveyMonkey™, SurveyGizmo) to collect testing data from partners. Some of the tools can provide participants with an identifying login and be pre-filled with participant information to ease the burden and increase participation rates. The data collected can include information on a variety of agents and test methods. Data can be downloaded to a spreadsheet for analysis. These reports provide data from thousands of tests performed by these

laboratories and test sites throughout the influenza season. In addition, a number of clinical laboratories perform their influenza testing as part of a respiratory virus molecular panel, access to this data allows for a more complete picture of circulating respiratory pathogens.

- Establish collaborative relationships with specimen providers to ensure and/or improve the quality and consistency of specimen submissions as outlined in the Sampling Section. For more information on implementation for establishing specimen provider networks, please reference the Sampling Implementation Guidance.
 - In addition to the standard communication methods, some states distribute a “handbook” containing instructions, forms, and summary data of laboratory surveillance needs in their jurisdiction. The *WI Laboratory-Based Surveillance Plan* is one such example (www.slh.wisc.edu/dotAsset/27271.pdf).
- Promote the value of participating in the surveillance system, provide incentives when permissible. Incentives do not need to be monetary; they can be test kits, training, certificates of appreciation, attendance at state conferences, and reference books, as well as the “added value” of improved surveillance data that can be used to improve clinical management recommendations.
- Provide specimen collection and shipping supplies and courier service to virologic surveillance participants. Most health care providers and clinical laboratories will not be able to absorb the cost of surveillance supplies or shipping.
- Provide timely updates to specimen providers via email, web or fax. Clinical laboratory partners and PHLs both benefit from exchanges of information related to current influenza activity, commercial test shortages, and emerging disease threats (e.g., 2013 novel coronavirus and H7N9).
- Provide workshops, teleconferences/webinars, and educational materials related to influenza surveillance, proper specimen collection, and use of RIDT.
- Provide proficiency assessment challenges or exercises related to influenza testing and specimen packaging if budget permits. These exercises may be coordinated with other state/local preparedness activities.

PHL and CDC Partnerships

Collaboration between the PHLs and CDC’s Influenza Division is imperative for effective virologic surveillance. The CDC, especially the Virus Surveillance and Diagnosis Branch, provide vital support to the PHLs, and relies on data and specimens submitted by the PHLs. Coordination between the CDC and PHLs is often facilitated by APhL.

- CDC provides didactic and hands-on training to PHLs in test methods, and national teleconferences to share surveillance guidance for laboratory testing and influenza coordinator activities.
- CDC provides technical support to PHLs, including assistance with assay troubleshooting and interpreting unusual results.

- CDC provides testing reagents and materials through the Influenza Reagent Resource (IRR), along with updates to the assay and implementation support when novel viruses emerge (e.g., H3N2v, H7N9).
- CDC partners with PHLs to complete the necessary validation studies for regulatory approval of new assays.
- CDC provides guidance to assist states develop emergency outbreak and pandemic response plans, and provides essential support in actual response situations.

Considerations for Building Effective Partnerships and Communications

1. Has your influenza surveillance program and PHL identified the appropriate contacts among public health, clinician, and laboratory partners within your jurisdiction?
2. If so, do you routinely and collaboratively review the list of contacts to ensure that all key partners are included (e.g. identify new partners, update for staffing changes, etc).
3. Does your laboratory maintain a database of current contact information and influenza testing capabilities for identified laboratories within your jurisdiction?
 - a. Note: Some SPHLs maintain multiple separate databases – one for the LRN, one for a statewide laboratory network for surveillance, etc., while some SPHLs have added influenza testing sites and included influenza testing capability, surveillance participation, etc. to the existing LRN database.
4. Does your influenza surveillance program and PHL designate one or more staff members to coordinate outreach activities (e.g., network or surveillance coordinator/manager/advisor)?
5. Do you maintain a communication plan that identifies and links system partners?
6. Do you collaborate with other laboratories and rapid influenza testing sites to acquire virologic testing result data and specimens for further virologic testing?
7. Do you have a method to collect influenza testing data from clinical laboratories/testing sites (e.g. survey tool, fax, or web portal)?
8. Does your laboratory and/or influenza surveillance program maintain the capability to exchange information and data via email, fax or other electronic tools with laboratories within your jurisdiction?
9. Do you provide reports on the current status of circulating influenza types and subtypes via website, newsletter or other means?
10. Do you provide an end of year summary to all stakeholders about the influenza season? Do you provide individualized reports to participating laboratories and/or providers?
11. Do you relay the importance of receiving specimens for confirmatory testing, subtyping, and identification of unsubtypeable specimens to your clinical partners, especially when the threat of a novel virus is high (e.g., H3N2v, H7N9)?
12. Does your influenza surveillance program and/or laboratory provide teleconferences, webinars, or in-person training and outreach to clinician and laboratory surveillance partners and potential partners?

Is there a mechanism for feedback and corrective action to providers who incorrectly or inappropriately send specimens to the PHL (e.g. improperly shipped; incorrect form, incomplete information, sent dry swab not in media)?

Quality Management System

National Surveillance System Quality Monitoring

The ultimate value of virologic surveillance data is dependent on the quality of specimens, laboratory procedures, and data analysis. CDC and state/local jurisdictions should establish performance metrics and monitor essential components of the national influenza virologic surveillance system to ensure quality and make improvements as needed. Listed in this section are key components that should be routinely assessed, but it should be noted that each quality management system will vary and jurisdictions need not be limited by this list. It is likely that existing data sources can be leveraged to assess the quality of many surveillance components.

State/Local Quality Management Responsibilities:

At the state and local level, quality management systems need to monitor both internal performance and performance in meeting national surveillance requirements, including those defined in this document. As discussed previously, influenza virologic surveillance systems are complex and vary across jurisdictions; quality management systems will likewise need to be tailored to each system. Regardless of the assessment mechanism(s), it is recommended that states have some method to evaluate the following elements related to influenza virologic surveillance, and make adjustments and improvements as needed.

- Compliance with ELC, PHEP, and other cooperative agreement and grant benchmarks for all epidemiology and laboratory components of the surveillance system.
- Specimen submissions through the provider networks including consistency, quality and number. Timely electronic transmission of specimen-level data. The Public Health Laboratory Interoperability Project (PHLIP) system is the preferred method of reporting.
- Percentage of influenza test results received by CDC from the PHL within two weeks of the test date.
- Capability to provide year-round molecular

New Hampshire Public Health Laboratories Influenza Quality Monitoring, 2013

When it became clear that the 2012-2013 influenza season was ramping up to be the busiest since the 2009 pandemic, staff at the NH Public Health Laboratories realized they needed to closely monitor influenza submissions, in order to ensure resources were appropriately allocated to meet the goals of the surveillance program. This was achieved by building a simple “Daily Flu Data” report in the laboratory LIMS, which effectively extracted all data associated with influenza specimens received from the start of the season in October 2012.

The flu report was run daily, and data was dumped into an Excel spreadsheet. Once populated in the spreadsheet, the data could be manipulated in a number of ways: specimens received by date, specimens received by provider, specimens received by county, etc. This manipulation allowed us to quickly see if submissions were increasing or decreasing and if we were obtaining representative samples from across the state. We were able to use this tool to reach out to health care providers and encourage additional submissions from providers in underrepresented areas of the state, while informing others that their submissions had exceeded our needs. By storing the spreadsheet in a shared folder on the network, any staff who needed to use the information was able to access it quickly and conveniently.

testing for the detection, typing and subtyping of seasonal influenza viruses and detection of novel influenza viruses.

- Systematic submission of representative influenza positive clinical materials and/or viral isolates for national virologic in accordance with annual CDC specimen submission guidance.
- Rapid referrals of all unsubtypable influenza A viruses to CDC.
- Proficiency in PCR methods for influenza virus detection, typing, and subtyping. The laboratory must operate in compliance with the Clinical Laboratory Improvement Act (CLIA) 88' requirements, which include participating in an external/blinded proficiency test for each assay. CDC provides a quality assessment (QA) panel to PHLs at least one time per year which helps PHLs fulfill this CLIA requirement. Participation in this CDC assessment also provides data that helps CDC assess and address training needs.
- Usage of IRR-provided reagents, materials and other resources used for national surveillance in comparison to the number of specimens tested and reported to CDC. IRR reagents are provided to PHL's to support testing for national surveillance. Prior authorization from CDC is needed if IRR-provided materials are needed to support special studies.
- Staff expertise to perform each influenza test method used at the PHL. Every PHL should have a competency assurance policy that addresses initial training, assay update training, and cross-training to ensure continuity of operations in a surge event such as the 2009 H1N1 pandemic.
- Staff expertise and ability to adopt influenza assay revisions, add additional testing markers or adopt assay interpretation updates. The detection of novel or variant viruses may result in new assay components or modified interpretation guidelines.
- Maintenance of an influenza specimen repository that can be utilized for assay verification and validation and competency testing as needed. Store a subset of positive and negative specimens containing a mix of influenza types and subtypes at -70°C.

CDC Quality Management Responsibilities:

- Cross-reference public health laboratory influenza testing data reported to CDC against virologic specimen submissions to CDC and CDC-designated laboratories.
- Monitor national surveillance data for timeliness, adequate testing and specimen submissions numbers, and representativeness to ensure the system is able to effectively inform situational awareness and vaccine virus selection efforts. When needed, provide targeted communications to PHLs that are not consistently complying with specimen submission expectations or to request additional specimens as needed. Targeted communications help reduce confusion about specimen requirements, and focus attention on key gaps or special needs.
- Monitor IRR reagent ordering history in relation to testing reported to CDC. Targeted follow-up to PHLs can be an effective method for addressing excessive reagent ordering which may be due to oversampling or unrecognized technical problems. When technical problems are identified, CDC and the PHL should collaborate to implement appropriate solutions as needed.

Considerations for Establishing and Maintaining Quality Management Systems:

1. Does your laboratory and surveillance program have mechanisms in place to monitor compliance with grant/cooperative agreement benchmarks and deliverables?
 - a. Example: Leadership should meet regularly to review grant line items, identify issues, and document progress. LIMS and tracking spreadsheets can be used to document and verify deliverables are being met.
2. Does your laboratory and/or surveillance program have processes for monitoring the quality, quantity, consistency, representativeness and timeliness of specimen submissions from specimen providers?
 - a. Example: Influenza Coordinator and PHL may regularly review specimen submission data for quality indicators such as number of specimens rejected for poor quality, number of inconclusive test results, etc.
 - b. Example: Influenza Coordinator and PHL may regularly review number of specimens received compared to number designated by sample size calculators. Sampling may be adjusted as appropriate.
3. Does your laboratory have mechanisms in place to ensure that representative specimens are being submitted to CDC or CDC-designated laboratories in accordance with annual specimen submission guidance and other CDC requests?
 - a. Example: Use LIMS and tracking spreadsheets to monitor the timeliness of influenza surveillance testing and submissions to CDC or CDC-designated laboratories. Regularly check to ensure specimens submitted to CDC are representative of the influenza activity in your jurisdiction (see examples in [Sampling Implementation](#)) and current CDC guidelines. Verify that shipment quantities and frequencies are in compliance with the CDC guidelines.

Surge Capacity for Influenza Surveillance, Novel Event Investigation and Outbreak Events

The virologic surveillance system should be flexible and scalable for rapid, effective response to support initial diagnostic needs and case counts in novel event investigations, and enhanced surveillance outbreak and pandemic scenarios. Pre-event and during an event, communication and coordination between epidemiology and laboratory leadership will be essential to develop, refine and change the strategy for virologic surge sampling and testing.

Pre-event:

- Ensure that PHL representatives are included in state preparedness and pandemic planning activities. Address the role and resource needs of the PHL in state/jurisdictional pandemic plans. Pandemic Planning Information can be found at :
www.cdc.gov/phpr/coopagreement.htm, www.cdc.gov/flu/pandemic-resources/tools/index.htm.
- Utilize the APHL “Emerging Infectious Disease Planning and Response Framework” to identify key partners and preparedness activities, including validation of new testing methodologies, biosafety, regulatory requirements, training, information dissemination, specimen collection and transport guidance.
www.aphl.org/aphlprograms/infectious/emerging/Documents/EID_2005April_Emerging-Infectious-Diseases-Planning-and-Response-Framework.pdf
- Develop and maintain a laboratory pandemic surge plan that is integrated into a laboratory wide Continuity of Operations Plans (COOP). The surge plan should address:
 - communication/coordination with epidemiologists for specimen triage,
 - algorithm changes to improve efficiency and throughput, or to meet specific surveillance needs,
 - resources (e.g., staff, cross-training, equipment, space, reagents and consumable supplies),
 - biosafety considerations for working with novel viruses,
 - options to mitigate the capacity gaps and bottlenecks identified in the state Lab Capacity Modeling report provided to participating states by APHL/CDC.
- Establish mechanisms to determine and implement a sampling strategy for investigation following detection of a novel event. Consider the potential scenarios that may define sampling approaches, such as the need to identify additional cases and detect person-to-person transmission. Consider targeted surveillance options including clinical severity criteria, exposure risk, number of hospitalized cases/deaths, and other event specific needs.
- Establish criteria for specimen triage, and decision points for performing diagnostic testing, and/or expanding virologic surveillance testing. Draft scenario specific scale up and ramp down criteria that can be quickly applied when a novel event or outbreak occurs.
- Define laboratory testing algorithms that may be implemented to accommodate the influx of surveillance and diagnostic specimens.

- Periodically assess laboratory contingency and crisis surge capacity, as defined in Surge Requirements Section. Laboratory Capacity modeling has been conducted in most state PHLs using a model developed by APHL and CDC (ref). These models estimated baseline capacity, identified likely sources of bottlenecks in a surge event, and evaluated the impact of various changes on overall throughput. Utilizing a surge algorithm with surge resources (staff, equipment, etc.) that are expected to be available to the laboratory during emergency periods of high testing demand provided a capacity increase of 127% compared to the Influenza A/B Typing Assay with reflex Influenza A Subtyping algorithm and baseline resources. The implementation of a super surge process strategy, which included changing from the Influenza A/B Typing with full Influenza A Subtyping Panel baseline to an Influenza A/B Typing only testing algorithm, along with the addition of staff and equipment, could increase national aggregate PHL daily capacity from approximately 5,000 specimens to approximately 14,250 specimens – an estimated change of 185%.
- Utilize sample size calculators to estimate number of samples to be collected and tested for various novel event investigation scenarios. Compare laboratory surge capacity to likely sample size expectations so that both epidemiology and laboratory leaders understand capacity gaps, if any. Collaboratively explore strategies to reduce sample size or increase capacity.

Event:

- Use sample size calculators to determine the appropriate sample size for the investigation, based on the scenario, acceptable Confidence Level and Error rate. Sustaining testing to provide daily case counts will not be possible and states should consider use of sample size calculators to adjust testing volumes as necessary to answer key surveillance objectives as the event evolves.
- Develop, refine and change state/local needs and/or CDC guidance as needed, based on the specific event:
 - Defined surveillance/investigation objectives
 - Targeted sampling approaches
 - Initial virus detection reporting criteria (laboratory to epidemiology)
 - Ramp up/ Ramp down criteria
- Revise testing algorithms to improve efficiency and throughput or meet specific surveillance needs.

Detailed guidance on pandemic response is beyond the scope of this document. During a large scale event, CDC, CSTE, and APHL will coordinate to provide timely direction and support. It is important that information disseminated by CDC, state health officials, and APHL to PH Laboratory Directors is disseminated to the laboratory staff. Management and technical staff should participate in CDC/APHL conference calls to obtain pertinent recommendations.

Financial Resources

Every state will need to determine how to achieve influenza surveillance goals to meet national and state needs. Federal resources (funding, reagents) distributed to states need to be directed principally to activities that support overall national priorities. State/local capabilities beyond those recommended as essential to meet national virologic surveillance goals will require securing sustainable state or jurisdictional funding. When optimizing services and justifying budget requests, PHLs and surveillance programs should work cooperatively to address:

- Essential elements defined in this roadmap for national surveillance.
- Options for shared services among PHLs.
- Scalability of the surveillance system based on available resources.
- State/local specific influenza surveillance expectations or operational issues.

Cost Accounting

Optimizing resources and justifying funding requests will require better cost accounting at the national, state and local level. Results from the 2011 Right Size Influenza Virologic Surveillance Landscape survey showed that 35% of state PHLs (16 out of 45 SPHLs) were unable to provide accurate estimates of influenza testing costs or were only able to provide rough estimates. There are many advantages to effective cost accounting including, but not limited to:

- Identify true cost of virologic surveillance
- Plan and allocate resources for each influenza season
- Justify surveillance program and laboratory testing budgets
- Assess which surveillance components are covered by various funding sources (e.g. federal vs state funds)
- Calculate the cost the PHL absorbs beyond the actual state or federal funds provided
- Ensure PHLs and programs are good stewards of existing resources
- Determine and justify the most efficient testing algorithm for various scenarios (see [Laboratory Testing Section](#) for additional information on testing algorithms)
- Write grant proposals
- Characterize impacts of funding reductions

The cost of performing influenza surveillance testing varies across jurisdictions. While there is no standard method that can be applied across all jurisdictions to assess costs, at a minimum a cost analyses should include four areas related to influenza surveillance¹³: labor, consumable materials, equipment, and overhead/shipping.

- **Labor** – Including laboratory, epidemiology/influenza coordinator, and information technology staff salaries, fringe/benefits costs, and capabilities.

¹³ APHL internal report on influenza cost factors, unpublished data.

- **Consumable Materials** – Including material costs for specimen collection materials (if provided to the specimen submitters by the surveillance program or laboratory), submitter incentives (if provided), reagents and testing kits used for extraction and rRT-PCR processes, as well as consumables both directly and indirectly associated with PCR testing. If the laboratory is performing any additional tests as defined in the Laboratory Section, costs per test should be determined for these consumables as well.
- **Equipment** – Including acquisition, service/maintenance, and depreciation costs for all equipment used for influenza testing.
- **Overhead & Miscellaneous** – Including costs associated with facilities, surcharges, utilities, sentinel provider networks (e.g. provider communication tools), information technology support, transportation of specimens to and from the laboratory, training, and travel.

It may also be helpful to reference CMS Medicare/Medicaid CPT codes and fee schedules when performing cost analysis. The federal standards for clinical diagnostic reimbursement for testing can help estimate laboratory costs for surveillance testing as well as serve as a comparator to true cost accounting. The CMS fee schedule will vary across states, and PHLs should reference the most recent CPT fee schedule for their state. Listed below are some of the relevant CPT codes for influenza surveillance testing that may assist with determining laboratory costs.

- CDC Flu rRT-PCR Dx Panel: 87501 x 2 if assay uses separate wells for Flu A and Flu B.
 - If a laboratory uses a commercial multiplex test, reference the 87502 code x 1 for up to two analytes.
- Influenza subtyping: 87503 x number of analytes (e.g., influenza A(H1N1), influenza A(H3N2) and influenza A(H1N1)pdm).
- Virus culture: 87252.
- Virus culture by shell vial: 87254
- Immunofluorescent identification (if culture positive) and hemadsorption: 87253

Allocating Available Funds

These questions address suggested processes for cost analysis and coordination needed to optimize funding allocation among those involved in influenza surveillance within the state.

1. Do you have a routine process for having all involved parties meet to discuss grant development, planning, fund allocation, and deliverable/benchmark monitoring?
2. Do you have a process for determining how much it costs your jurisdiction to perform influenza virologic surveillance?
 - a. Example: Perform a detailed cost analysis for both surveillance program and laboratory components. See the cost accounting section above for some helpful tips.
3. Do you have a method or process for allocating funds across program and laboratory elements?

- a. Example: Appropriate representatives of the laboratory and surveillance program meet at the beginning of each season and periodically throughout season to discuss allocation of funds and monitor expenditures throughout the season.
4. Are ELC, PHEP, and quality management benchmarks considered in prioritization of funding allocation?

Table 1 is provided as a tool to facilitate funding allocation discussions and to help identify potential funding gaps. Use of this table, or a similar state- developed tool, can help elucidate the actual costs of influenza surveillance and provide a basis for discussion and priority setting. This table can be modified to fit your jurisdiction's funding sources and surveillance components. Depending on the level of detail desired, you can complete this table by listing dollar amounts, percentages or simply using checkmarks to indicate which surveillance components are funded by each of these sources in your jurisdiction.

Table 1. Resource Allocation Tracking Table

	ELC	PHEP	State	Other: _____
Surveillance coordinator				
Epidemiology staff				
Laboratory staff				
Laboratory testing reagents, supplies (not provided by CDC)				
LIMS/electronic reporting/IT support				
Specimen collection supplies (e.g. VTM, swabs)				
Specimen transport (e.g., shipping boxes, courier or commercial carrier costs)				
Sentinel provider incentives				
Equipment & equipment maintenance costs, including service contracts				
Laboratory Overhead (e.g. travel/training, autoclave/waste, printing/publications/education/press releases)				
Supporting local capacity (e.g. local PHLs and programs)				
Other: _____				
Other: _____				

Resource Justification

As previously mentioned, state/local capabilities beyond those recommended as essential to meet national virologic surveillance goals will require securing alternate, supplemental, sustainable state or local funds. As federal and states funds decline in this post-pandemic period, it may be necessary to explore options for alternate non-traditional funding sources, such as research grants, or academic partnerships for special studies. Justifying resource needs requires an accurate estimate of surveillance system costs and funding needs/gaps.

Funding Fact Sheet Tool for States

Fact sheets and success/impact stories are useful tools when requesting funds and resources to meet jurisdictional surveillance requirements. Appendix D provides a funding justification “fact sheet” template that can be modified and used as a tool by public health laboratory leaders to highlight a specific jurisdiction’s program impact and funding needs for influenza virologic surveillance. To see other example fact sheets, please visit www.aphl.org/policy/facts/Pages/default.aspx.

Intended Use: Public health laboratory leaders can customize this document using the editable version of Appendix D located at www.aphl.org/xxxx. This is intended to be used to highlight surveillance program successes, impact, and funding needs to non-public health audiences such as state and local health officers, policy makers and other government officials.

Instructions for Use: To create a jurisdiction-specific fact sheet, go to www.aphl.org/xxxx for an editable version of Appendix D that leadership can modify to highlight their own program’s success and funding needs. For example, the current template highlights the influenza A(H3N2)v outbreak and the PHLs’ response to the outbreak. If a state was impacted by influenza A(H3N2)v, the template anecdote may only need minor changes; however if a state has a better example of how their program has impacted their state and/or national surveillance it may be necessary to adapt or revise this section.

Users will notice that the fact sheet uses basic, non-scientific language; the level of technical detail included should be tailored to the target audience. In the example language in Appendix D, the target audience would be a lay person that has no prior knowledge of influenza testing, surveillance, or public health laboratories. For example, the term such as “influenza” is replaced with “flu,” a widely recognized, colloquial reference.


The template also includes a box to highlight funding needs. It is recommended that this box only include the funding needs being requested of the specific target audience. Some jurisdictions may prefer to use this fact sheet solely to promote impact and success, in which case the funding box can be deleted. Lastly, keep the fact sheet focused on a specific topic and/or request. The recommended maximum length is one page, front and back.

Appendices

Appendices

- A. Surveillance Sampling Process Map
- B. Sample Size Calculator Quick Glance Tables
- C. Laboratory methods
- D. Funding Fact Sheet Template
- E. Glossary of Terms and Acronyms
- F. Additional Resources and References

Appendix A: Surveillance Sampling Process Map



Appendix B: Sample Size Calculator Instructions for Use

Sample Size Calculator Instructions for Use:

User guide will be added.

Situational Awareness

	Average Population	Confidence Level (5% Margin of Error)		
		95	90	85
Less than 2 Million	1,094,706	118	87	68
2-5 Million	3,530,463	132	94	73
5-10 Million	7,193,033	135	96	74
10-20 Million	15,214,189	137	97	74

Novel Event Detection at national detection thresholds using 100% unscreened MA-ILI specimens

	Average Population	Confidence Level (5% Margin of Error)		
		95 (1/200)	95 (1/165)	90 (1/165)
Less than 2 Million	1,094,706	21	17	13
2-5 Million	3,530,463	68	55	43
5-10 Million	7,193,033	137	112	86
10-20 Million	15,214,189	287	237	182

Novel Event Detection at national thresholds, assuming 10% Flu+/MA-ILI, using approx. 60% MA-ILI specimens and 40% specimens pre-screened Flu +

	Average Population	Confidence Level (5% Margin of Error)					
		95 (1/200)		95 (1/165)		90 (1/165)	
		~60% MA-ILI	~40% Flu +	~60% MA-ILI	~40% Flu +	~60% MA-ILI	~40% Flu +
Less than 2 Million	1,094,706	13	1	6	2	4	1
2-5 Million	3,530,463	41	3	15	4	13	3
5-10 Million	7,193,033	82	6	22	9	16	7
10-20 Million	15,214,189	47	24	27	21	22	16

Appendix C: Laboratory methods

Requirements for Influenza Detection and Identification Methods

The below chart provides several influenza testing methodologies, described in the [Laboratory Testing Implementation](#) section. The chart provides only high level equipment, and material and reagent needs, more detailed needs should be obtained from the standard operating procedure being implemented within your laboratory.

	rRT-PCR	Culture	Shell Vial	Direct Specimen Immunofluorescence	RIDT	Antiviral Resistance	Neuraminidase Inhibition	Hemagglutination Inhibition	Serology
Equipment									
Biological safety cabinet	X	X	X	X	X	X	X	X	DEPENDENT ON SPECIFIC PROCEDURE
Centrifuge with carriers and containment	X		X						
Centrifuge, cytospin				X					
Cold block	X								
Microcentrifuge, benchtop	X				X	X			
Microscope, light		X							
Microscope, fluorescence			X	X					
Heating block						X			
Incubator		X	X	X			X		
Real Time PCR instrument	X					X			
Gel documentation system						X			
PyroMark™ vacuum prep tool and workstation						X			
Roller apparatus		X							
Splash shield with goggles					X				
Mirror stand to read plate reactions								X	
Liquid dispensers and diluters	X	X	X	X		X	X	X	
Pyrosequencing instrument						X			
Thermocycler-conventional						X			
Transilluminator						X			
Victor X4 or Equivalent Fluorescent Reader							X	X	
CDC JASPR Software (IC50)							X	X	

Requirements for Influenza Detection and Identification Methods (continued)

	rRT-PCR	Culture	Shell Vial	Direct Specimen Immunofluorescence	RIDT	Antiviral Resistance	Neuraminidase Inhibition	Hemagglutination Inhibition	Serology
Materials & Reagents									
Miscellaneous laboratory glassware	X	X	X	X		X	X	X	
NA Drugs (e.g. oseltamivir)							X		
NA Sensitive & NA Resistant Influenza Virus Controls						X	X		
Opaque flat bottom 96well plates							X		
12 column & single reservoirs							X		
Cell Culture Flasks		X	X						
Cell culture sensitive to influenza viruses		X	X						
Cell culture medium		X	X						
Ethanol, 100%/molecular grade	X					X	X		
Microcentrifuge tubes, DNase/RNase free	X					X			
Molecular grade water	X					X	X		
Monoclonal antibodies to influenza A and B viruses			X	X					
Fluorescent conjugate (if MAbs not conjugated)			X	X					
Fixative (e.g., acetone) to prepare ("fix") cells			X	X					
Containers to rinse stained slides				X					
Incubation chamber/container				X					
Influenza primers and probes	X					X			
Influenza positive and human specimen controls	X								
Neuraminidase inhibition drug							X		
Neuraminidase assay kit							X		
Nucleic acid extraction/isolation kits, multiple	X					X			
rRT-PCR mastermix	X					X			
Pyrosequencing Reaction mix						X			
Red blood cell solution, standardized		X						X	
Phosphate buffered saline (PBS)		X	X	X			X	X	
Pyrosequencing instrument						X			
Transilluminator						X			
U-bottom microtiter plates								X	
Plate cover tapes								X	
Antisera against influenza A and B, treated								X	
Expertise Level	★★ ★★	★★ ★★	★★ ★	★★ ★★	★	★★ ★★	★★ ★★	★★ ★	★★ ★★

Appendix D: Funding Fact Sheet Template

PREVENT FLU ILLNESS AND DEATHS

<JURISDICTION NAME> PUBLIC HEALTH ACTIVITIES PREVENT FLU

Every year over <3,000> people are hospitalized, around <200> deaths occur in <state>, more than <20,000> doctors room and emergency room visits are made, and more than <500,000> work and school days are lost because of influenza (respiratory flu) even with everything we are already doing to prevent flu illnesses. We can reduce the number of illnesses and deaths from flu with more effective vaccines, maintaining critical laboratory testing and improving the quality and use of public health data about flu. The work of our state public health laboratory and CDC contributes to the development of new vaccines as flu viruses change each year. We could improve our public health flu prevention and detection efforts by optimizing our state's ability to receive and test samples in the public health laboratory and share the information with state and federal health officials. At a minimum, we need to maintain our state's current ability to monitor respiratory flu in our community. Improvements in the quality of reliable public health data can contribute to healthcare cost savings through reduced antibiotic usage, reduced disease transmission, and reduced hospitalizations.

<JURISDICTION>'S FUNDING NEEDS FOR EFFECTIVE FLU DETECTION AND PREVENTION

Governmental public health laboratories and health departments play a crucial role in flu detection, characterization, prevention and pandemic preparedness. Advances in testing and a highly-trained public health laboratory workforce allowed for a successful response to the 2009 flu pandemic and enabled the early detection and investigation of a

number of new potential pandemic strains since then. However, declining federal and state funding to support public health laboratory testing, coupled with increasing shortages of public health laboratory staff with necessary expertise, threaten

the success of current programs, and their ability to address ongoing pandemic threats. An effective flu monitoring system that meets both state and federal public health needs requires funding to:

STATE FUNDING

FY2013 <\$50k> (Enacted)

FY2014 <\$85k> (State Required Amount)

- Improve and maintain <Jurisdiction>'s ability to collect, test, and interpret a sufficient number of flu samples to allow state and federal health officials to make informed decisions.
- Enhance laboratory identification of new flu viruses to support policy decisions such as school closures to help control the spread of disease and to support patient treatment decisions.
- Support efforts to ensure sufficient number of flu samples are received by the governmental public health laboratory to provide real-time details on flu activity in our state.
- Enhance public health outreach, training and coordination with health care providers, hospitals and private sector laboratories.
- Support development and maintenance of electronic reporting of laboratory results allowing public health official and health care providers to respond to test results rapidly.


- Expand the ability to test samples by increasing and cross-training public health laboratory staff to maintain routine flu monitoring and surge capacity for another pandemic.
- <ADD/REMOVE/MODIFY FUNDING NEEDS STATEMENTS AS APPROPRIATE FOR YOUR JURISDICTION>.

STATE IMPACT – Detecting a New Flu Virus

During the summers of 2011 and 2012, a new flu virus*, not covered by current vaccines, was detected, mostly in people who came in contact with sick pigs at agricultural fairs. Due to diligent work by the health department and public health laboratory, we were able to limit the spread of this virus to people and allow the public to enjoy the fairs. The public health laboratory played an integral role in detecting this new flu virus and promptly shared this information with state and national health officials so preventative measures could be implemented. Our public health laboratory shared samples containing the new virus with CDC to help national flu experts prepare for and respond to this potential new threat. <CAN BE REPLACED WITH JURISDICTION-SPECIFIC STORY>

CONTACT

For more information, contact <NAME, <TITLE>, <PHONE NUMBER, <EMAIL>



During the 2009 flu pandemic, the laboratory test being performed at public health laboratories was “the *sine qua non* of every action in public health regarding the virus, from closing a neighborhood school to shutting down international flights. Effects near and far rippled from what the test could tell about the virus.”

--Thomas R. Frieden, MD, MPH, Director, CDC
adapted from *Lessons from a Virus*

**To learn more about the influenza A(H3N2) variant virus referenced here, please visit:*
www.cdc.gov/flu/swineflu/h3n2v-cases.htm

Appendix E: Glossary of Terms and Acronyms

The following terms and acronyms are defined here according to their usage in this document; terms may have additional meanings beyond these descriptions.

Term/Acronym	Definition (as used in this document)
APHL	Association of Public Health Laboratories; the national nonprofit organization that represents governmental public health; www.aphl.org
CDC	Centers for Disease Control and Prevention; Federal organization within the U.S. Department of Health and Human Services, to protect health and promote quality of life through the prevention and control of disease, injury, and disability.; www.cdc.gov
CLIA '88	Clinical Laboratory Improvement Amendments of 1988; the Clinical Laboratory Improvement Amendments (CLIA) were passed by the U.S. Congress in 1988 to establish quality standards for all laboratory testing to ensure the accuracy, reliability and timeliness of patient test results regardless of where the test was performed; www.cdc.gov/clia/
CSTE	Council of State and Territorial Epidemiologists; an organization of member states and territories representing public health epidemiologists; www.cste.org
DOT	Department of Transportation; U.S. government agency with responsibilities that include regulating the transport of dangerous or hazardous materials
Drifted viruses	Small changes in the virus that happen continually over time. Antigenic drift is a mechanism for variation in viruses that involves the accumulation of mutations within the genes that code for antibody-binding sites, which reduces or inhibits the binding of neutralizing antibodies.
EIP	Emerging Infections Program; a program administered by the CDC in which a network of 10 state health departments and their partners conduct specialized surveillance, prevention, and control of emerging infectious diseases.
ELC	Epidemiology and Laboratory Capacity cooperative agreements (grants) to states.
ELR	Electronic Laboratory Reporting; the electronic transmission to public health of laboratory reports which identify reportable conditions
ELR for Meaningful Use	Activities overseen by Centers for Medicare & Medicaid Services in support of interoperable electronic health records, including electronic laboratory reporting, which can be used to achieve measurable outcomes.
FLuView	CDC website, provides data and analysis of current influenza activity; www.cdc.gov/flu/weekly/summary.htm
HL7	Health Level 7; standards developed by a non-profit, ANSI-accredited organization that provides standards for the exchange, integration, sharing, and retrieval of electronic health information; www.hl7.org/
HPAI	Highly pathogenic avian influenza; influenza viruses that can cause disease in chickens when they are infected, but does not relate to disease-causing capabilities related to other species
IATA	International Air Transport Association; responsibilities include regulating the air transport of dangerous or hazardous materials; www.iata.org

ILI	Influenza-Like Illness; defined as fever (temperature of 100°F [37.8°C] or greater) and cough and/or sore throat; used as a measure of illness that may be caused by influenza viruses
ILINet	U.S. Outpatient Influenza-like Illness Surveillance Network (ILINet) ; healthcare providers in all states, the District of Columbia and the U.S. Virgin Islands who report to CDC the total number of patients seen and the number of those patients with influenza-like illness (ILI) by age group
IRR	Influenza Reagent Resource; organization established by the U.S. CDC to provide registered users with reagents, tools and information to study and detect influenza virus; http://influenzareagentresource.org
LIMS	Laboratory Information Management System; also known as a Laboratory Information System (LIS), a software system to support laboratory operations, possibly including data tracking and exchange, sample tracking, and informatics.
LOINC	Logical Observation Identifiers Names and Codes; a universal code system to allow the exchange and aggregation of electronic health data from many independent systems; http://loinc.org
LRN	Laboratory Response Network; a national network of more than 150 local, state and federal public health, food testing, veterinary diagnostic, and environmental testing laboratories to respond to public health emergencies. http://emergency.cdc.gov/lrn/
MDCK	Madin-Darby Canine Kidney; a cell culture line used primarily for culture of influenza viruses
MMWR	Morbidity and Mortality Weekly Report; often called “the voice of CDC,” the weekly publication provides timely, reliable, authoritative, accurate, objective, and useful public health information and recommendations; www.cdc.gov/mmwr
Neuraminidase Inhibition	Preventing the normal function of a protein present in influenza viruses (neuraminidase) that allows the virus to be released from infected cells; also a method to determine one component of the subtype of an influenza virus.
NLS	National Laboratory System; a proposed nationwide laboratory system to provide communication, coordination and testing capacity to effectively detect and report outbreaks and exposures; www.cdc.gov/mlp/nls.aspx
NLTN	National Laboratory Training Network; a joint program of APHL and the CDC to develop and deliver education programs for professionals in both public and private sector laboratories; www.aphl.org/training/nltm/pages/default.aspx
Novel influenza virus	Reassortant or animal origin virus found in humans or previously unidentified antigenic virus subtype.
NREVSS	National Respiratory and Enteric Virus Surveillance System; a laboratory-based system managed by CDC to monitor patterns in the detection of respiratory syncytial virus (RSV), human parainfluenza viruses, adenoviruses and rotavirus; www.cdc.gov/surveillance/nrevss/
PCR	Polymerase Chain Reaction; method to detect DNA.
Phenotypic	The observable characteristics, e.g., morphology, behavior, biochemical properties, of an organism, resulting from the expression of the organism's genes and environmental factors, as compared to genotypic, which refers to the genetic makeup of the organism.

PHEP	Public Health Emergency Preparedness; program of cooperative agreements established by the U.S. government to provide funding, guidance, and technical assistance to state and local public health departments to improve the readiness to respond to a variety of threats, including infectious diseases, natural disasters, biological or chemical events, etc.
PHL	Public Health Laboratory; laboratory that performs testing and other laboratory services on behalf of the entire jurisdiction as part of the public health system to protect the public against diseases and other health hazards.
PHLIP	Public Health Laboratory Interoperability Project; a collaboration between CDC and APHL to establish reliable laboratory data exchange between state public health laboratories and the CDC; www.aphl.org/aphlprograms/informatics/collaborations/phlip/pages/default.aspx
Pyrosequencing	Method of determining the order of nucleotides in DNA; can be used in determining the presence of selected mutations that denote resistance of influenza viruses to antiviral medications.
RIDT	Rapid influenza diagnostic test; Tests to detect influenza viruses that are commercially available and can be completed within 15-30 minutes.
rRT-PCR	Real-time reverse transcription polymerase chain reaction; method to detect RNA by creating complementary DNA.
SNOMED	Systematized Nomenclature of Medicine; collection of medical terminology that includes codes for diseases, procedures, microorganisms, etc., to allow consistent indexing, storage, retrieval, and aggregation of clinical data across medical records.
SPHL	State Public Health Laboratory; laboratory that performs testing and other laboratory services on behalf of the entire jurisdiction, where the jurisdiction is the state, part of the public health system protecting the public against diseases and other health hazards.
Subtype	Groupings within the influenza A viruses, denoted by the hemagglutinin (“H”) and neuraminidase (“N”) of the virus, e.g., influenza A (H1N1) or influenza A (H3N2).
Threshold	The level or point at which an event occurs or decision is made, the “cut-off point”.
Unsubtypable	Any influenza positive specimen that cannot be definitively typed and subtyped as a circulating seasonal influenza virus. Influenza positive specimens producing non-standard or inconclusive results as defined in the CDC Human Influenza Virus Real-Time RT-PCR Diagnostic Panel Instructions for Use package insert.
USPS	United States Postal Service; responsibilities include regulating the transport of dangerous or hazardous materials in the U.S. Mail; www.usps.com/
Warm Base	Minimum state of readiness. In this context, applies to laboratory capacity.
WHO Collaborating Laboratory	Laboratory testing site that participates in the World Health Organization global influenza surveillance system, providing data and samples to the CDC, a WHO Collaborating Center for Influenza.

Appendix F: Additional Resources and References

APHL's Emerging Infectious Diseases Planning and Response Framework

A checklist to be used by public health laboratory leaders and scientists that outlines the various elements public health laboratories must address with each disease outbreak or emerging threat. www.aphl.org/aphlprograms/infectious/emerging/Documents/EID_2005April_Emerging-Infectious-Diseases-Planning-and-Response-Framework.pdf

CDC's FluView

FluView provides regular summary reports of influenza activity including interactive maps, viral surveillance data, antigenic characterization, antiviral resistance, novel influenza activity, and many other valuable data summaries.

www.cdc.gov/flu/weekly/fluactivitysurv.htm

CDC's Seasonal Influenza Website

CDC's website provides up to date information on current seasonal influenza activity as well as critical updates regarding any emerging viruses or issues.

www.cdc.gov/flu/

Clinical Description & Lab Diagnosis of Influenza,

www.cdc.gov/flu/professionals/diagnosis/index.htm

Guidance for Clinicians on the Use of Rapid Influenza Diagnostic Tests,

www.cdc.gov/flu/professionals/diagnosis/clinician_guidance_ridt.htm

Rapid Diagnostic Testing for Influenza,

www.cdc.gov/flu/professionals/diagnosis/rapidlab.htm

Google Flu Trends, www.google.org/flutrends/

Joint Commission: Strategies for Improving Rapid Influenza Testing in Ambulatory Settings (SIRAS), www.jointcommission.org/siras.aspx

Interim Guidance on Use of Intervals, Triggers, and Actions for Novel Influenza A (H1N1) Response

www.flu.gov/planning-preparedness/federal/operationalplans.html# (Appendix A)

Laboratory Efficiencies Initiative (LEI) Resources and Tools

This website provides resources and tools for the LEI have been designed to address the needs of public health laboratories. Current available resources assist with services changes, legal considerations for sharing services and third-party billing, such as "A Practical Guide to Assessing and Planning Implementation of Public Health Laboratory Service Changes."

www.aphl.org/aphlprograms/Iss/Laboratory-Efficiencies-Initiative/Pages/Resources-and-Tools.aspx

Laboratory System Improvement Program (L-SIP)

APHL's Laboratory System Improvement Program (L-SIP) advances the efficacy of state and local public health laboratory systems through a guided process of performance evaluation, system improvements, and periodic evaluation and reassessment. Participating member laboratories receive resources and technical assistance to guide them on their way to system excellence.

www.aphl.org/aphlprograms/lss/performance/pages/default.aspx

Public Health Emergency Preparedness Cooperative Agreements (PHEP)

CDC's PHEP website provides valuable resources including cooperative agreement guidance, training videos, and CDC's "National Standards for Public Health Preparedness Capabilities" document.

www.cdc.gov/phpr/coopagreement.htm

WHO Interim Global Surveillance Standards for Influenza

"This document proposes surveillance objectives and describes global standards for a minimal basic respiratory disease surveillance system for the monitoring of influenza."

www.who.int/influenza/resources/documents/INFSURVMANUAL.pdf

WHO Global Influenza Surveillance Network – Manual for the laboratory diagnosis and virological surveillance of influenza

"WHO has therefore developed this manual in order to strengthen the laboratory diagnosis and virological surveillance of influenza infection by providing standard methods for the collection, detection, isolation and characterization of viruses."

http://whqlibdoc.who.int/publications/2011/9789241548090_eng.pdf

References

- APHL. (2007). Retrieved from Definition of a State Public Health Laboratory System:
http://www.aphl.org/AboutAPHL/publications/Documents/Definition_of_a_State_PHL_System_2007.pdf
- APHL. (2010). *The Core Functions of State Public Health Laboratories*. Retrieved from
http://www.aphl.org/aboutaphl/publications/documents/com_2010_corefunctionsphls.pdf
- APHL. (2011). *Lessons from a Virus*.
- APHL. (2012). *The Influenza Virologic Surveillance Landscape of State and Local Public Health Laboratories*. Retrieved from
http://www.aphl.org/AboutAPHL/publications/Documents/ID_2012June_The-Influenza-Virologic-Surveillance-Landscape-of-State-and-Local-Public-Health-Laboratories.pdf
- APHL. (2013). *Laboratory Efficiencies Initiative*. Retrieved from
<http://www.aphl.org/aphlprograms/lss/Laboratory-Efficiencies-Initiative/Pages/default.aspx>
- APHL. (2013). *Laboratory System Improvement Program*. Retrieved from
<http://www.aphl.org/aphlprograms/lss/performance/pages/default.aspx>
- CDC. (2011, March). Retrieved from National Standards for State and Local Planning:
http://www.cdc.gov/phpr/capabilities/DSLR_capabilities_July.pdf
- CDC. (2012, July 2). *Emergency Preparedness and Response: The Laboratory Response Network Partners in Preparedness*. Retrieved from <http://www.bt.cdc.gov/lrn/>
- CDC. (2013, April 2). *Office of Public Health Preparedness and Response: Funding and Guidance for State and Local Health Departments*. Retrieved from <http://www.cdc.gov/phpr/coopagreement.htm>
- CDC & CSTE. (2008, February). *Competencies for Applied Epidemiologists in Governmental Public Health Agencies (AECs)*. Retrieved from
http://www.cdc.gov/appliedepicompetencies/downloads/Applied_Epi_Comps.pdf
- Institute of Medicine (US) Forum on Medical and Public Health Preparedness for Catastrophic Events. (2010). *The Communications Challenge: Definitions, Standards, and Metrics. Medical Surge Capacity: Workshop Summary*. Washington (DC): National Academies Press (US). Retrieved from
<http://www.ncbi.nlm.nih.gov/books/NBK32856/>